

ASPIRIN- aspirin tablet, film coated
Chain Drug Marketing Association, Inc.

Quality Choice 44-157 RESERVED 83324-066

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments?

1-800-426-9391

Principal display panel

**QC[®]
QUALITY
CHOICE**

NDC 83324-066-01

†Compare to the
Active Ingredient in
Genuine Bayer[®] Aspirin

Aspirin
(NSAID)

Pain Reliever
Fever Reducer 325 mg

Coated Tablets

100 Tablets

Actual Size


**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

† This product is not manufactured
or distributed by Bayer AG, owner of
the registered trademark Genuine
Bayer[®] Aspirin.

50844 REV0122D15712

Distributed by CDMA, Inc.
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362

NDC 83324-066-01



†Compare to the Active Ingredient in Genuine Bayer® Aspirin


Aspirin (NSAID)

Pain Reliever
Fever Reducer **325 mg**

Coated Tablets


100 Tablets

Actual Size



Distributed by CDHA, Inc.
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin.
50844 REV0122D15712



6 35515 90431 6

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

No Print/No Varnish Lot & EXP

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 325 mg (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves: ■ minor pain of arthritis
■ headache ■ muscle pain ■ menstrual pain
■ toothache ■ pain and fever of colds

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ facial swelling
■ shock ■ hives ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older
■ take a blood thinning (anticoagulant) or steroid drug
■ have 3 or more alcoholic drinks every day while using this product ■ have had stomach ulcers or bleeding problems
■ take more or for a longer time than directed
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued)

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma
■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs.
Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ ringing in the ears or a loss of hearing occurs
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur ■ redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
■ children under 12 years: do not use unless directed by a doctor

Other information
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Inactive ingredients corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments? 1-800-426-9391

--- STOP PEELING ---

Adhesive Only

Quality Choice 44-157

ASPIRIN

aspirin tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83324-066
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY35J)	

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASPIRIN;44;157
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83324-066-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/03/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/03/2024	

Labeler - Chain Drug Marketing Association, Inc. (011920774)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(83324-066)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(83324-066)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(83324-066)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(83324-066)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(83324-066)

