

AOBBIY TATTOO NUMBING CREAM- tattoo numbing cream cream
Shenzhen Situya Trading Co., Ltd.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredient

LIDOCAINE 4%

Purpose

Local Anesthetic

Use

temporarily relieves local and anorectal discomfort associated with the anorectal disorder or swelling.

Warnings

For external use only.

Do not use

pregnant or breastfeeding, ask your doctor before use.
in case of accidental overdose, seek immediate medical help or contact the Poison Control Center.
seal is broken or missing.

When Using

do not exceed the recommended daily dosage unless directed by a doctor. Certain sensitive individuals may have an allergic reaction to some of the ingredients of this product.
do not put this cream into the rectum using your fingers, applicators, or any medical devices.

Stop Use

the symptom being treated does not improve, or if redness, irritation, swelling, pain, or other symptoms appear or worsen. Seek medical help right away.

Keep Out Of Reach Of Children

In case of accidental ingestion, seek medical help right away.

Directions

Apply up to 3 times a day.

Children (under 12 years of age): consult a doctor.

Adults: before applying this product, clean the affected area with mild soap and warm water and rinse thoroughly. Pat dry the area with a soft cloth or tissue.

Other information

Keep away from direct sunlight or heat.

Store at room temperature 59°-86°F (15°-30°C).

Inactive ingredients

Aqua, Propylene Glycol, Carbomer 934, Lecithin (Soybean), Tocopheryl Acetate, Benzyl Alcohol.

Correct Use Steps Before Tattooing


1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
2. Apply a thick layer of the numbing cream onto the treatment area.
3. Wrap tightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
4. Wipe away the residue with a soft cloth or tissue.
5. Begin the procedure.

Notice


Due to the difference in temperature and skin type, the numbness effect duration will differ. Testing in a small area first is recommended to ensure the best numbness effect.

PRINCIPAL DISPLAY PANEL

60 mm*60 mm* 58 cm



AOBBiY



AOBBiY


Tattoo

NUMBING CREAM

4% Lidocaine with Arnica, Eucalyptus oil and Emu Oil

6 Hours Maximum Strength Pain Relief

Net Wt: 1.7 oz/50ml



DRUG FACTS

Active Ingredient	Purpose
Lidocaine 4%	Local Anesthetic

Uses: temporarily relieves local and anorectal discomfort associated with the anorectal disorder or swelling.

Warning (For external use only)

Do not use this product if

- you are allergic to Lidocaine.
- in case of accidental overdose, seek immediate medical help or contact the Poison Control Center.
- seal is broken or missing.

When using this product

- do not exceed the recommended daily dosage unless directed by a doctor.
- Certain sensitive individuals may have an allergic reaction to some of the ingredients of this product.
- do not put this cream into the rectum using your fingers, applicators, or any medical devices.

Stop use and ask a doctor if the symptom being treated does not improve, or if redness, irritation, swelling, pain, or other symptoms appear or worsen. Seek medical help right away.




Correct Use Steps Before Tattooing:

1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
2. Apply a thick layer of the numbing cream onto the treatment area.
3. Wrap tightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
4. Wipe away the residue with a soft cloth or tissue.
5. Begin the procedure.

Notice: Due to the difference in temperature and skin type, the numbness effect duration will differ. Testing in a small area first is recommended to ensure the best numbness effect.

DISTRIBUTED BY:
AOBBiY LLC
Sheridan, WY 82801



DRUG FACTS (Continued)

KEEP OUT OF REACH OF CHILDREN.
In case of accidental ingestion, seek medical help right away.

Directions


- Apply up to 3 times a day
- Children (under 12 years of age): consult a doctor.
- Adults: before applying this product, clean the affected area with mild soap and warm water and rinse thoroughly. Pat dry the area with a soft cloth or tissue.

Other Information


- Keep away from direct sunlight or heat.
- Store at room temperature 59°-86°F (15°-30°C).

Inactive Ingredients

Aqua, Propylene Glycol, Carbomer 934, Lecithin (Soybean), Tocopheryl Acetate, Benzyl Alcohol.



AOBBiY



KEEP OUT OF REACH OF CHILDREN.
In case of accidental ingestion, seek medical help right away.

DIRECTIONS

1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
2. Apply a thick layer of the numbing cream onto the treatment area.
3. Wrap tightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
4. Wipe away the residue with a soft cloth or tissue.
5. Begin the procedure.

DISTRIBUTED BY:
AOBBiY LLC, Sheridan, WY 82801



AOBBiY

NUMBING CREAM

4% Lidocaine with Arnica, Eucalyptus oil and Emu Oil

6 Hours Maximum Strength Pain Relief

Net Wt: 30ml



DRUG FACTS

Active Ingredient	Purpose
Lidocaine 4%	Local Anesthetic


Uses: temporarily relieves local and anorectal discomfort associated with the anorectal disorder or swelling.

Warning (For external use only)

Do not use this product if you are allergic to lidocaine.

When using this product

- do not exceed the recommended daily dosage unless directed by a doctor. Certain sensitive individuals may have an allergic reaction to some of the ingredients of this product.
- do not put this cream into the rectum using your fingers, applicators, or any medical devices.



KEEP OUT OF REACH OF CHILDREN.
In case of accidental ingestion, seek medical help right away.

DIRECTIONS

1. Clean your hands and the treatment area with soft soap and warm water, rinse thoroughly, then gently pat dry.
2. Apply a thick layer of the numbing cream onto the treatment area.
3. Wrap tightly with plastic wrap and wait 45-60 minutes for the numbness to peak. (The peak lasts up to 6 hours.)
4. Wipe away the residue with a soft cloth or tissue.
5. Begin the procedure.

DISTRIBUTED BY:
AOBBiY LLC, Sheridan, WY 82801



AOBBiY

NUMBING CREAM

4% Lidocaine with Arnica, Eucalyptus oil and Emu Oil

6 Hours Maximum Strength Pain Relief

Net Wt: 100ml



DRUG FACTS

Active Ingredient	Purpose
Lidocaine 4%	Local Anesthetic

Uses: temporarily relieves local and anorectal discomfort associated with the anorectal disorder or swelling.

Warning (For external use only)

Do not use this product if you are allergic to lidocaine.

When using this product

- do not exceed the recommended daily dosage unless directed by a doctor. Certain sensitive individuals may have an allergic reaction to some of the ingredients of this product.
- do not put this cream into the rectum using your fingers, applicators, or any medical devices.

AOBBiY TATTOO NUMBING CREAM

tattoo numbing cream cream

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:82739-016

Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 mL	
Inactive Ingredients				
	Ingredient Name	Strength		
	CARBOMER 934 (UNII: Z135WT9208)			
	.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)			
	BENZYL ALCOHOL (UNII: LKG8494WBH)			
	LECITHIN, SOYBEAN (UNII: 1DI56QDM62)			
	WATER (UNII: 059QF0KO0R)			
	PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82739-016-01	30 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/15/2023	
2	NDC:82739-016-02	50 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/15/2023	
3	NDC:82739-016-03	75 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/15/2023	
4	NDC:82739-016-04	100 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/15/2023	
5	NDC:82739-016-05	150 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/15/2023	
Marketing Information				
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	OTC monograph not final	part348	09/15/2023	

Labeler - Shenzhen Situya Trading Co., Ltd. (706154255)

Establishment

Name	Address	ID/FEI	Business Operations
Shenzhen Situya Trading Co., Ltd.		706154255	manufacture(82739-016) , label(82739-016)