# MAXIMUM STRENGTH NIGHTTIME COLD AND FLU- acetaminophen, diphenhydramine hcl,phenylephrine hcl liquid THE KROGER CO

-----

## Maximum Strength Nighttime Cold and Flu 6 fl oz. (180 mL)

## **Drug Facts**

Active ingredients (in each 20 mL)	Purposes	
Acetaminophen 650 mg	Pain reliever/fever reducer	
Diphenhydramine HCl 25 mg	Antihistamine/cough suppressant	
Phenylephrine HCl 10 mg	Nasal decongestant	

#### Uses

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - runny nose
  - sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

## **Warnings**

## **Liver Warning:**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions . Symptoms may include:

- Skin reddening
- Blisters
- rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:**if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

cough that occurs with too much phlegm (mucus)

### Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

## When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks

be careful when driving a motor vehicle or operating machinery

## Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

Keep out of reach of children.

## Overdose warning:

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

#### **Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL= milliliter
- adults and children 12 years of age and older:20 mL in dosing cup provided every 4 hours
- children under 12 years of age:do not use

#### Other information

- each 20 mL contains:sodium 6 mg
- low sodium
- store at room temperature
- do not refrigerate
- dosing cup provided

## Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, natural and artificial flavor, potassium citrate ,propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum.

#### Questions or comments?

1-888-287-1915

#### PRINCIPAL DISPLAY PANEL -

NDC# 30142-736-06

Compare to Mucinex  $^{\rm ®}$  Fast-Max  $^{\rm ®}$  Maximum Strength Might Time Cold & Flu Active Ingredients

Maximum Strength

## Night TimeCold & Flu

- AcetaminophenPain Reliever/Fever Reducer
   Diphenhydramine HCl Antihistamine/Cough Suppressant
   Phenylephrine HCl Nasal Decongestant
- Relieves:

- Sore Throat; Itchy Throat-Cough
- Nasal Congestion Sneezing & Runny Nose
- Headache Body Pain

#### For Ages 12+

6 FL OZ (180 mL)

\*This product is not manufactured or distributed by Reckitt Benckiser, the distributor of Mucinex ® Fast- Max ® Maximum Strength Nighttime Cold & Flu

Tamper evident: do not use if printed seal under cap is broken or missing.

‡Maximum Strength per 4 hour dose.

## Distributed by:













#### MAXIMUM STRENGTH NIGHTTIME COLD AND FLU

acetaminophen, diphenhydramine hcl, phenylephrine hcl liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-736
Route of Administration	ORAL		

l	Active Ingredient/Active Moiety			
ı	Ingredient Name	<b>Basis of Strength</b>	Strength	
	ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 20 mL	

DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
EDETATE DISODIUM (UNII: 7FLD91C86K)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
POTASSIUM CITRATE (UNII: EE90ONI6FF)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
PROPYL GALLATE (UNII: 8D4SNN7V92)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SORBITOL (UNII: 506T60A25R)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
XANTHAN GUM (UNII: TTV12P4NEE)			

Product Characteristics		
Color	blue	Score
Shape		Size
Flavor	FRUIT	Imprint Code
Contains		

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:30142-736- 06	180 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/04/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/04/2020	

## Labeler - THE KROGER CO (006999528)

Revised: 7/2024 THE KROGER CO