

MEDI-FIRST PROTECTIVE COATING BANDAGE- benzethonium chloride and benzocaine aerosol, spray
Unifirst First Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Medi-First Bandage Spray

Drug Facts

Active ingredients

Benzethonium Chloride 0.2% w/w

Benzocaine 3.2% w/w

Purpose

Topical antiseptic

Topical anesthetic

Uses

☐ First aid to help protect against infection in

- ☐ minor cuts and scrapes
- burns

Warnings

☐ For external use only

Flammable

- keep away from fire or flame
-
- contents under pressure
-
- do not puncture or incinerate container
-
- do not expose to temperatures above 120° F

Do not use

- in or near eyes or other mucus membranes
-
- in case of serious burns
-
- in case of deep or puncture wounds
-
- for a prolonged period of time

-
- on a large portion of the body

Stop use and ask a doctor if

- condition worsens or symptoms persist for more than 7 days
-
- condition clears up and recurs within a few days
-
- redness, swelling, or irritation occurs

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- clean the affected area
-
- shake can well before using
-
- hold 4-6 inches from surface and spray area until wet
-
- may be covered with sterile bandage. If bandaged, let dry first
-
- for adult institutional use only
-
- not intended for use on children

Other information

- avoid inhaling
-
- use only as directed
-
- intentional misuse by deliberately concentrating and inhaling contents may be harmful or fatal

Inactive ingredients

Ethyl alcohol, isobutane, n-butane, propane, PVP-VA copolymer

Questions or comments?

1-800-634-7680

Medi-First Bandage Spray Label

Medi-First®

Protective Coating

Bandage Spray

- Provides a protective coating for minor burns, cuts, skin irritation, and minor insect bites
- Water soluble

Aerosol

WARNING:FLAMMABLE

Contents Under Pressure, Read Back Panel Carefully

Net Weight 3 OZ (85g)

47.0297

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- clean the affected area
- shake can well before using
- hold 4-6 inches from surface and spray area until wet
- may be covered with a sterile bandage. If bandaged, let dry first
- for adult institutional use only
- not intended for use on children

Other information

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal

Inactive ingredients

Ethyl alcohol, isobutane, n-butane, propane, PVP-VA copolymer

Questions or comments? 1-800-634-7680

Reorder #45017

Manufactured for Medique Products
Fort Myers, FL 33967 USA



MEDI-FIRST®

Protective Coating
Bandage Spray

- Provides a protective coating for minor burns, cuts, skin irritation, and minor insect bites
- Water soluble

Aerosol

WARNING: FLAMMABLE

CONTENTS UNDER PRESSURE, READ BACK PANEL CAREFULLY

NET WEIGHT 3 OZ (85g)

Drug Facts

Active ingredients

Benzethonium Chloride 0.2% w/w..... Topical antiseptic
Benzocaine 3.2% w/w Topical anesthetic

Purposes

Uses

First aid to help protect against infection in

- minor cuts and scrapes
- burns

Warnings

For external use only

Flammable

- keep away from fire or flame
- contents under pressure
- do not puncture or incinerate container
- do not expose to temperatures above 120° F

Do not use

- in or near eyes or other mucus membranes
- in case of serious burns
- in case of deep or puncture wounds
- for a prolonged period of time
- on a large portion of the body

Stop use and ask a doctor if

- condition worsens or symptoms persist for more than 7 days
- condition clears up and recurs within a few days
- redness, swelling, or irritation occurs

MEDI-FIRST PROTECTIVE COATING BANDAGE

benzethonium chloride and benzocaine aerosol, spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-450(NDC:52982-105)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	170 mg in 85 g
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	2720 mg in 85 g

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ISOBUTANE (UNII: BXR49TP611)	

BUTANE (UNII: 6LV4FOR43R)	
PROPANE (UNII: T75W9911L6)	
COPOVIDONE K25-31 (UNII: D9C330MD8B)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-450-17	85 g in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	04/05/2013	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	04/05/2013	

Labeler - Unifirst First Aid Corporation (832947092)

Registrant - Dixon Investments, Inc. (115315822)

Revised: 7/2019

Unifirst First Aid Corporation