HEARTBURN RELIEF- famotidine tablet, film coated Meijer Distribution Inc

Meijer Distribution, Inc. Heartburn Relief Drug Facts

Active ingredient (in each tablet)

Famotidine 10 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness**, **sweating**, **or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, hypromellose, iron oxide red, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, talc, titanium dioxide, triacetin

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Pepcid® AC active ingredient

ORIGINAL STRENGTH

heartburn relief

Famotidine Tablets, 10 mg

Acid Reducer

SEE NEW WARNINGS

Just One Tablet! Prevents & Relieves Heartburn Due to Acid Indigestion

30 Tablets

actual size

by Johnson & Johnson, owner of the registered *This product is not manufactured or distributed

Questions or comments? 1-800-7 19-9260

Drug Facts (continued)



DIST

CBY MEJER DISTRIBUTION GRAND RAPIDS, MI 49544 www.meijer.com

롱

preg elati niz ed starch, tal c, titan ium dio xi de, triaceti n microcrystalline cell ulose, poly dextrose, p d yethylene gly col, in n oxide red, ir on oxid e y ello w, ma gnes iu m stearate, In act ive ingredients camauba wax, hypromellose,

anutaiom mon't baton q ■ (7.77-%3) プ 2S-°OS its anota∎ ■ Neep the carton. It con tains important in formation. ■read the direction sandwamings before use not farm of ni 19A1V

- ■children un der 12 yeans:aak a doctor
- don ot use m ore than 2 tablets in 2 4 hours eating food or drinking beverages in atcause hear toum
- of water at any time from 1 5 to 60 minutes before assige with viblest twellow tablet with a glass of water. Do not chew.
- to relieve symptoms, swallow 1 tablet with aglass ■squits and children 1 Zyearsan dover

Directions

away. (1-800-222-1222) get m edical help or contact a Poison Con trol Center ing ht Keepouto freach of children. In case of overdose,

If pregnant or bre ast-fe eding, ask a health professional

■you need to take this product for more than 14 days ■ your near cournings or worsens Stopuse and as ka doctorif

with certain prescrip tion drugs. taking a prescription drug. Acid reduce rs may in teract Ask a doctor or pharma cist before useif you are

Drug Facts (continued)

- stom ach pain
 kidn ey disease

tra demark Pepcid[®].

- mu dhead Atiw yhe lucithea, gai saath with headb um
 - m frequent che st pain
- sweating; pain spreading to arme, nedcorshoulders; drest pain or should erp ain with shortness of breath;
- 🔳 hear fourn with lighthead edness, sweating, or dizzin ea m ore serious condition.
- had heartb um over3 mon ths. This may be a sign of a Ask a doctor before useif you have
 - with other acid reducers
- with blood, or bloody or black stools. These may be m if you h ave trouble or pains want and boot g niw olls want and both ave it uoy it eu tono (

or other acid reducers

Alergyalert. Do not use if you are all ergic to farm obdine **SgrinnsW**

certain food and beverages

- and sours tomach brought on byea ting ord in inking prevents heartburn associa ted with acid in dig estion so nus foursich
- relieve sheartburn associated with acid indigestion and

Famotidine 10 mg......Prid reducer Active ingredient (in each tablet) Purpose

Drug Facts



CONVENIENT RECLOSING TAB

Tips for Managing Heartburn

- · Do not lie flat or bend over after eating
- . Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
 Raise the head of your bed · Avoid heartburn-causing foods such as rich, spicy, fatty
- or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables . Eat slowly and avoid big meals
- · If overweight, lose weight · Quit smoking

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.



SEE NEW WARNINGS NDC 41250-493-65

Compare to Pepcid® AC active ingredient*



ORIGINAL STRENGTH

heartburn relief

Famotidine Tablets, 10 mg **Acid Reducer**

Just One Tablet! Prevents & Relieves **Heartburn Due to Acid Indigestion**

30 Tablets



HEARTBURN RELIEF

famotidine tablet, film coated

Product Information	1
----------------------------	---

Product TypeHUMAN OTC DRUGItem Code (Source)NDC:41250-493

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8) FAMOTIDINE 10 mg

Inactive Ingredients

Ingredient Name
Strength

CARNAUBA WAX (UNII: R12CBM0 EIZ)

HYPROMELLO SE, UNSPECIFIED (UNII: 3NXW29 V3WO)

FERRIC O XIDE RED (UNII: 1K09 F3G675)

FERRIC O XIDE YELLO W (UNII: EX438 O 2MRT)

MAGNESIUM STEARATE (UNII: 70097M6130)

MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)

 $\textbf{POLYDEXTROSE} \, (\textbf{UNII: VH2XOU12IE})$

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)

TRIACETIN (UNII: XHX3C3X673)

Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	8 mm
Flavor		Imprint Code	L141
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-493-65	30 in 1 CARTON	04/17/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:41250-493-72	1 in 1 CARTON	04/17/2019	

2	0 in 1 BOTTLE; Type 0: Not a Combination Product					
Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
ANDA	ANDA075400	0 4/17/20 19				

Labeler - Meijer Distribution Inc (006959555)

Revised: 1/2020 Meijer Distribution Inc