

COLD AND FLU NIGHTTIME SEVERE, 24-7 LIFE- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl tablet, film coated
Lil' Drug Store products, Inc.

Cold & Flu Nighttime Severe, 24/7 Life by 7-Eleven

Drug Facts

Drug Facts

Active Ingredients

Active Ingredients (in each caplet)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Doxylamine succinate 6.25 mg
Phenylephrine HCl 5 mg

Purpose

Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine
Nasal decongestant

Uses

Uses

- temporarily relieves common cold and flu symptoms:
- minor aches and pains
- sinus congestion and pressure
- sore throat
- fever
- headache
- nasal congestion
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- runny nose and sneezing
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

Warnings

Warnings

Liver warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- heart disease
- glaucoma
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic beverages
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Directions

- **do not take more than directed**
- do not take more than 8 caplets in 24 hours
- adults and children 12 years and over: take 2 caplets with water every 4 hours
- children under 12 years: ask a doctor

Other information

Other information

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER**

IS TORN OR BROKEN

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

Inactive ingredients

black iron oxide, corn starch, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

Questions or comments? 1-877-507-6516

Cold & Flu Nighttime Severe, 16ct, 24/7 Life - PDP/Package

[24/7 Life logo]

Cold & Flu Nighttime Severe

Acetaminophen / Pain Reliever/Fever Reducer
Dextromethorphan HBr /Cough Suppressant
Doxylamine Succinate / Antihistamine
Phenylephrine HCl / Nasal Decongestant

Relieves: Headache, Fever, Sore Throat,
Minor Aches & Pains, Nasal Congestion,
Sinus Pressure, Sneezing, Runny Nose,
Cough

compare to **Vicks**® **NyQuil**®
**VapoCOOL™ SEVERE Cold &
Flu + Congestion** active ingredients*

16

CAPLETS

Actual Size [caplet image]

24/7life
By 7-ELEVEN™

Cold & Flu Nighttime Severe

Acetaminophen / Pain Reliever/Fever Reducer
Dextromethorphan HBr /Cough Suppressant
Doxylamine Succinate / Antihistamine
Phenylephrine HCl / Nasal Decongestant

Relieves: Headache, Fever, Sore Throat, Minor Aches & Pains, Nasal Congestion, Sinus Pressure, Sneezing, Runny Nose, Cough

Compare to Vicks' NyQuil® VapoCOOL™ SEVERE Cold & Flu + Congestion (active ingredients)

16 CAPLETS

Actual Size 

PARENTS: Learn about our medicine store www.247lifemedicine.com

Satisfaction Guaranteed 1-800-255-0711
Distributed by: 7-Eleven, Inc.
 Irving, TX 75063 www.7-ELEVEN.com
*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks', NyQuil®, and VapoCOOL™ are trademarks of The Procter & Gamble Company.
66169CSE-10-22-1 50844 REV072248721

Lot # & Exp. Area
No Print/No Varnish
TAMPER EVIDENT: DO NOT USE IF PACKAGING IS OPENED OR IF BLISTER WITH TORN, BROKEN OR MISSING ANY SIGNS OF TAMPERING.
015254817251316

Drug Facts (continued)

- temporarily restores freer breathing through the nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: hives, rash, skin redness. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- heart disease
- glaucoma
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

Drug Facts (continued)

- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic beverages
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Step use and ask a doctor if

- nevousness, dizziness, or sleeplessness occurs
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- do not take more than 5 caplets in 24 hours
- adults and children 12 years and over: take 2 caplets with water every 4 hours
- children under 12 years: ask a doctor

Drug Facts **KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

Active Ingredients (in each caplet)

| Active Ingredients | Purpose |
|------------------------------|-----------------------------|
| Acetaminophen 325 mg | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg | Cough suppressant |
| Doxylamine succinate 6.25 mg | Antihistamine |
| Phenylephrine HCl 5 mg | Nasal decongestant |

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - sinus congestion and pressure
 - sore throat
 - fever
 - headache
 - nasal congestion
 - cough due to minor throat and bronchial irritation
 - cough to help you sleep
 - runny nose and sneezing
- relieves swelling of nasal passages
- promotes nasal and/or sinus drainage

Drug Facts (continued)

- each caplet contains: sodium 3 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive Ingredients black iron oxide, corn starch

Drug Facts (continued)

- crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyorbate 80, polyvinyl alcohol, polyvidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-877-807-6616

COLD AND FLU NIGHTTIME SEVERE, 24-7 LIFE

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl tablet, film coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:66715-6516 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg |
| DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL) | DOXYLAMINE SUCCINATE | 6.25 mg |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6) | |
| FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EA3S2JM) | |
| STARCH, CORN (UNII: O8232NY3S) | |
| CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561) | |
| TALC (UNII: 7SEV7J4R1U) | |
| FERROSFERRIC OXIDE (UNII: XM0M87F357) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) | |

| | |
|--|--|
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | green | Score | no score |
| Shape | OVAL | Size | 19mm |
| Flavor | | Imprint Code | 44;677 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:66715-6516-8 | 2 in 1 CARTON | 04/19/2022 | |
| 1 | | 8 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 04/19/2022 | |

Labeler - Lil' Drug Store products, Inc. (093103646)

Revised: 12/2024

Lil' Drug Store products, Inc.