

POTASSIUM CHLORIDE IN LACTATED RINGERS AND DEXTROSE- dextrose monohydrate, sodium chloride, sodium lactate, potassium chloride, calcium chloride injection, solution
Baxter Healthcare Corporation

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use POTASSIUM CHLORIDE IN LACTATED RINGER'S AND DEXTROSE (5%) INJECTION safely and effectively. See full prescribing information for POTASSIUM CHLORIDE IN LACTATED RINGER'S AND DEXTROSE (5%) INJECTION.

POTASSIUM CHLORIDE IN LACTATED RINGER'S AND DEXTROSE injection, for intravenous use
Initial U.S. Approval: 1985

RECENT MAJOR CHANGES

Dosage and Administration (2.1, 2.2, 2.3, 2.4) 10/2025

Contraindications (4) 10/2025

Warnings and Precautions (5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 5.10, 5.11) 10/2025

INDICATIONS AND USAGE

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is indicated for use as a source of water, electrolytes, and calories or as an alkalizing agent in adults and pediatric patients. (1)

DOSAGE AND ADMINISTRATION

- The recommended dosage and duration are based on the patient's age, weight, clinical condition, and concomitant therapy. (2.1)
- To reduce the risk of air embolism, adhere to the preparation instructions. (2.2, 5.2)
- Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is for intravenous use. (2.3)
- Use a peripheral vein to administer if the final dextrose concentration is 5% or less, and the osmolarity is less than 900 mOsm/L. (2.3)
- Consider using a central vein to administer hypertonic solutions with osmolarity of 900 mOsm/L or more to avoid venous irritation, including phlebitis. (2.3)
- Do not administer Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection simultaneously with ceftriaxone in neonates (28 days of age or younger) due to serious risks. (2.4)
- See full prescribing information for information dosage considerations, preparation, administration, and drug incompatibilities. (2)

DOSAGE FORMS AND STRENGTHS

Injection: Potassium Chloride (20 mEq) in Lactated Ringer's and Dextrose (5%) Injection, USP packaged in a single-dose VIAFLEX plastic container: 1000 mL. (3)

CONTRAINDICATIONS

- Concomitant treatment with ceftriaxone in neonates (28 days of age or younger). (4)
- Patients with known hypersensitivity to any of the components of Potassium Chloride in Lactated Ringer's solution and Dextrose (5%) Injection (4)
- Patients with clinically significant hyperkalemia (4).
- Patients with clinically significant hyperglycemia (4).

WARNINGS AND PRECAUTIONS

- Serious Risks with Inappropriate Use with Ceftriaxone: Deaths have occurred in neonates (28 days of age or younger) who received concomitant intravenous calcium-containing solutions with ceftriaxone. In patients older than 28 days, ceftriaxone and Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may be administered sequentially if the infusion lines are thoroughly flushed between infusions. (4, 5.1, 8.4)
- Air Embolism: Use a non-vented infusion set or close the vent on a vented set and use a dedicated line without any connections. Pressure infusion is not recommended to increase flow rates, but if necessary, remove all air from the bag prior to initiating infusion. (5.2)
- Hypersensitivity Reactions: Stop the Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection infusion immediately if signs or symptoms of a hypersensitivity reaction develop. (5.3)
- Potassium Imbalances, Hyponatremia, Neonatal Hypoglycemia, Hyperglycemia and Hyperosmolar Hyperglycemic State, Hypercalcemia, Fluid Overload, Acid-Base Imbalances, Interference with Interpretation of Serum Lactate Levels in Patients with Severe Metabolic Acidosis: See Full Prescribing Information for risk management recommendations. (5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 5.10, 5.11)

ADVERSE REACTIONS

Common adverse reactions include infusion site reactions and symptoms of hypersensitivity reactions (e.g., pruritus, dyspnea, urticaria, rash, cough). (6)

To report SUSPECTED ADVERSE REACTIONS, contact Baxter Healthcare at 1-866-888-2472 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Drugs that Affect Electrolyte and/or Fluid Balance: Avoid concomitant use. If concomitant use cannot be avoided, closely monitor electrolyte concentrations and fluid balance. (7.1)
- Lithium: Avoid concomitant use. If concomitant use is unavoidable, monitor serum lithium concentrations more frequently. (7.2)
- Digoxin: Consider reducing the volume or rate of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection due to the increased risk of digoxin toxicity with calcium-containing solutions. (7.3)
- Drugs with pH-Dependent Renal Elimination: Renal clearance of acidic drugs may be increased. In contrast, renal clearance of alkaline drugs may be decreased. (7.4)

USE IN SPECIFIC POPULATIONS

- Closely monitor plasma electrolyte concentrations in young pediatric patients with immature kidney

- function. (8.4)
- Neonates are at increased risk of developing hypo- or hyperglycemia. Closely monitor to ensure adequate glycemic control. (8.4)
- Geriatric patients are more likely to have decreased renal function. Consider monitoring renal function and starting the infusion at the low end of the dosing range. (8.5)
- Avoid in patients with severe renal impairment. (8.6)
- Patients with severe hepatic impairment may have impaired lactate metabolism. Closely monitor serum lactate levels and acid-base status. (8.7)

Revised: 10/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosage Considerations
- 2.2 Important Preparation Instructions
- 2.3 Important Administration Instructions
- 2.4 Drug Incompatibilities

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Serious Risk with Concomitant Use with Ceftriaxone
- 5.2 Air Embolism
- 5.3 Hypersensitivity Reactions
- 5.4 Potassium Imbalances
- 5.5 Hyponatremia
- 5.6 Neonatal Hypoglycemia
- 5.7 Hyperglycemia and Hyperosmolar Hyperglycemic State
- 5.8 Hypercalcemia
- 5.9 Fluid Overload
- 5.10 Acid/Base Imbalances
- 5.11 Interference of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection with Interpretation of Serum Lactate Levels in Patients with Severe Metabolic Acidosis

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

- 7.1 Drugs that Affect Electrolyte and/or Fluid Balance
- 7.2 Lithium
- 7.3 Digoxin
- 7.4 Drugs with pH-Dependent Renal Elimination
- 7.5 Interference of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection with Interpretation of Serum Lactate Levels in Patients with Severe Metabolic Acidosis

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is indicated for use as a source of water, electrolytes, and calories or as an alkalinizing agent in adults and pediatric patients.

2 DOSAGE AND ADMINISTRATION

2.1 Dosage Considerations

The recommended dosage and duration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is based on the patient's age, weight, clinical condition, and concomitant therapy. Evaluate the patient's clinical status and monitor changes in blood glucose and electrolyte concentrations especially during prolonged use of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection to optimize clinical status.

2.2 Important Preparation Instructions

Visually inspect the Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection solution for particulate matter and discoloration. Do not administer Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection unless the solution is clear and the container seals are intact.

If additives are determined to be compatible with Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection then using aseptic technique, mix thoroughly; do not store solutions containing additives. After mixing, do not use if there is discoloration or formation of precipitates.

To reduce the risk of air embolism, adhere to the following preparation instructions for Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection [see *Warnings and Precautions (5.2)*]:

- Use a non-vented infusion set or close the vent on a vented set.
- Use a dedicated line without any connections (do not connect flexible containers in series).
- The use of pressure infusion is not recommended as a method to increase flow rates. However, if pressure infusion is required, ensure that any air within the bag is fully evacuated prior to initiation of infusion.
- If using a pumping device to administer Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection, turn off the pump before the container is empty.

Preparation Instructions

1. Tear overwrap downside at slit and remove solution container.
2. Visually inspect the container.
 - o If the outlet port protector is damaged, detached, or not present, discard container as solution path sterility may be impaired.
 - o Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually.
3. Check for minute leaks by squeezing inner bag firmly. If leaks are found, discard solution as sterility may be impaired.
4. If supplemental medication is desired, follow directions below [see *Dosage and Administration (2.3)*].

Preparation for Administration

1. Suspend container from eyelet support.
2. Remove protector from outlet port at bottom of container.
3. Attach administration set according to its accompanying directions.

2.3 Important Administration Instructions

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is for intravenous use.

Use immediately after removing the overwrap. Discard the unused portion.

- Use a peripheral vein to administer Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection if the final dextrose concentration is 5% or less, and the

osmolarity is less than 900 mOsm/L.

- Consider using a central vein to administer hypertonic solutions with osmolarity of 900 mOsm/L or more to avoid venous irritation [see Warnings and Precautions (5.7)].

Some additives may be incompatible [see *Dosage and Administration (2.4)* and *Contraindications (4)*].

To Add Medication Before Solution Administration

1. Prepare medication site.
2. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
3. Mix solution and medication thoroughly. For high density medication such as potassium chloride, squeeze ports while ports are upright and mix thoroughly.

To Add Medication During Solution Administration

1. Close clamp on the set.
2. Prepare medication site.
3. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
4. Remove container from IV pole and/or turn to an upright position.
5. Evacuate both ports by squeezing them while container is in the upright position.
6. Mix solution and medication thoroughly.
7. Return container to in use position and continue administration.

2.4 Drug Incompatibilities

Do not administer Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection simultaneously with ceftriaxone in neonates (28 days of age or younger) due to serious risks [see *Contraindications (4)* and *Warnings and Precautions (5.1)*]. However, in patients older than 28 days, ceftriaxone and Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may be administered sequentially if the infusion lines are thoroughly flushed between infusions with a compatible fluid [see *Warnings and Precautions (5.1)*].

Do not administer Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection simultaneously with citrate anticoagulated/preserved blood through the same administration set because of the likelihood of coagulation precipitated by the calcium content of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection.

3 DOSAGE FORMS AND STRENGTHS

Injection: Potassium Chloride (20 mEq) in Lactated Ringer's and Dextrose (5%) Injection, USP as a clear, sterile, and nonpyrogenic solution packaged in a single-dose VIAFLEX plastic container: 1,000 mL.

4 CONTRAINDICATIONS

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is contraindicated in:

- Neonates (28 days of age or younger) who are receiving concomitant treatment with ceftriaxone, even if separate infusion lines are used, due to the risk of fatal ceftriaxone-calcium salt precipitation in the neonate's bloodstream [see *Warnings and Precautions (5.1)* and *Specific Populations (8.4)*].
- Patients with known hypersensitivity to any of the components of Potassium Chloride in Lactated Ringer's solution and Dextrose (5%) Injection [see *Warnings and Precautions (5.3)*].
- Patients with clinically significant hyperkalemia [see *Warnings and Precautions (5.4)*].
- Patients with clinically significant hyperglycemia [see *Warnings and Precautions (5.7)*].

5 WARNINGS AND PRECAUTIONS

5.1 Serious Risk with Concomitant Use with Ceftriaxone

Precipitation of ceftriaxone-calcium can occur when ceftriaxone is mixed with calcium-containing solutions, such as Potassium Chloride in Lactated Ringer's and Dextrose

(5%) Injection, in the same intravenous administration line. Deaths have occurred in neonates (28 days of age or younger) who received concomitant intravenous calcium-containing solutions with ceftriaxone resulting from calcium-ceftriaxone precipitates in the lungs and kidneys, even when separate infusion lines were used.

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is contraindicated in neonates who receive ceftriaxone [see *Contraindications (4)*, *Use in Specific Populations (8.4)*]. However, in patients older than 28 days, ceftriaxone and Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may be administered sequentially if the infusion lines are thoroughly flushed between infusions with a compatible fluid.

5.2 Air Embolism

Cases of air embolism have been reported with pressurized administration of intravenous fluids. Air embolism may result in stroke, organ ischemia and/or infarction, and death.

Use a non-vented infusion set or close the vent on a vented set and use a dedicated line without any connections.

If administration is controlled by a pumping device, care must be taken to discontinue the pumping action before the container is empty.

If administration is controlled by a pumping device, care must be taken to discontinue the pumping action before the container is empty.

Pressure infusion is **not** recommended to increase flow rates, but if necessary, ensure all air is removed from the bag before infusion.

Refrain from applying excessive pressure (>300mmHg) causing distortion to the container such as wringing or twisting. Such handling could result in breakage of the container [see *Dosage and Administration (2.2)*].

5.3 Hypersensitivity Reactions

Hypersensitivity reactions, including anaphylaxis and angioedema, have been reported with Lactated Ringer's solution. Stop the Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection infusion immediately and treat patient accordingly if signs or symptoms of a hypersensitivity reaction develop. Initiate appropriate treatment as clinically indicated.

5.4 Potassium Imbalances

Hyperkalemia

Potassium-containing solutions, including Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection, may increase the risk of hyperkalemia. This risk is increased in patients predisposed to hyperkalemia including those with severe renal impairment, acute dehydration, extensive tissue injury or burns, heart failure, or in those using concomitant drugs that are associated with hyperkalemia.

Avoid use of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients with, or at increased risk for, hyperkalemia. If use cannot be avoided in these patients, closely monitor serum potassium concentrations.

5.5 Hyponatremia

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may cause hyponatremia. Hyponatremia can lead to acute hyponatremic encephalopathy characterized by headache, nausea, seizures, lethargy and vomiting. The risk of hospital-acquired hyponatremia is increased in younger pediatric patients, geriatric patients, patients treated with diuretics, patients with cardiac or pulmonary failure, or patients with the syndrome of inappropriate antidiuretic hormone (SIADH) (e.g., postoperative patients, patients concomitantly treated with arginine vasopressin analogs, or certain antiepileptic, psychotropic, or cytotoxic drugs) [see *Drug Interactions (7.1)*, *Use in Specific Populations (8.4)*]. Avoid Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients with or at risk for hyponatremia. If use cannot be avoided in these patients, closely monitor serum sodium concentrations.

Rapid correction of hyponatremia may result in serious neurologic complications such as osmotic demyelination syndrome (ODS). To avoid complications, monitor serum sodium and chloride concentrations, fluid status, acid-base balance, and neurologic status.

5.6 Neonatal Hypoglycemia

Neonates, especially preterm neonates with low birth weight, are at increased risk of developing hypoglycemia. Closely monitor blood glucose concentration during treatment with intravenous dextrose solutions to ensure adequate glycemic control in order to avoid potential long-term adverse effects.

5.7 Hyperglycemia and Hyperosmolar Hyperglycemic State

Administration of solutions containing dextrose and lactate in patients with impaired glucose tolerance including those with diabetes mellitus may worsen hyperglycemia. Hyperglycemia is associated with an increase in serum osmolality, which can result in hypovolemia and electrolyte imbalances due to osmotic diuresis.

Patients with underlying central nervous system disease or renal impairment who receive dextrose infusions may be at greater risk of developing hyperosmolar hyperglycemic state.

While using Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection, closely monitor blood glucose concentrations and treat hyperglycemia to maintain glucose concentrations within normal limits. Anti-diabetic drugs may need to be started or dosages of these drugs may need to be increased to maintain optimal blood glucose concentrations.

5.8 Hypercalcemia

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection contains calcium salts and may cause hypercalcemia. Avoid administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients with hypercalcemia, those with calcium-containing renal calculi or history of such calculi, those with conditions predisposing to hypercalcemia, or those treated with concomitant thiazide diuretics or vitamin D.

5.9 Fluid Overload

Depending on the administered volume and the infusion rate, administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection can cause fluid overload, including pulmonary edema.

Avoid Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients at risk for fluid and/or solute overload including patients with severe renal impairment. If use cannot be avoided in these patients, monitor fluid balance, electrolyte concentrations and acid base balance, especially during prolonged use.

5.10 Acid/Base Imbalances

Because lactate is metabolized to bicarbonate, administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may result in, or worsen, metabolic alkalosis. Closely monitor the acid-base balance in patients with, or at risk of, alkalosis.

In patients with severe hepatic impairment, decreased lactate metabolism may result in worsening anion gap metabolic acidosis. Avoid Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients with severe hepatic impairment. If use cannot be avoided in these patients, closely monitor serum bicarbonate levels.

5.11 Interference of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection with Interpretation of Serum Lactate Levels in Patients with Severe Metabolic Acidosis

Administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may result in interference with the interpretation of serum lactate levels in patients with severe metabolic acidosis, including lactic acidosis [see *Drug Interactions (7.5)*].

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections of the labeling:

- Serious Risk with Concomitant Use with Ceftriaxone [see *Warnings and Precautions (5.1)*]
- Air Embolism [see *Warnings and Precautions (5.2)*]
- Hypersensitivity Reactions [see *Warnings and Precautions (5.3)*]
- Potassium Imbalances [see *Warnings and Precautions (5.4)*]

- Hyponatremia [see Warnings and Precautions (5.5)]
- Neonatal Hypoglycemia [see Warnings and Precautions (5.6)]
- Hyperglycemia and Hyperosmolar Hyperglycemic State [see Warnings and Precautions (5.7)]
- Hypercalcemia [see Warnings and Precautions (5.8)]
- Fluid Overload [see Warnings and Precautions (5.9)]
- Acid/Base Imbalances [see Warnings and Precautions (5.10)]

The following adverse reactions have been identified during postapproval use of Lactated Ringer's products. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure:

General Disorders and Administration Site Conditions:

Phlebitis, extravasation, infusion site inflammation, infusion site swelling, infusion site rash, infusion site pruritus, infusion site erythema, infusion site pain, infusion site burning, and infusion site hypoaesthesia.

Hypersensitivity Reactions and Infusion Reactions:

Angioedema, chest pain/discomfort, bradycardia or tachycardia, hypotension, respiratory distress, bronchospasm, dyspnea, cough, urticaria, rash, pruritus, erythema, flushing, throat irritation, paresthesia, oral hypoesthesia, dysgeusia, nausea, anxiety, pyrexia, and headache, laryngeal edema, sneezing, injection site infection.

Metabolism and Nutrition Disorders:

Hyperkalemia, hyponatremia, and hypervolemia.

Nervous System Disorders:

Hyponatremic encephalopathy.

7 DRUG INTERACTIONS

7.1 Drugs that Affect Electrolyte and/or Fluid Balance

Hyperkalemia

Administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection to patients concomitantly treated or recently treated with drugs that are associated with hyperkalemia increases the risk of severe and potentially fatal hyperkalemia, especially in the presence of other hyperkalemia risk factors. Avoid use of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients receiving drugs that are associated with hyperkalemia (e.g., potassium-sparing diuretics, ACE inhibitors, angiotensin II receptor antagonists, or calcineurin inhibitors). If concomitant use cannot be avoided, closely monitor serum potassium concentrations during concomitant use [see Warnings and Precautions (5.4)].

Hyponatremia

Administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection to patients treated concomitantly with drugs associated with hyponatremia may increase the risk of developing hyponatremia. These drugs include diuretics and those that cause SIADH (e.g., arginine vasopressin analogs, certain antiepileptic, psychotropic, or cytotoxic drugs). Avoid use of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients receiving such drugs. If use cannot be avoided, closely monitor serum sodium concentrations during concomitant use [see Warnings and Precautions (5.5)].

Hypercalcemia

Avoid the use of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients treated with thiazide diuretics or vitamin D because these drugs can increase the risk of hypercalcemia. If use cannot be avoided, closely monitor serum calcium concentrations during concomitant use [see Warnings and Precautions (5.8)].

Hypernatremia and Fluid Retention

Administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection to patients treated concomitantly with drugs associated with sodium and fluid retention (e.g., corticosteroids or corticotropin) may increase the risk of hypernatremia and volume overload. Avoid use of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients receiving such drugs. If use cannot be avoided, closely monitor serum electrolytes, fluid balance, and acid-base balance during concomitant use.

7.2 Lithium

Renal sodium and lithium clearance may be increased during concomitant use of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection and lithium and

may result in decreased lithium concentrations. Avoid use of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients receiving lithium. If use cannot be avoided, increase the frequency of monitoring of serum lithium concentrations during concomitant use.

7.3 Digoxin

Administration of calcium via use of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may increase digoxin's effects and lead to digoxin toxicity including serious or fatal cardiac arrhythmias. In digoxin-treated patients, consider reducing the volume and/or rate of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection administration.

7.4 Drugs with pH-Dependent Renal Elimination

Due to the alkalinizing action of lactate (formation of bicarbonate), Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may interfere with the elimination of drugs with pH-dependent renal elimination. Renal clearance of alkaline drugs may be decreased. In contrast, renal clearance of acidic drugs may be increased.

7.5 Interference of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection with Interpretation of Serum Lactate Levels in Patients with Severe Metabolic Acidosis

Because administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may interfere with the interpretation of serum lactate levels in patients with severe metabolic acidosis, including lactic acidosis, assessment of the patient's clinical status should not solely rely on the measurement of serum lactate.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection as a source of water and electrolytes has been used for decades during labor and delivery. Appropriate administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is not expected to cause major birth defects, miscarriage, or adverse maternal or fetal outcomes. Animal reproduction studies have not been conducted with this drug.

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

8.2 Lactation

Risk Summary

Lactated Ringer's as a source of water and electrolytes and Dextrose (5%) Injection have been used for decades and is not expected to cause harm to a breastfed infant. There are no data on the presence of Lactated Ringer's and Dextrose (5%) Injection in human milk, the effects on the breastfed infant, or the effects on milk production. Potassium is present in human breast milk. There are no data on the effects of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection on a breastfed infant or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection and any potential adverse effects on the breastfed infant from Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection or from the underlying maternal condition.

8.4 Pediatric Use

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is contraindicated in neonates (28 days of age or younger) who are receiving ceftriaxone due to reported deaths that occurred when neonates received ceftriaxone and intravenous calcium-containing solutions concomitantly [*see Warnings and Precautions (5.1)*].

The safety and effectiveness of Lactated Ringer's Injection for use as a source of water, electrolytes, and calories or as an alkalinizing agent have been established in pediatric patients of all ages, including neonates.

The safety profile of Potassium Chloride in Dextrose Injection in pediatric patients is similar to adults.

Closely monitor plasma electrolyte concentrations in young pediatric patients with immature kidney function who may have decreased ability to maintain fluid and electrolyte balance [see *Warnings and Precautions (5.4, 5.5, 5.8, 5.9)*]. Administration of a lactate-containing intravenous solution, including Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection to pediatric patients should account for liver and kidney maturation (the kidney function affects the biotransformation and renal excretion of lactate) [see *Warnings and Precautions (5.9)*]. Pediatric patients are at increased risk for developing hyponatremic encephalopathy [see *Warnings and Precautions (5.5)*].

Neonates, especially preterm neonates with low birth weight, are at increased risk of developing hypo- or hyperglycemia and therefore need close monitoring during treatment with intravenous glucose solutions including Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection to ensure adequate glycemic control to avoid potential long-term adverse reactions [see *Warnings and Precautions (5.6, 5.7)*]. In very low birth weight neonates, excessive or rapid administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may result in increased serum osmolarity and risk of intracranial hemorrhage.

8.5 Geriatric Use

Geriatric patients treated with Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection are at increased risk of developing electrolyte imbalances. Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is substantially excreted by the kidney, and the risk of adverse reactions to Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may be greater in patients with renal impairment than in patients with normal renal function. Because geriatric patients are more likely to have decreased renal function, consider monitoring renal function in geriatric patients and consider starting the infusion at the low end of the dosing range.

8.6 Renal Impairment

Administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection to patients with or at risk of severe renal impairment, may result in hyperkalemia and/or fluid overload [see *Warnings and Precautions (5.4, 5.7, 5.9)*]. Avoid Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection in patients with severe renal impairment. If use cannot be avoided in such patients, monitor for development of these adverse reactions.

8.7 Hepatic Impairment

In patients with severe hepatic impairment, lactate metabolism may be impaired, and Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection may not produce alkalization. Closely monitor serum lactate levels and acid-base status in such patients.

10 OVERDOSAGE

Excessive administration of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection can cause:

- Hyperkalemia and hyponatremia, especially in patients with severe renal impairment.
- Fluid overload (which can lead to pulmonary and/or peripheral edema).
- Hyperglycemia, hyperosmolarity, and osmotic diuresis, dehydration, and electrolyte loss.
- Metabolic alkalosis with or without hypokalemia.
- Loss of bicarbonate with an acidifying effect.
- Hypercalcemia.

Overdose interventions include Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection discontinuation, treatment of electrolyte imbalances, treatment of hyperglycemia, and close monitoring of fluid balance, electrolyte concentrations, and acid-base balance [see *Warnings and Precautions (5.4, 5.5, 5.6, 5.7, 5.9, 5.10)*]

11 DESCRIPTION

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection, USP is a sterile, nonpyrogenic solution for fluid and electrolyte replenishment and caloric supply in a single dose container for intravenous administration. It contains no antimicrobial agents.

Composition, osmolarity, pH, ionic concentration and caloric content are shown in Table 1. Dextrose is derived from corn.




Table 1

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection, USP	Size (mL)	Composition (g/L)					*Osmolarity (mOsmol/L) (calc)	pH	Ionic Concentration (mEq/L)					Caloric Content (kcal/L)
		Dextrose, USP	Sodium Chloride, USP	Sodium Lactate	Potassium Chloride, USP	Calcium Chloride, USP			Sodium	Potassium	Calcium	Chloride	Lactate	
20 mEq Potassium added	1000	50	6	3.1	1.79	0.2	565	5.0 (3.5 to 6.5)	130	24	3	129	28	170

* Normal physiologic osmolarity range is approximately 280 to 310 mOsmol/L.

The chemical name, structural formula, and molecular weight of the active ingredients are shown in Table 2.

Table 2

Chemical Name	Structural Formula	Molecular Weight
Sodium Chloride, USP	$\text{Na}^+ \text{Cl}^-$	58.44
Sodium Lactate		112.06
Potassium Chloride, USP	$\text{K}^+ \text{Cl}^-$	74.55
Calcium Chloride, USP	$\cdot 2\text{H}_2\text{O}$ 	147.02
Dextrose, USP		198.17

The VIAFLEX plastic container is fabricated from polyvinyl chloride (PL 146 Plastic). Small amounts of its chemical components (e.g., di-2-ethylhexyl phthalate [DEHP]) may leach out within the expiration period. Animal tests, USP biological tests for plastic containers, and tissue culture toxicity studies have demonstrated that the container meets safety requirements. The container is overwrapped to provide protection from the physical environment and an additional moisture barrier.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection is a source of water, electrolytes, and calories, and produces an alkalinizing effect.

- Sodium, the major cation of the extracellular fluid, functions primarily in the control of water distribution, fluid balance, and osmotic pressure of body fluids. Sodium is also associated with chloride and bicarbonate in the regulation of the acid-base equilibrium of body fluid.
- Potassium, the principal cation of intracellular fluid, participates in carbohydrate utilization and protein synthesis and is critical in the regulation of nerve conduction and muscle contraction, particularly in the heart.
- Chloride, the major extracellular anion, closely follows the metabolism of sodium, and changes in the acid-base balance of the body are reflected by changes in the

chloride concentration.

- Calcium, an important cation, provides the framework of bones and teeth in the form of calcium phosphate and calcium carbonate. In the ionized form, calcium is essential for the functional mechanism of the clotting of blood, normal cardiac function, and regulation of neuromuscular irritability.
- Sodium lactate provides sodium and lactate ions. The lactate anion is in equilibrium with pyruvate and has an alkalinizing effect resulting from simultaneous removal by the liver of lactate and hydrogen ions. The sodium ion combines with bicarbonate ion produced from carbon dioxide of the body and thus retains bicarbonate to combat metabolic acidosis (bicarbonate deficiency).
- Dextrose provides a source of calories. Dextrose may decrease losses of body protein and nitrogen, promotes glycogen deposition and decreases or prevents ketosis if sufficient doses are provided.

12.2 Pharmacodynamics

The exposure-response relationship and time course of pharmacodynamic response for the safety and effectiveness of Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection have not been fully characterized.

12.3 Pharmacokinetics

Elimination

Metabolism/Excretion

Potassium: Normally about 80 to 90% of the potassium intake is excreted in the urine; the remainder is excreted in feces and to a smaller extent, in perspiration.

Sodium and Chloride: The distribution and excretion of sodium (Na⁺) and chloride (Cl⁻) are largely under the control of the kidney which maintains a balance between intake and output.

Lactate: In the liver, lactate is metabolized to carbon dioxide and water by oxidative metabolism and consumption of hydrogen cations.

Dextrose: Dextrose injected parenterally undergoes oxidation to carbon dioxide and water.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity, genetic toxicology, and animal fertility studies have not been conducted with Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection.

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

Potassium Chloride in Lactated Ringer's and Dextrose (5%) Injection, USP is supplied as a clear solution in single-dose VIAFLEX plastic containers as follows:

Code	Size (mL)	NDC	Product Name
2B2224	1000	0338-0811-04	20 mEq/L Potassium Chloride in Lactated Ringer's and 5% Dextrose Injection, USP

Storage and Handling

Store at room temperature (recommended 25°C/77°F).

Brief exposure up to 40°C (104°F) does not adversely affect the product. Exposure to heat should be minimized. Avoid excessive heat.

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Baxter, PL 146 and Viaflex are trademarks of Baxter International Inc. or its subsidiaries.

07-19-00-9427

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

LOT

EXP



2B2224
NDC 0338-0811-04
DIN 00786314

1

20 mEq Potassium Chloride

2

3

(20 mEq/L) Potassium Chloride in
Lactated Ringer's and 5% Dextrose
Injection USP

4

5

1000 mL

EACH 100 mL CONTAINS 5 g DEXTROSE HYDROUS USP 600 mg SODIUM CHLORIDE USP 310 mg SODIUM LACTATE 179 mg POTASSIUM CHLORIDE USP 20 mg CALCIUM CHLORIDE USP pH 5.0 (3.5 TO 6.5) mEq/L SODIUM 130 POTASSIUM 24 CALCIUM 3 CHLORIDE 129 LACTATE 28 HYPERTONIC OSMOLARITY 565 mOsmol/L (CALC) STERILE NONPYROGENIC SINGLE DOSE CONTAINER **NOT FOR USE IN THE TREATMENT OF LACTIC ACIDOSIS** ADDITIVES MAY BE INCOMPATIBLE CONSULT WITH PHARMACIST IF AVAILABLE WHEN INTRODUCING ADDITIVES USE ASEPTIC TECHNIQUE MIX THOROUGHLY Do NOT STORE Dosage INTRAVENOUSLY AS DIRECTED BY A PHYSICIAN SEE DIRECTIONS CAUTIONS SQUEEZE AND INSPECT INNER BAG WHICH MAINTAINS PRODUCT STERILITY DISCARD IF LEAKS ARE FOUND MUST NOT BE USED IN SERIES CONNECTIONS Do NOT ADMINISTER SIMULTANEOUSLY WITH BLOOD Do NOT USE UNLESS SOLUTION IS CLEAR Rx ONLY STORE UNIT IN MOISTURE BARRIER OVERWRAP AT ROOM TEMPERATURE (25°C/77°F) UNTIL READY TO USE AVOID EXCESSIVE HEAT SEE INSERT

6

7

VIAFLEX PLUS CONTAINER PL 146 PLASTIC
BAXTER VIAFLEX AND PL 146 ARE TRADEMARKS OF
BAXTER INTERNATIONAL INC
FOR PRODUCT INFORMATION 1-800-933-0303

8

Baxter

BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN USA

DISTRIBUTED IN CANADA BY
BAXTER CORPORATION
TORONTO ONTARIO CANADA

9

Container Label

LOT

EXP

2B2224
NDC 0338-0811-04
DIN 00786314

20 mEq

Potassium Chloride

(20 mEq/L) Potassium Chloride in
Lactated Ringer's and 5% Dextrose
Injection USP

1000 mL

EACH 100 mL CONTAINS 5 g DEXTROSE HYDROUS USP 600 mg SODIUM CHLORIDE USP 310 mg SODIUM LACTATE 179 mg POTASSIUM CHLORIDE USP 20 mg CALCIUM CHLORIDE USP pH 5.0 (3.5 TO 6.5) mEq/L SODIUM 130 POTASSIUM 24 CALCIUM 3 CHLORIDE 129 LACTATE 28 HYPERTONIC

OSMOLARITY 565 mOsmol/L (CALC) STERILE NONPYROGENIC
 SINGLE DOSE CONTAINER **NOT FOR USE IN THE TREATMENT OF
 LACTIC ACIDOSIS** ADDITIVES MAY BE INCOMPATIBLE CONSULT WITH
 PHARMACIST IF AVAILABLE WHEN INTRODUCING ADDITIVES USE
 ASEPTIC TECHNIQUE MIX THOROUGHLY DO NOT STORE DOSAGE
 INTRAVENOUSLY AS DIRECTED BY A PHYSICIAN SEE DIRECTIONS
 CAUTIONS SQUEEZE AND INSPECT INNER BAG WHICH MAINTAINS
 PRODUCT STERILITY DISCARD IF LEAKS ARE FOUND MUST NOT BE
 USED IN SERIES CONNECTIONS DO NOT ADMINISTER
 SIMULTANEOUSLY WITH BLOOD DO NOT USE UNLESS SOLUTION IS
 CLEAR Rx ONLY STORE UNIT IN MOISTURE BARRIER OVERWRAP AT
 ROOM TEMPERATURE (25°C/77°F) UNTIL READY TO USE AVOID
 EXCESSIVE HEAT SEE INSERT

VIAFLEX PLUS CONTAINER

PL 146 PLASTIC

BAXTER VIAFLEX AND PL 146 ARE TRADEMARKS OF
 BAXTER INTERNATIONAL INC

FOR PRODUCT INFORMATION 1-800-933-0303

Baxter

BAXTER HEALTHCARE CORPORATION
 DEERFIELD IL 60015 USA

MADE IN USA

DISTRIBUTED IN CANADA BY
BAXTER CORPORATION
 TORONTO ONTARIO CANADA

POTASSIUM CHLORIDE IN LACTATED RINGERS AND DEXTROSE				
dextrose monohydrate, sodium chloride, sodium lactate, potassium chloride, calcium chloride injection, solution				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0811	
Route of Administration	INTRAVENOUS			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G) (ANHYDROUS DEXTROSE - UNII:5SLOG7R0OK)		DEXTROSE MONOHYDRATE	5 g in 100 mL	
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)		SODIUM CHLORIDE	600 mg in 100 mL	
SODIUM LACTATE (UNII: TU7HW0W0QT) (SODIUM CATION - UNII:LYR4M0NH37, LACTIC ACID - UNII:33X04XA5AT)		SODIUM LACTATE	310 mg in 100 mL	
POTASSIUM CHLORIDE (UNII: 660YQ98I10) (POTASSIUM CATION - UNII:295O53K152, CHLORIDE ION - UNII:Q32ZN48698)		POTASSIUM CHLORIDE	179 mg in 100 mL	
CALCIUM CHLORIDE (UNII: M410D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)		CALCIUM CHLORIDE	20 mg in 100 mL	
Inactive Ingredients				
Ingredient Name		Strength		
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0811-04	1000 mL in 1 BAG; Type 0: Not a Combination Product	04/05/1985	
Marketing Information				
Marketing	Application Number or Monograph		Marketing Start	Marketing End

Category	Citation	Date	Date
NDA	NDA019367	04/05/1985	

Labeler - Baxter Healthcare Corporation (005083209)

Establishment			
Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		059140764	ANALYSIS(0338-0811) , MANUFACTURE(0338-0811) , LABEL(0338-0811) , PACK(0338-0811) , STERILIZE(0338-0811) , API MANUFACTURE(0338-0811)

Establishment			
Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		194684502	ANALYSIS(0338-0811)

Establishment			
Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		189326168	ANALYSIS(0338-0811) , MANUFACTURE(0338-0811) , LABEL(0338-0811) , PACK(0338-0811) , STERILIZE(0338-0811)

Revised: 10/2025

Baxter Healthcare Corporation