# MENSTRUAL PAIN RELIEF COMPLETE- acetaminophen, pyrilamine maleate, caffeine tablet, film coated Shield Pharmaceuticals Corp

-----

# **Menstrual Pain Relief Complete**

# **Drug Facts**

## Active ingredient (in each caplet)

Acetaminophen 500 mg Caffeine 60 mg Pyrilamine maleate 15 mg

## **Purpose**

Pain reliever Diuretic Antihistamine

#### □ Uses

for the temporary relief of these symptoms associated with menstrual periods:

- cramps
- bloating
- water
- weight gain
- headache
- backache
- muscle aches
- fatigue

# Warnings

**Liver warning:**This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)

shock

If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

#### Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- liver disease glaucoma
- a breathing problem such as emphysema or chronic

## Ask a doctor or pharmacist before use if youare

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

## When using this product

- avoid alcoholic beverages
- excitability may occur, especially in children
- be careful when driving a motor vehicle or operating machiner
- alcohol, sedatives, and tranquilizers may increase drowsiness
- you may get drowsy
- limit the use of caffeine-containing medications, foods, or beverages while taking this
  product because too much caffeine may cause nervousness, irritability,
  sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this
  product contains about as much caffeine as a cup of coffee.

# Stop use and ask a doctor if

- new symptoms occur \_ redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

# adults and children 12 years and over:

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 caplets with water
- repeat every 6 hours, as needed
- do not exceed 6 caplets per day

children under 12 years: ask a doctor

#### Other information

• store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

#### Inactive ingredients

calcium phosphate, colloidal silicon dioxide, croscarmellose sodium, FD&C yellow no. 6, hypromellose, isopropyl alcohol, magnesium stearate, methylene chloride, microcrystalline cellulose, polyethylene glycols, polyvinyl pyrrolidone, povidone, pregelatinized starch, sodium starch glycolate\*, stearic acid, talc, titanium dioxide \*may contain this ingredient

#### Questions or comments?

**1-800-373-6981** (toll-free)

Distributed by:

Shield Pharmaceuticals Corp.

Ronkonkoma, NY 11779


#### PRINCIPAL DISPLAY PANEL

Menstrual Pain Relief Complete

Acetaminophen/Caffeine/Pyrilamine maleate

Pain reliever/Diuretic/Antihistamine

Label



Ask a doctor before use if you have  I difficulty in unination due to enlargement of the prostate gland  I liver disease I glaucoma I abreathing problem such as emphysema or chronic bronchitis  Ask a doctor or pharmacist before use if you are  I taking the blood thinning drug warfarin I aking sedatives or tranquilizers  When using this product  I avoid alcoholic beverages I excitability may occur, especially in children I be careful when driving a motor vehicle or operating machinery I alcohol; sedatives or tranquilizers may increase drowsiness I you may get drows; I limit the use of caffeine-containing medications, bods, or beverages while taking the use of caffeine-containing medications, bods, or beverages while taking the account of the containing medications, bods, or beverages while taking the containing medications, bods, or beverages while taking the account of ordiffer.  Stop use and ask a doctor if I new symptoms occur. I endease or lasts more than 10 days  I freganan tor breast-feeding, ask a health professional before use.  Stop use and ask a doctor if I be accounted to see the containing about as much caffeine as a cup of ordiffer.  Stop use and ask a doctor if I before use.  Stop use and ask a doctor if I before use.  Stop use out of reach of children. In case of overdose, get medical altention is ordical for adults as well as for children even if you do not notice any signs or symptoms.  Directions  do not take more than the recommended dose  adults and children 12 years and over.  I repeat every 6 hours, as needed  on ont ake more than the recommended dose  on ont ake more than the recommended dose  submission of capters be a doctor  Other information  store at 29% (77%) excursions permitted between 159-30%  (59-86%)  Inactive information  store at 29% (77%) excursions permitted between 159-30%  (59-86%)  Inactive information  stored every folloria, sportory disorion, registering as each well with a signation or continuents?  1-800-35-36-36-36-36-36-36-36-36-36-36-36-36-36-
--

# **MENSTRUAL PAIN RELIEF COMPLETE**

acetaminophen, pyrilamine maleate, caffeine tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83059-0119
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	500 mg	
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	60 mg	
PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)	PYRILAMINE MALEATE	15 mg	

Inactive Ingredients		
Ingredient Name	Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
CALCIUM PHOSPHATE (UNII: 97Z1W3NDX)		
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
HYPROMELLOSES (UNII: 3NXW29V3WO)		
ISOPROPYL ALCOHOL (UNII: ND2M416302)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
METHYLENE CHLORIDE (UNII: 588X2YUY0A)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
STARCH, CORN (UNII: 08232NY3SJ)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
PYRROLIDONE (UNII: KKL5D39EOL)		
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)		
POVIDONE (UNII: FZ 989GH94E)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
TALC (UNII: 7SEV7J4R1U)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		

Product Characteristics			
Color	white	Score	no score
Shape	CAPSULE	Size	17mm
Flavor		Imprint Code	SD33
Contains			

Packaging				
# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1 NDC:83059- 0119-8	80 in 1 BOTTLE; Type 0: Not a Combination Product	12/05/2025		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	12/05/2025		

Labeler - Shield Pharmaceuticals Corp (118724924)

Revised: 12/2025 Shield Pharmaceuticals Corp