

MEDI-FIRST IBUPROFEN- ibuprofen tablet, coated
MEDI-FIRST PLUS IBUPROFEN- ibuprofen tablet, coated
MEDIQUE IPRIN- ibuprofen tablet, coated
DOVER ADDAPRIN- ibuprofen tablet, coated
OTIS CLAPP ULTRAPRIN- ibuprofen tablet, coated
Unifirst First Aid Corporation

UniFirst First Aid Ibuprofen

Drug Facts

Active ingredient

Ibuprofen 200 mg (NSAID)

*nonsteroidal antiinflammatory drug

Purpose

Pain reliever/fever reducer

Uses

Temporarily relieves minor aches and pains associated with

- headache ■ toothache ■ backache ■ menstrual cramps
- common cold ■ muscular aches ■ minor arthritis pain

Temporarily reduces fever

Warnings

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack or stroke warning:

NSAIDS, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

If pregnant or breast feeding, ask a health professional before use. It is especially important not

to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not use more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults and children:(12 years and older)

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years:

Do not give to children under 12 years of age.

Other information

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 40°C (above 104°F)
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

carnauba wax*, corn starch, hypromellose*, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, polyvinyl alcohol*, povidone K30*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide

**may contain*

Questions or comments? 1-800-634-7680

Medi-First Ibuprofen Label

100 tablets (50 x 2)

Medi-First®

Ibuprofen 200 mg

Pain Reliever/Fever Reducer (NSAID)

Aches, Fever • Ibuprofen (NSAID) 200 mg

Pull to Open

Compare active ingredient to:

Advil®

Registered Trademark of Pfizer Consumer Healthcare

This package is for Households without Young Children

Tamper Evident Unit Dose Packets

Drug Facts
Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID).....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
Temporarily relieves minor aches and pains associated with:
■ headache ■ toothache ■ backache ■ menstrual cramps
■ common cold ■ muscular aches ■ minor arthritis pain
Temporarily reduces fever.

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ skin redness ■ asthma (wheezing)
■ facial swelling ■ rash ■ shock ■ dizziness
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued)
Have 3 or more alcoholic drinks every day while using this product
Take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery
Ask a doctor before use if
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ stomach bleeding warning applies to you
■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
■ under a doctor's care for any serious condition
■ taking any other drug (continued on opposite panel)

MEDI-FIRST®
Ibuprofen 200mg
Pain Reliever/Fever Reducer
✓ Aches, Fever - Ibuprofen (NSAID) 200mg
Pull To Open
Compare active ingredient to:
Advil®
Registered Trademark of Pfizer Consumer Healthcare
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
Tamper Evident Unit Dose Packets

Drug Facts (continued)
When using this product
■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ you have stomach pain that does not get better
■ you have symptoms of heart problems or stroke
■ chest pain ■ trouble breathing
■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts for more than 10 days
■ fever gets worse or lasts for more than 3 days
■ redness or swelling is present in the painful area
■ any new or unexpected symptoms occur
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ do not use more than directed
■ the smallest effective dose should be used
■ do not take longer than 10 days, unless directed by a doctor (see Warnings)
Adults and children (12 years and older) Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.
Children under 12 years: Do not give to children under 12 years of age.
Other information
■ read all product information before using
■ store at 68-77°F (20-25°C)
■ avoid excessive heat 104°F (above 40°C)
■ tamper evident sealed packets
■ do not use any opened or torn packets (continued on opposite panel)

Drug Facts (continued)
Inactive ingredients
carnauba wax, corn starch, hypromellose, iron oxide red, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, polyvinyl alcohol, povidone (K-30), silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide
*may contain
Questions or comments? 1-800-634-7680

3 47682 80833 4
MEDI-FIRST®
Revider #80833
Manufactured for
Medique Products
Fort Myers, FL 33907
An FDA-Registered Facility
1-800-634-7680
www.mediqueproducts.com

Retain carton for complete product information

Medi-First Plus Ibuprofen Label

100 tablets (50 x 2's)

Medi-First® Plus

Ibuprofen

Ibuprofen 200 mg (NSAID)

This package is for Households without Young Children

Pull To Open

Pain Reliever/Fever Reducer

Compare active ingredient to:

Advil®

Registered Trademark of Pfizer Consumer Healthcare

Tamper Evident Unit Dose Packets

Drug Facts
Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID).....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
Temporarily relieves minor aches and pains associated with:
■ headache ■ toothache ■ backache ■ menstrual cramps
■ common cold ■ muscular aches ■ minor arthritis pain
Temporarily reduces fever.

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ skin redness ■ asthma (wheezing)
■ facial swelling ■ rash ■ shock ■ dizziness
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued)
Have 3 or more alcoholic drinks every day while using this product
Take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery
Ask a doctor before use if
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ stomach bleeding warning applies to you
■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
■ under a doctor's care for any serious condition
■ taking any other drug (continued on opposite panel)

MEDI-FIRST®
Plus
Ibuprofen 200mg (NSAID)
Pull To Open
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
Pain Reliever/Fever Reducer
Compare active ingredient to:
Advil®
Registered Trademark of Pfizer Consumer Healthcare
Tamper Evident Unit Dose Packets

Drug Facts (continued)
When using this product
■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ you have stomach pain that does not get better
■ you have symptoms of heart problems or stroke
■ chest pain ■ trouble breathing
■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts for more than 10 days
■ fever gets worse or lasts for more than 3 days
■ redness or swelling is present in the painful area
■ any new or unexpected symptoms occur
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ do not use more than directed
■ the smallest effective dose should be used
■ do not take longer than 10 days, unless directed by a doctor (see Warnings)
Adults and children (12 years and older) Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.
Children under 12 years: Do not give to children under 12 years of age.
Other information
■ read all product information before using
■ store at 68-77°F (20-25°C)
■ avoid excessive heat 104°F (above 40°C)
■ tamper evident sealed packets
■ do not use any opened or torn packets (continued on opposite panel)

Drug Facts (continued)
Inactive ingredients
carnauba wax, corn starch, hypromellose, iron oxide red, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, polyvinyl alcohol, povidone (K-30), silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide
*may contain
Questions or comments? 1-800-634-7680

3 47682 90833 1
Revider #90833
Manufactured for Medi-First Plus
Fort Myers, FL 33907 USA
An FDA-Registered Facility
1-800-634-7680
www.mediqueproducts.com

Retain carton for complete product information

Medique Iprin Label

Medique®

Collect MediBucks

See inside flap for more details

I-Prin

Drug Facts		Drug Facts (continued)
Active ingredient (in each tablet) Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer <small>*nonsteroidal anti-inflammatory drug</small>		When using this product ■ take with food or milk if stomach upset occurs
Purpose Pain reliever/fever reducer		Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ redness or swelling is present in the painful area ■ any new or unexpected symptoms occur
Uses Temporarily relieves minor aches and pains associated with ■ headache ■ toothache ■ backache ■ menstrual cramps ■ common cold ■ muscular aches ■ minor arthritis pain		If pregnant or breast-feeding , ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Temporarily reduces fever.		Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ skin redness ■ asthma (wheezing) ■ facial swelling ■ rash ■ stroke ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.		Directions ■ do not use more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor (see Warnings)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed		Adults and children (12 years and older) Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.		Children under 12 years: Do not give to children under 12 years of age.
Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery		Other information ■ read all product information before using ■ store at 68-77°F (20-25°C) ■ stomach bleeding warning applies to you ■ avoid excessive heat 104°F (above 40°C) ■ tamper evident sealed packets ■ do not use any opened or torn packets
Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke ■ you are taking a diuretic		Inactive ingredients camsauba wax [®] , com starch, hypromellose [®] , iron oxide red, lactose [®] , magnesium stearate [®] , microcrystalline cellulose [®] , polydextrose [®] , polyethylene glycol, polyvinyl alcohol [®] , povidone (K-30) [®] , silicon dioxide, sodium starch glycolate, stearic acid, talc [®] , titanium dioxide
Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug		*may contain Questions or comments? 1-800-634-7680

Retain carton for complete product information

Dover **Addaprin™**
PAIN RELIEVER-FEVER REDUCER
IBUPROFEN 200 MG. TABLETS (NSAID)

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

Dover Pharmaceutical
 Products of the highest quality
 and effectiveness

TAMPER EVIDENT
 Sealed Packets

UNIT DOSE PACKS
500 TABLETS
(250 PACKETS OF 2)

Otis Clapp Ultraprin Label

OC Otis Clapp

Quality & Integrity Since 1840

Ultraprin™

Pain Reliever-Fever Reducer (NSAID)

Ibuprofen Tablets USP 200 mg

For Deep Seated Pain

See Warnings and Directions on Side Panel

Tear Out Along Perforation To Dispense

Professional Healthcare

500 Tablets (250 Packets of 2)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-718-13	250 in 1 BOX	01/26/2017	
1	NDC:47682-718-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-718-48	125 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-718-33	50 in 1 BOX	01/26/2017	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-718-30	4 in 1 BOX	01/26/2017	
4		2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:47682-718-50	25 in 1 BOX	04/16/2019	
5		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/26/2017	

MEDI-FIRST PLUS IBUPROFEN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-709
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	red (Reddish Brown)	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-709-48	125 in 1 BOX	01/26/2017	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-709-33	50 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/26/2017	

MEDIQUE IPRIN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-700
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POVIDONE K30 (UNII: U725QWY32X)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	red (Reddish Brown)	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-700-69	3 in 1 BOX	01/26/2017	
1	NDC:47682-700-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-700-64	12 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-700-47	100 in 1 BOX	01/26/2017	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-700-13	250 in 1 BOX	01/26/2017	
4		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/26/2017	

DOVER ADDAPRIN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-714
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

TALC (UNII: 7SEV7J4R1U)

Product Characteristics

Color	red (Reddish Brown)	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-714-13	250 in 1 BOX	01/26/2017	
1	NDC:47682-714-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/26/2017	

OTIS CLAPP ULTRAPRIN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-702
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, CORN (UNII: O8232NY3SJ)	

FERRIC OXIDE RED (UNII: 1K09F3G675)

Product Characteristics

Color	red (Reddish Brown)	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-702-13	250 in 1 BOX	02/01/2017	04/03/2017
1	NDC:47682-702-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	11/17/2014	04/03/2017

Labeler - Unifirst First Aid Corporation (832947092)

Establishment

Name	Address	ID/FEI	Business Operations
Prestige Packaging		170837962	relabel(47682-700, 47682-702, 47682-709, 47682-714, 47682-718) , repack(47682-700, 47682-702, 47682-709, 47682-714, 47682-718)

Revised: 4/2020

Unifirst First Aid Corporation