

**DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hbr,  
phenylephrine hcl liquid  
BI-MART**

-----  
**DAYTTIME COLD & FLU**

***Drug Facts***

***Active ingredients (in each 15 mL dose cup)***

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer

Cough Suppressant

Nasal Decongestant

***Uses***

temporarily relieves common cold and flu symptoms:

- minor aches and pains
- headache
- sore throat
- fever
- nasal congestion
- cough due to minor throat and bronchial irritation

***Warnings***

**Liver warning:** This product contains acetaminophen. Sever liver damage may occur if:

- adults and children over 12 years of age take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- children 6 to 12 years of age take more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- take with other drug containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause sever skin reactions. Symptoms may include:

- skin reddening
- blisters
- ras

If skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- If you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough accompanied by excessive phlegm (mucus)
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema.

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product**

- do not use more than directed (see overdose warning)
- avoid alcoholic drinks

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough get worse or last more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:**

Taking more than the recommended dose (overdose) may cause serious health problems including liver damage. In case of accidental overdose, seek professional assistance or contact a Poison Control Center (1-800-222-1222) immediately. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed (see Overdose warning)
- measure only with dose cup provided and keep dose cup with product
- mL=milliliter
- do not exceed 4 doses per 24 hours

---

**Age**

adults & children 12 years and over  
children 6 to under 12 years  
children under 6 years

---

**Dose**

30 mL every 4 hours  
15 mL every 4 hours  
do not use

---

***Other information***

- **each 15 mL contains:**sodium 5 mg
- store at room temperature 15-30°C (59-86°C) and do not refrigerate


***Inactive ingredients***

Citric acid, FD&C Yellow No 6, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

***Questions or Comments?***

1-844-912-4012

TAMPER-EVIDENT FEATURE: DO NOT USE IF PRINTED  
NECK WRAP IS BROKEN OR MISSING.

  
0 71357 91244 5

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE  
If you are not completely satisfied with your BI-MART  
QUALITY PRODUCT, regardless of the reason, return the  
unused portion and your entire purchase price will be  
cheerfully refunded.

\*This product is not manufactured or  
distributed by The Procter & Gamble  
Company, distributor of Vicks®  
DayQuil® Cough & Flu.

Distributed by:  
BI-MART  
Eugene, OR 97402  
527-12-BM 04/25  
Rev. 1

no print / no varnish area  
lot no. & exp. date

BI-MART

NDC 37835-527-12

Compare to the active  
ingredients in Vicks® DayQuil®  
Cold & Flu\*

DAYTIME

COLD & FLU

Acetaminophen (Pain reliever/fever reducer)  
Dextromethorphan HBr (Cough suppressant)  
Phenylephrine HCl (Nasal decongestant)

Headache, Fever, Sore Throat,  
Minor Aches & Pains,  
Nasal Congestion,  
Cough

Ages 6 Years and Over

12 FL OZ (355 mL)

**Drug Facts**

**Active Ingredients** (In each 15 mL dose cup)  
Acetaminophen 325 mg.....Pain Reliever/Fever Reducer  
Dextromethorphan HBr 10 mg.....Cough Suppressant  
Phenylephrine HCl 5 mg.....Nasal Decongestant

**Uses** Temporarily relieves common cold/flu symptoms: • minor aches and pains • headache • sore throat • fever • nasal congestion • cough due to minor throat and bronchial irritation

**Warnings**  
**Live warnings:** This product contains acetaminophen. Severe liver damage may occur if:  
• adults and children over 12 years of age take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product  
• children 6 to 12 years of age take more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product  
• take with other drugs containing acetaminophen  
• adult has 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions.  
Symptoms may include: • skin redness • hives • rash. If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** • with any other drug containing acetaminophen (prescription or over-the-counter). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • if you have ever had an allergic reaction to this product or any of its ingredients.  
**Ask a doctor before use if you have** • liver disease • heart disease • high blood pressure • diabetes • thyroid disease • trouble urinating due to an enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, or emphysema.  
**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.  
**When using this product** • do not use more than directed (see overdose warning) • avoid alcoholic drinks.

DRUG FACTS CONTINUED ON THE BACK

**Purpose**

**Step use and ask a doctor if** • you get nervous, dizzy or drowsy • pain, nasal congestion or cough get worse or last more than 5 days (children) or 7 days (adults) • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding** ask a health professional before use.  
**Keep out of the reach of children.**  
**Overdose warning:** Taking more than the recommended dose (overdose) could cause serious health problems, including liver damage. In case of accidental overdose, seek professional assistance or contact a Poison Control Center (1-800-222-1222) immediately. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** • take only as directed (see overdose warning)  
• measure only with dose cup provided and keep dose cup with product  
• mL=milliliter • do not exceed 4 doses per 24 hours

Age	Dose
adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

**Other information** • Each 15 mL dose contains: sodium 5 mg  
• store at room temperature 15-30°C (59-86°F) and do not refrigerate

**Inactive ingredients** citric acid, FD&C Yellow #6, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum.

**Questions or comments?** 1-844-913-4012

# DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl liquid

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37835-527
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1VS297W6MV)	PHENYLEPHRINE	5 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID (UNII: 2968PHW8QP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

## Product Characteristics

Color	orange	Score	
-------	--------	-------	--

Shape		Size		
Flavor	ANISE	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-527-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		05/01/2025	

**Labeler -** BI-MART (027630078)

**Registrant -** BI-MART (027630078)

Establishment			
Name	Address	ID/FEI	Business Operations
Rnv LLC		118917568	manufacture(37835-527)