

ALL DAY BACK AND MUSCLE PAIN RELIEF- naproxen sodium tablet, coated
Walmart Inc.

Walmart Naproxen Sodium 220 mg Tablets

Drug Facts

Active ingredient (in each tablet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold

temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

you have difficulty swallowing

it feels like the pill is stuck in your throat

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

do not take more than directed
the smallest effective dose should be used

drink a full glass of water with each dose

-
- take 1 tablet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 tablets within the first hour
 - do not exceed 2 tablets in any 8- to 12-hour period
 - do not exceed 3 tablets in a 24-hour period
- Adults and children 12 years and older
- ask a doctor
- Children under 12 years
-

Other information

- **each tablet contains:** sodium 20 mg
- store at 20-25°C (68-77°F).
- avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide

Questions or comments? Call 1-877-290-4008

equate™ NDC 79903-609-20

All Day Back & Muscle Pain Relief
Naproxen Sodium
Tablets USP,
220 mg

Pain reliever/
Fever reducer
(NSAID)

• For temporary relief of
minor back and muscle
aches and pains

220 mg EACH **200 TABLETS**

12 HOURS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Uses
temporarily relieves minor aches and pains due to:
minor pain of arthritis, muscular aches, backache, menstrual cramps, headache, toothache, the common cold
temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed, **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.**

Directions do not take more than directed the smallest effective dose should be used drink a full glass of water with each dose. Adults and children 12 years and older: take 1 tablet every 8 to 12 hours while symptoms last for the first dose you may take 2 tablets within the first hour do not exceed 2 tablets in any 8- to 12-hour period do not exceed 3 tablets in a 24-hour period. Children under 12 years: ask a doctor.

Other information TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING
each tablet contains: sodium 20 mg store at 20-25°C (68-77°F), avoid high humidity and excessive heat above 40°C (104°F).

Questions or comments? Call 1-888-287-1915
Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments please call 1-888-287-1915.
DISTRIBUTED BY: Wal-Mart Inc., Bentonville, AR 72716
PRODUCT OF INDIA
60BR 1123 Code No. 60DRUGS615

Lot No.:
Exp Date:

equatē™

NDC 79903-609-20



All Day Back & Muscle Pain Relief

Naproxen Sodium Tablets USP, 220 mg

Pain reliever/
Fever reducer
(NSAID)

• For temporary relief of minor back and muscle aches and pains



220 mg EACH

200 TABLETS



40175901

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet) Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
 - temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Drug Facts (continued)

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years

- ask a doctor

Other information


- each tablet contains: sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide

Questions or comments? Call 1-888-287-1915

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an unclassified reaction or side effect, please call 1-888-287-1915.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716
PRODUCT OF INDIA
*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve® Tablets. 6081103



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING



Lot No.:
Exp Date:

equatē™

NDC 79903-609-09

All Day Back & Muscle Pain Relief

Naproxen Sodium Tablets USP, 220 mg

Pain reliever/
Fever reducer
(NSAID)

• For temporary relief of minor back and muscle aches and pains



220 mg EACH

90 TABLETS



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis, muscular aches, backache, menstrual cramps, headache, toothache, the common cold
 - temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Directions do not take more than directed, the smallest effective dose should be used, drink a full glass of water with each dose. Adults and children 12 years and older take 1 tablet every 8 to 12 hours while symptoms last, for the first dose you may take 2 tablets within the first hour, do not exceed 2 tablets in any 8- to 12-hour period, do not exceed 3 tablets in a 24-hour period. Children under 12 years ask a doctor.

Other information TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

- each tablet contains: sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F).

Questions or comments? Call 1-888-287-1915. Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments please call 1-888-287-1915.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716
PRODUCT OF INDIA
609R 1123 40175601
Code No.: GD0RUG6515

Lot No.:
Exp Date:



ALL DAY BACK AND MUSCLE PAIN RELIEF

naproxen sodium tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79903-609
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE (UNII: FZ989GH94E)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	141
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-609-09	1 in 1 CARTON	12/15/2023	
1		90 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:79903-609-20	1 in 1 CARTON	12/15/2023	
2		200 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090545	12/15/2023	

Labeler - Walmart Inc. (051957769)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(79903-609)