

**ASPIRIN- aspirin tablet, coated**  
**FSA Store Inc.**

-----  
**1018-CRM-2024-1016**

***Drug Facts***

**Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID)\*

\* nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

temporarily relieves:

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for:

- diabetes
- gout
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours

- children under 12 years: consult a doctor

**Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

**Inactive ingredients**

corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

**Questions or comments?**

1-844-705-4384

**PRINCIPAL DISPLAY PANEL**

NDC 81522-018-11

caring mill™

†Compare to the active ingredient of Bayer® Aspirin

Regular Strength

Aspirin

Pain Reliever / Fever Reducer

(NSAID)

Actual Size

100 COATED TABLETS 325 MG EACH



## ASPIRIN

aspirin tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:81522-018
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL;011
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81522-018-11	1 in 1 CARTON	11/16/2023	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/16/2023	

**Labeler** - FSA Store Inc. (049283340)

Revised: 10/2024

FSA Store Inc.