

REDICARE BLOOD CLOTTING- blood clotting spray aerosol, spray
Redicare LLC

Blood Clotting Spray

Active Ingredients

Benzethonium Chloride 0.2% w/w

Lidocaine 4% w/w

Purpose

Topical Antiseptic

Topical Antiseptic

Uses

- for help in the control of superficial bleeding
- for temporary relief of pain and to help protect against skin infection in minor cuts, minor scrapes, and minor burns

WARNINGS

For external use only.

Flammable

- Keep away from flame. Contents under pressure.
- Do not puncture or incinerate.
- Do not expose to temperatures above 120°(F)

DO NOT USE

- in eyes or other mucous membranes
- on large portions of the body
- in case of deep or puncture wounds
- on raw surfaces
- on blistered areas
- on animal bites
- on serious burns

DO NOT USE

- in eyes or other mucous membranes
- on large portions of the body
- in case of deep or puncture wounds
- on raw surfaces
- on blistered areas

- on animal bites
- on serious burns

Stop use and ask doctor if:

- Redness, swelling, or irritation occurs
- Infection occurs
- The condition persists or worsens
- Symptoms persist for more than 7 days

Stop use and ask doctor if:

- Redness, swelling, or irritation occurs
- Infection occurs
- The condition persists or worsens
- Symptoms persist for more than 7 days

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- shake well before using
- hold 6-8 inches from moist injured area
- spray until the area is covered
- Scab-like cover will gradually disappear as healing takes place or may be removed with soap and water
- for adult institutional use only
- not intended for use on children

Inactive Ingredients

Isobutane, Isopropyl Myristate, Karaya, N-Butane, Propane, Silica, Silicone, Sorbitan Monooleate, Talc, Tragacanth

202 x 406 - 2up)

Drug Facts

Active Ingredients
 Benzethonium Chloride 0.2% w/w.....Topical Antiseptic
 Lidocaine 4% w/w.....Topical Anesthetic

Purpose

Uses for help in control of superficial bleeding and temporary relief of pain of
 ■ minor cuts and scrapes ■ minor burns

Warnings
 For external use only

Flammable: ■ keep away from fire or flame ■ contents under pressure
 ■ do not puncture or incinerate container ■ do not expose to temperatures above 120°F

Do not use: ■ in or near mucous membranes ■ on serious burns ■ on raw surfaces
 ■ on deep puncture wounds ■ for a prolonged period of time ■ on blistered areas
 ■ on a large portion of the body ■ on animal bites

Stop use and ask a doctor if: ■ redness, swelling, or irritation occurs ■ infection occurs ■ condition persists or worsens ■ symptoms persist for more than 7 days

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
 ■ shake can well before using ■ hold can 6-8 inches from injured area
 ■ spray until area is covered ■ for adult use only ■ not intended for use on children
 ■ scab-like cover will gradually disappear as healing takes place or may be removed with soap and water

Other Information
 ■ avoid inhaling ■ use only as directed
 ■ intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal

Inactive Ingredients
 Isobutane, Isopropyl Myristate, Karaya, n-Butane, Propane, Silica, Silicone, Sorbitan Monooleate, Talc, Tragacanth

FLAMMABLE CONTENTS UNDER PRESSURE

Redicare LLC
 Congers, NY 10920
 866.561.5650

CAT 0105010
 NDC 71105-107-03
 info@redicare.com
 www.redicare.com



4 552701 050106



Blood Clotting Spray

Blood Stopper Spray

Helps control superficial bleeding

Net Wt. 3 oz. (85g)

REDICARE BLOOD CLOTTING

blood clotting spray aerosol, spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71105-107
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZ ETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	170 mg in 85 g
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	3400 mg in 85 g

Inactive Ingredients

Ingredient Name	Strength
3-(TRIETHOXYSILYL)PROPYLAMINE (UNII: L8S6UBW552)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
PROPANE (UNII: T75W9911L6)	
TRAGACANTH (UNII: 2944357O2O)	
TALC (UNII: 7SEV7J4R1U)	
ISOBUTANE (UNII: BXR49TP611)	
BUTANE (UNII: 6LV4FOR43R)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	

KARAYA GUM (UNII: 73W9IQY50Q)

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71105-107-03	85 g in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	10/01/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	10/01/2023	

Labeler - Redicare LLC (800149346)

Revised: 8/2025

Redicare LLC