

**MOTRIN IB- ibuprofen tablet, film coated**  
**Lil' Drug Store Products, Inc.**

-----  
**Motrin IB®**

***Drug Facts***

**Active ingredient (in each caplet)**

Ibuprofen 200 mg (NSAID)<sup>1</sup>

---

<sup>1</sup> nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

#### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

#### **When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

#### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**

- 
- take 1 caplet every 4 to 6 hours while symptoms

adults and children 12 years and older	<p>persist</p> <ul style="list-style-type: none"> <li>▪ if pain or fever does not respond to 1 caplet, 2 caplets may be used</li> <li>▪ do not exceed 6 caplets in 24 hours, unless directed by a doctor</li> </ul>
children under 12 years	<ul style="list-style-type: none"> <li>▪ ask a doctor</li> </ul>

### Other information

- store between 20 - 25°C (68 - 77°F)
- **do not use if carton or pouch is open or torn**
- see bottom panel for lot number and expiration date

### Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide

### Questions or comments?

Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST

**Repackaged and distributed by:** Lil' Drug Store Products, Inc.  
1201 Continental Place NE, Cedar Rapids, IA 52402

### PRINCIPAL DISPLAY PANEL - 4 Caplet Carton

*See New Warnings Information*

**Motrin IB<sup>®</sup>**

***Ibuprofen Tablets USP, 200 mg  
Pain Reliever/Fever Reducer (NSAID)***

***\*Capsule-Shaped Tablets***

***4 COATED  
CAPLETS\****

***Lil'  
Drug Store<sup>®</sup>***

See New Warnings Information

# Motrin<sup>®</sup> IB

Ibuprofen Tablets USP, 200 mg  
Pain Reliever / Fever Reducer (NSAID)



\*Capsule-Shaped Tablets

4 COATED  
CAPLETS\*



Code 128  
1464

4 COATED  
CAPLETS\*

Motrin<sup>®</sup> IB

Important: Read all product information before using.  
Keep this box for important information.

### Drug Facts

#### Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)\*.....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

#### Purpose

#### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - menstrual cramps
  - temporarily reduces fever
  - muscular aches
  - backache
- minor pain of arthritis
- the common cold

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

#### Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

LIFT PANEL FOR CONTINUED DRUG FACTS

### Drug Facts (continued)

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

#### Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

#### When using this product

- take with food or milk if stomach upset occurs


### Drug Facts (continued)

- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.**



**Drug Facts (continued)**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older	<ul style="list-style-type: none"> <li>■ take 1 caplet every 4 to 6 hours while symptoms persist</li> <li>■ if pain or fever does not respond to 1 caplet, 2 caplets may be used</li> <li>■ do not exceed 6 caplets in 24 hours, unless directed by a doctor</li> </ul>
children under 12 years	<ul style="list-style-type: none"> <li>■ ask a doctor</li> </ul>

**Other information**

- store between 20 - 25°C (68 - 77°F)
- do not use if carton or pouch is open or torn
- see bottom panel for lot number and expiration date

**Inactive ingredients**    carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6, hypromellose, iron oxide, magnesium stearate, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide

**Questions or comments?**

Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST

Manufactured by: McNeil Healthcare LLC, LAS PIEDRAS, PUERTO RICO 00771  
 For: McNeil Consumer Healthcare, DIVISION OF MCNEIL-PPC, INC.  
 FORT WASHINGTON, PA 19034 USA ©MCN-PPC, Inc. 2010 www.motrin.com

Repackaged and distributed by: Lif' Drug Store Products, Inc.  
 1201 Continental Place NE, Cedar Rapids, IA 52402  
 1-877-507-8516 (M-F 8AM-4:30PM CST) www.lifdrugstore.com Rev 3-10 v3

EXP  
LOT



**MOTRIN IB**

ibuprofen tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-9712(NDC:50580-110)
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>Ibuprofen</b> (UNII: WK2XYI10QM) (Ibuprofen - UNII:WK2XYI10QM)	Ibuprofen	200 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>carnauba wax</b> (UNII: R12CBM0EIZ)	
<b>silicon dioxide</b> (UNII: ETJ7Z6XBU4)	
<b>starch, corn</b> (UNII: O8232NY3SJ)	

<b>FD&amp;C yellow no. 6</b> (UNII: H77VEI93A8)
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)
<b>ferric oxide red</b> (UNII: 1K09F3G675)
<b>magnesium stearate</b> (UNII: 70097M6I30)
<b>polydextrose</b> (UNII: VH2XOU12IE)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)
<b>propylene glycol</b> (UNII: 6DC9Q167V3)
<b>shellac</b> (UNII: 46N107B71O)
<b>stearic acid</b> (UNII: 4ELV7Z65AP)
<b>titanium dioxide</b> (UNII: 15FIX9V2JP)

**Product Characteristics**

<b>Color</b>	ORANGE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	Motrin;IB
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-9712-1	1 in 1 CARTON	05/10/2010	04/27/2021
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:66715-9712-2	2 in 1 CARTON; Type 0: Not a Combination Product	05/10/2010	04/27/2021

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019012	05/10/2010	04/27/2021

**Labeler - Lil' Drug Store Products, Inc. (093103646)**