

MOORE MEDICAL BISMUTH- bismuth subsalicylate tablet, chewable
Moore Medical LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Moore Medical Bismuth

Drug Facts

Active ingredient (in each tablet)

Bismuth Subsalicylate 262mg

(each tablet contains 102 mg salicylate)

Purpose

Upset stomach reliever/antidiarrheal

Uses

Temporarily relieves

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - belching
 - fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are:

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- bloody or black stool
- an ulcer
- a bleeding problem

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist if you are taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

When using this product

a temporary and harmless darkening of the tongue and/or stool may occur. Stool darkening should not be confused with melena.

If pregnant or breast-feeding,

ask a health professional before use.

Keep this and all drugs out of reach of children.

In case of accidental overdose, contact a physician or poison control center immediately.

Directions

- ■ **do not use more than directed**
 - **chew or crush tablets completely before swallowing**
 - **do not swallow tablets whole**
 - use until diarrhea stops but not more than 2 days
 - drink plenty of clear fluids to help prevent dehydration, which may accompany diarrhea
 - do not exceed 16 tablets in 24 hours

Adults and children: (12 years and older) Take 2 tablets every 1/2 to 1 hour or 4 tablets every hour as needed

Children under 12 years: Do not give to children under 12 years of age.

Other information

- phenylketonurics: contains phenylalanine 1.1mg per tablet
- each tablet contains 73mg elemental calcium
- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

acacia gum, aspartame, calcium carbonate, D&C red #27, dextrans, flavoring, magnesium stearate, maltodextrin, microcrystalline cellulose, silicon dioxide

Questions or comments?

Call 800.234.1464

Moore Medical Bismuth Label

moore medical

Supporting Health and Care

Reorder No. 82473

UPN# 0-0607415-82473-5

Moore Brand®

Bismuth Tablets

100 Packets/2 Tablets

Controls diarrhea within 24 hours,
relieving associated abdominal cramps;
provides soothing relief for:
indigestion, upset stomach, heartburn and nausea.

Bismuth Subsalicylate 262 mg

Chew or crush tablets completely before swallowing.

Do not swallow tablets whole.

To Reorder:

Call: 800.234.1464

Fax: 800.944.6667

Online: www.mooremedical.com

Retain carton for complete product information

mooremedical
Supporting Health & Care

Drug Facts	
Active ingredient (in each tablet) Bismuth Subsalicylate 262 mg (each tablet contains 102 mg salicylate)	Purpose Upset stomach reliever/antidiarrheal
Uses	
Temporarily relieves <ul style="list-style-type: none"> ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: <ul style="list-style-type: none"> ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness 	
Warnings	
<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.</p> <p>Allergy alert: Contains salicylate. Do not take if you are: ■ allergic to salicylates (including aspirin) ■ taking other salicylate products</p> <p>Do not use if you have ■ bloody or black stool ■ an ulcer ■ a bleeding problem</p> <p>Ask a doctor before use if you have ■ fever ■ mucus in the stool</p> <p>Ask a doctor or pharmacist if you are taking any drug for ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis</p> <p>Stop use and ask a doctor if ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days</p> <p>When using this product a temporary and harmless darkening of the tongue and/or stool may occur. Stool darkening should not be confused with melena.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep this and all drugs out of reach of children. In case of accidental overdose, contact a physician or poison control center immediately.</p>	
Directions	
<ul style="list-style-type: none"> ■ do not use more than directed ■ chew or crush tablets completely before swallowing ■ do not swallow tablets whole ■ use until diarrhea stops but not more than 2 days ■ drink plenty of clear fluids to help prevent dehydration, which may accompany diarrhea ■ do not exceed 16 tablets in 24 hours 	
Adults and children: (12 years and older)	Chew 2 tablets every 1/2 to 1 hour or 4 tablets every hour as needed
Children under 12 years:	Do not give to children under 12 years of age.
Other Information:	
<ul style="list-style-type: none"> ■ phenylketonurics: contains phenylalanine 1.1mg per tablet ■ each tablet contains 73mg elemental calcium ■ store at room temperature 59°-86°F (15°-30°C) ■ tamper-evident sealed packets ■ do not use any opened or torn packets 	
Inactive ingredients acacia gum, aspartame, calcium carbonate, D&C red #27, dextrates, flavoring, magnesium stearate, maltodextrin, microcrystalline cellulose, silicon dioxide	
Questions or Comments? Call 800.234.1464	

Reorder No. 82473

UPN# 0-0607415-82473-5

MooreBrand® Bismuth Tablets

100 Packets/2 Tablets

**Controls diarrhea within 24 hours,
relieving associated abdominal cramps;
provides soothing relief for: indigestion,
upset stomach, heartburn and nausea.**

Bismuth Subsalicylate 262 mg

**Chew or crush tablets completely before swallowing.
Do not swallow tablets whole.**

To Reorder:
Call: 800.234.1464
Fax: 800.944.6667
Online: www.mooremedical.com

MOORE MEDICAL BISMUTH

bismuth subsalicylate tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55670-474
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD18YE)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
ASPARTAME (UNII: Z0H242BBR1)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	

DEXTRATES (UNII: G263MI44RU)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	16mm
Flavor		Imprint Code	RH;046
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55670-474-47	100 in 1 BOX	04/01/2014	07/31/2024
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	04/01/2014	07/31/2024

Labeler - Moore Medical LLC (051420107)

Revised: 3/2022

Moore Medical LLC