

**TABCIN EXTRA STRENGTH- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl, capsule, liquid filled  
Pharmadel LLC**

-----  
**Tabcin Antigrip Softgel (HHH)(DA)**

**Drug Facts**

**Active Ingredients & Purposes**

- Acetaminophen 250mg
- Chlorpheniramine maleate 2 mg
- Dextromethorphan hydrobromide 10 mg
- Phenylephrine hydrochloride 5 mg

---

<b>Active Ingredients</b>	<b>Purposes</b>
Acetaminophen 250mg.....	Pain reliever/ fever reducer
Chlorpheniramine maleate 2 mg.....	Antihistamine
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Phenylephrine hydrochloride 5 mg.....	Nasal decongestant

---

**Uses**

For the temporary relief of symptoms due to the common cold/ flu:

- runny nose
- sneezing
- cough due to minor bronchial irritation
- nasal congestion
- minor aches and pain
- headache
- sore throat
- and to reduce fever

**Warnings**

**Liver warning**

This product contains **acetaminophen**. Severe liver damage may occur if you take

- more than **10 softgels in 24 hours**, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, and is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- a persistent or chronic cough such as occurs with smoking, asthma, or emphysema, or if cough is accompanied by excessive phlegm (mucus) unless directed by a doctor

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed the recommended dosage**
- may cause excitability, especially in children
- may cause drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness effect □avoid alcoholic beverages while taking this product
- use caution when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occurs
- the pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

- cough comes back or occurs with a rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **adults and children 12 years and over:** take 2 softgels with water every 4 hours. Do not exceed 10 softgels in 24 hours or as directed by a doctor.
- **children under 12 years:** do not use

***Other information***

- store at room temperature 15°-30°C (59°F-86°F)
- avoid excessive heat

***Inactive ingredients***

FD&C blue no.1, gelatin, glycerin, polyethylene glycol 400, povidone k30, propylene glycol, propylparaben, methylparaben, sodium hydroxide, sorbitol, titanium dioxide, water

***Questions or comments***

**1-866-359-3478** (M-F) 9 AM to 5 PM EST or [www.pharmadel.com](http://www.pharmadel.com)

**Principal Display Panel**

NDC 55758-407-12

# Antigripal **tabcin**™

## **EXTRA STRENGTH COUGH & COLD**

- *Fever*
- *Nasal Congestion*
- *Muscle and Backaches*
- *Runny Nose*



**Acetaminophen, Chlorpheniramine maleate,  
Dextromethorphan HBr, Phenylephrine HCl**

**12 Softgels**

### **TABCIN EXTRA STRENGTH**

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl, capsule, liquid filled

#### **Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55758-407
<b>Route of Administration</b>	ORAL		

#### **Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	

## Product Characteristics

<b>Color</b>	blue	<b>Score</b>	score with uneven pieces
<b>Shape</b>	CAPSULE (Oblong)	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	SN5
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55758-407-12	1 in 1 CARTON	10/02/2023	08/01/2027
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/02/2023	08/01/2027

**Labeler** - Pharmadel LLC (030129680)

Revised: 12/2025

Pharmadel LLC