

**DOUBLE TUSSIN INTENSE COUGH RELIEVER DM- dextromethorphan hydrobromide /
guaifenesin liquid**

Reese Pharmaceutical Co

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DRUG FACTS

Active ingredient per 5ml teaspoon

Dextromethorphan Hydrobromide 20mg Cough Suppressant

Guaifenesin 300mg Expectorant

Uses. temporarily relieves cough due to minor throat and bronchial irritation as may occur with a common cold • helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus
• helps make coughs more productive

Do not use if you are now taking a prescription monoamine oxidase (inhiMor~MAIO)

Certain drugs for depression, psychiatric or emotional conditioners or Parkinson's disease

or for 2 weeks after stopping MAIO drug, If you do not know if your prescription drug contains an MAIO, ask your doctor or pharmacist before using this product

Ask doctor before use if you have

persistent or chronic cough, such as occurs with smoking, asthma, bronchitis or emphysema

cough is accompanied by excessive phlegm (mucous)

Stop use and ask doctor if

Symptoms are accompanied by fever, rash or persistent headache

cough persists for more than 1 week or tends to recur

A persistent cough may be a sign of a serious condition

If pregnant or breastfeeding, ask a health professional before use

Keep out of reach of children, In case of overdose, get medical help or contact a Poison Center immediately

Directions

• Adults and children 12 years of age and over:

take 1 tablet every 4 hours as needed

• Children 6 to 10 under 12 years of age: take 1/2 tablet every 4 hours as needed

• Children under 6 years of age: consult a doctor

Do not exceed 6 doses in a 24 hour period or as directed by a doctor

Other information store at 15°- 30° C (59°- 86°F)

Purpose

Dextromethorphan Hydrobromide Cough Suppressant

Guaifenesin Expectorant

Cherry Flavor, Citric Acid, Menthol, Polyethylene Glycol, Propylene Glycol, Purified Water,

Sodium Benzoate, Sodium Saccharin, Sorbitol.

NDC 10956-674-04

Double DM Tussin

INTENSE
STRENGTH

DOUBLE STRENGTH

Intense Cough Reliever

- Sugar Free
- Alcohol Free
- Non-Drowsy
- New Improved Taste!

COUGH
SUPPRESSANT
EXPECTORANT

FREE
DOSAGE
CUP

4 FL oz (120 ml)

Stronger Than
Leading Brands

CAUTION: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

Drug Facts

Active ingredients **Purpose**
(in each 5ml teaspoon)
Dextromethorphan HBr 20mg.....Cough Suppressant
Guaifenesin 300mg.....Expectorant

Uses
■ Temporarily relieves cough due to minor throat and bronchial irritation as may occur with a common cold
■ Helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus.
■ Helps make coughs more productive.

Warnings
Do not use ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before using this product.
Ask a doctor before use if you have
■ persistent or chronic cough, such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough is accompanied by excessive phlegm (mucus)

Stop use and ask a doctor if
■ symptoms are accompanied by fever, rash, or persistent headache
■ cough persists for more than 1 week or tends to recur
A persistent cough may be a sign of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions
■ Adults and children 12 years of age and over take 1 teaspoon every 4 hours as needed
■ Children 6 to under 12 years take 1/2 teaspoon every 4 hours as needed
■ Children under 6 consult a doctor
Do not exceed 6 doses in a 24 hour period or as directed by a doctor.

Other information
■ store between 15°-30° C (59°-86°F)

Inactive ingredients Cherry Flavor, Citric Acid, Menthol, Polyethylene Glycol, Propylene Glycol, Purified Water, Sodium Benzoate, Sodium Saccharin, Sorbitol.

PRECAUCIÓN: NO LO USE SI EL SELLO IMPRESO BAJO LA TAPA ESTÁ ROTO O AUSENTE.

Hechos Acerca Del Medicamento

Ingredientes activos **Propósito**
(en cada cucharadita de 5ml)
Bronhidrato de dextrometorfano 20mg..... Supresor de las tos
Guaifenesina 300mg..... Expectorante

Usos
■ Ayuda a atajar la flema (mucosidad) a afinar las secreciones bronquiales para liberar las vías bronquiales de la molesta mucosidad
■ hacer que la tos sea más productiva
■ Alivia temporariamente la tos producida por resfriados comunes.

Advertencias
No tome
■ este producto si actualmente está tomando medicamentos de prescripción que contienen el inhibidor de Oxidasa Monoamina (MAOI) (ciertos medicamentos para la depresión, estados psiquiátricos o emocionales o Mal de Parkinson), o durante las 2 semanas siguientes a haber dejado de tomar un medicamento con MAOI, consulte a un profesional de la salud.
Consulte a un médico antes de usarlo si tiene
■ la tos persistente o crónica como la que ocurre al fumar, con el asma, bronquitis crónica o enfisema
■ la tos se acompaña de flema excesiva (moco)

Deje de tomar y consulte a su médico
■ los síntomas se acompañan de fiebre, erupción cutánea o dolor de cabeza persistente
■ la tos persiste por más de 1 semana o tiende a reaparecer.
Una tos persistente pueda ser signo de un trastorno serio.
Si está embarazada o dando pecho, consulte a un profesional de salud antes de tomar este medicamento. Manténgalo lejos del alcance de los niños. En caso de una sobredosis, consiga asistencia médica inmediatamente.

Hechos Acerca Del Medicamento (cont.)

Instrucciones
■ Adultos y niños mayores de 12 años tome 1 cucharadita cada 4 horas como sea necesario.
■ Niños de 6 a 12 años tome 1/2 cucharadita cada 4 horas como sea necesario.
■ Niños menores de 6 años consulte a su Médico como sea necesario.
No exceda más de 6 dosis durante un periodo de 24 horas o como se lo indique su Médico.

Otra Información
■ Guarde a la temperatura ambiente de 15°-30°C (59°-86°F)

Ingredientes inactivos Sabor de cherry, Ácido cítrico, mentol, polietilenglicol, propilenglicol, agua purificada, benzoato sódico, sacarinato sódico, sorbitol.

Rev. 07/10

Mas Fuerte
Que Las Marcas
Principales

DISTRIBUTED BY

REESE
PHARMACEUTICAL
COMPANY

CLEVELAND, OH 44106
1-800-321-7178
reese@apk.net
www.reese-pharmaceutical.com

OUR GUARANTEE

This product is satisfaction
guaranteed or your money back

0 23513 67404 9

DOUBLE TUSSIN INTENSE COUGH RELIEVER DM

dextromethorphan hydrobromide / guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:10956-674
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 5 mL
Guaifenesin (UNII: 495W7451VQ) (Guaifenesin - UNII:495W7451VQ)	Guaifenesin	300 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10956-674-01	1 in 1 CARTON		
1	NDC:10956-674-04	120 mL in 1 BOTTLE, PLASTIC		
2	NDC:10956-674-08	240 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/02/2011	

Labeler - Reese Pharmaceutical Co (004172052)

Registrant - Reese Pharmaceutical Co (004172052)

Establishment

Name	Address	ID/FEI	Business Operations
Reese Pharmaceutical Co		004172052	relabel, repack

Revised: 1/2011

Reese Pharmaceutical Co