

MEDIQUE DIAMODE- loperamide hydrochloride tablet
PRIVATE LABEL DISTRIBUTOR- loperamide hcl 2 mg tablet
Unifirst First Aid Corporation

MEDIQUE[®] Diamode

Drug Facts

Active ingredient (in each caplet)

Loperamide Hydrochloride 2mg

Purpose

Antidiarrheal

Uses

Controls the symptoms of diarrhea, including Travelers diarrhea.

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl

Do not use

- if you have bloody or black stool

Ask a doctor before use if you have

- a fever
- mucus in stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are

- taking antibiotics

When using this product

- tiredness, drowsiness or dizziness may occur
- be careful when driving or operating machinery

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If you are pregnant or breast feeding, ask a health professional before use.

Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a physician or Poison Control Center right away.

Directions

- do not use more than directed
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Adults and children (12 years and older): Take 2 caplets after the first loose stool followed by 1 caplet after each subsequent loose stool but no more than 4 caplets in 24 hours.

Children under 12 years: Do not give to children under 12 years of age.

Other information

- store between 68° - 77°F (20° - 25°C)
- tamper-evident sealed packets
- do not use any opened or torn packet

Inactive ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow # 10, FD&C Blue # 1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Questions or comments? 1-800-634-7680

Principal Display Panel - 200R Medique Diamode Label

Medique®

Diamode

Controls the Symptoms of Diarrhea

Pull to Open

Antidiarrheal • Loperamide HCl 2 mg

50 Caplets

(50 x 1)

Tamper Evident Unit Dose Packets

<p>Drug Facts</p> <p>Active ingredient (in each caplet) Loperamide Hydrochloride 2mg</p> <p>Purpose Antidiarrheal</p> <p>Uses Controls the symptoms of diarrhea, including Traveler's diarrhea.</p> <p>Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl. Do not use if you have bloody or black stool. Ask a doctor before use if you have a fever, mucus in stool, or a history of liver disease. Ask a doctor or pharmacist before use if you are taking antibiotics.</p> <p>When using this product Drowsiness, dizziness, or dizziness may occur. Be careful when driving or operating machinery.</p> <p>Stop use and ask a doctor if symptoms get worse. diarrhea lasts more than 2 days. you get abdominal swelling or bulging. These may be signs of a serious condition. If you are pregnant or breast feeding, ask a health professional before use.</p>	<p>Medique®</p> <p>Diamode</p> <p>Controls the Symptoms of Diarrhea</p> <p>Pull to Open</p>	<p>Drug Facts (continued)</p> <p>Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a physician or Poison Control Center right away.</p> <p>Directions do not use more than directed. drink plenty of clear fluids to help prevent dehydration caused by diarrhea.</p> <p>Adults and children (12 years and older): Take 2 caplets after the first loose stool followed by 1 caplet after each subsequent loose stool but no more than 4 caplets in 24 hours.</p> <p>Children under 12 years: Do not give to children under 12 years of age.</p> <p>Other information store at room temperature 68° - 77°F (20° - 25°C). tamper-evident sealed packets. do not use any opened or torn packet.</p> <p>Inactive ingredients anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow # 10, FD&C Blue # 1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch.</p> <p>Questions or comments? 1-800-634-7680</p>	<p>Medique®</p> <p>Diamode</p> <p>Controls the Symptoms of Diarrhea</p> <p>Reorder #20050 Manufactured for Medique Products Fort Myers, FL 33967 www.mediqueproducts.com 1-800-634-7680</p> <p>3 47682 20050 3</p>
---	---	---	--

Retain carton for complete product information

Antidiarrheal • Loperamide HCl 2mg

50 Caplets (50 x 1)

Tamper Evident Unit Dose Packets

Private Label Distributor Physicians Care Anti-Diarrheal

Physicians Care®

Anti-Diarrheal

Controls Symptoms of Diarrhea

Loperamide Hydrochloride 2 mg

Pull To Open

Easy to Swallow

COATED CAPLETS

Compare Active ingredient to:

Imodium AD

Registered Trademark of McNeil Consumer & Specialty Pharmaceuticals

Tamper-Evident Unit Dose Packets

Physicians Care®

Registered Trademark of ACME UNITED Corporation

50 PACKETS (1 Caplet 2 Mg Each)

Also Available from PhysiciansCare®
in individual dose packets:

- PhysiciansCare® Aspirin
- PhysiciansCare® Ibuprofen
- PhysiciansCare® Non-Aspirin (Acetaminophen)
- PhysiciansCare® Antacid
- PhysiciansCare® Non-Drowsy Sinus
- PhysiciansCare® Allergy
- PhysiciansCare® Cold & Cough

PhysiciansCARE®

Anti-Diarrheal

Controls Symptoms of Diarrhea

Loperamide Hydrochloride 2 mg

Pull To Open

Easy to Swallow
COATED CAPLETS

Compare active ingredient to:
Imodium® AD
Registered Trademark of McNeil Consumer & Specialty Pharmaceuticals

Tamper-Evident Unit Dose Packets

Physicians Care®
Registered Trademark of ACME UNITED Corporation

50 PACKETS (1 Caplet 2 Mg Each)

Drug Facts

Active ingredient (in each caplet) **Purpose**
Loperamide Hydrochloride 2 mg Antidiarrheal

Uses
Controls the symptoms of diarrhea, including Travelers diarrhea.

Warnings
Allergy Alert: Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl

Do not use • if you have bloody or black stool

Ask a doctor before use if you have • a fever • mucus in stool • a history of liver disease

Ask a doctor or pharmacist before use if you are • taking antibiotics

When using this product
• tiredness, drowsiness or dizziness may occur
• be careful when driving or operating machinery

Stop use and ask a doctor if
• symptoms get worse
• diarrhea lasts more than 2 days
• you get abdominal swelling or bulging. These may be signs of a serious condition.

If you are pregnant or breast feeding, ask a health professional before use.

Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a physician or Poison Control Center right away.

Drug Facts (continued)

Directions
• do not use more than directed
• drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Adults and children: (12 years and older)
Take 2 caplets after the first loose stool followed by 1 caplet after each subsequent loose stool but no more than 4 caplets in 24 hours.

Children under 12 years:
Do not give to children under 12 years of age.

Other Information
• store at room temperature 68° - 77°F (20° - 25°C)
• tamper-evident sealed packets
• do not use any opened or torn packet

Inactive ingredients
Anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow # 10, FD&C Blue # 1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Questions or Comments?
Call 1-800-835-2263

Distributed by Acme United Corporation,
Fairfield, CT 06824
1-800-835-2263 www.acmeunited.com

Retain carton for complete product information

PhysiciansCARE®

Anti-Diarrheal

Controls Symptoms of Diarrhea

Easy to Swallow
COATED CAPLETS

Compare active ingredient to:
Imodium® AD
Registered Trademark of McNeil Consumer & Specialty Pharmaceuticals

Tamper-Evident Unit Dose Packets

Physicians Care®
Registered Trademark of ACME UNITED Corporation

50 PACKETS (1 Caplet 2 Mg Each)

Compare to the Active Ingredients in

Imodium® A-D

This box contains a generic product called Diamode

Anti-Diarrheal Relief

20 Packets of 2 Tablets

Remove This Panel

Read Drug Facts and Instructions on Packet

Drug Facts
Active ingredient (in each caplet) Purpose Loperamide Hydrochloride 2mg Antidiarrheal
Uses Controls the symptoms of diarrhea, including Travelers diarrhea
Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl. Do not use • if you have bloody or black stool Ask a doctor before use if you have • a fever • mucus in stool • a history of liver disease Ask a doctor or pharmacist before use if you are • taking antibiotics
When using this product • tiredness, drowsiness or dizziness may occur • be careful when driving or operating machinery
Stop use and ask a doctor if • symptoms get worse • diarrhea lasts more than 2 days • you get abdominal swelling or bulging. These may be signs of a serious condition. If you are pregnant or breast feeding , ask a health professional before use. Keep this and all drugs out of the reach of children . In case of accidental overdose, contact a physician or Poison Control Center right away.
Directions • do not take more than directed • drink plenty of clear fluids to help prevent dehydration caused by diarrhea Adults and children: (12 years and older) Take 2 caplets after the first loose stool followed by 1 caplet after each subsequent loose stool but no more than 4 caplets in 24 hours. Children under 12 years: Do not give to children under 12 years of age.
Other information • store between 68° and 77°F (20° and 25°C) • tamper-evident sealed packets • do not use any opened or torn packet
Inactive ingredients anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow # 10, FD&C Blue # 1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch
Questions or comments? 1-888-753-6575

Compare to the Active Ingredient in
Imodium[®] A-D

Anti-Diarrheal
Relief

20 Packets
of 2 Tablets

Compare to the Active Ingredients in
Imodium[®] A-D

This box contains a generic product called Diamode

Anti-Diarrheal
Relief

MEDIQUE DIAMODE

loperamide hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-200
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSPROVIDONE (UNII: 68401960MK)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

Product Characteristics

Color	green (green)	Score	2 pieces
Shape	OVAL (Caplet)	Size	10mm
Flavor		Imprint Code	123
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-200-69	6 in 1 BOX, UNIT-DOSE	12/30/2008	
1		1 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-200-50	50 in 1 BOX, UNIT-DOSE	12/30/2008	
2		1 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-200-33	100 in 1 BOX, UNIT-DOSE	12/30/2008	

3	NDC:47682-200-46	1 in 1 PACKET; Type 0: Not a Combination Product	
4	NDC:47682-200-46	1 in 1 PACKET; Type 0: Not a Combination Product	12/30/2008
5	NDC:47682-200-64	24 in 1 BOX, UNIT-DOSE	12/30/2008
5	NDC:47682-200-46	1 in 1 PACKET; Type 0: Not a Combination Product	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074091	12/30/2008	

PRIVATE LABEL DISTRIBUTOR

loperamide hcl 2 mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-021
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
CROSPVIDONE (UNII: 2S7830E561)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	green	Score	2 pieces
Shape	OVAL (Caplet)	Size	10mm
Flavor		Imprint Code	123
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-021-50	50 in 1 BOX, UNIT-DOSE	12/30/2008	12/31/2024
1		1 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-021-25	25 in 1 BOX, UNIT-DOSE	10/01/2015	12/31/2024
2		1 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-021-30	30 in 1 BOX, UNIT-DOSE	10/01/2015	12/31/2024
3		1 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-021-75	75 in 1 BOX, UNIT-DOSE	10/01/2015	12/31/2024
4		1 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074091	12/30/2008	

Labeler - Unifirst First Aid Corporation (832947092)

Revised: 4/2025

Unifirst First Aid Corporation