ADULT LOW DOSE ASPIRIN- aspirin tablet, delayed release FELICIFIC INC.

Adult Low Dose Aspirin Delayed Release Tablets, 81 mg

Drug Facts

Active ingredients (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

• temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed release action, this product will not provide fast relief of headaches or symptoms needing immediate relief.

Warnings

Allergy alert:

Aspirin may cause a severe allergic reaction which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

Stomach bleeding warning:

- This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

• if you are allergic to aspirin or any other pain reliever/fever reducer

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior

with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- under a doctor's care for any serious condition
- taking any other drug

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- an allergic reaction occurs.

Seek medical help right away.

- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling present in the painful area
- new symptoms occur

These could be sign of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - $\circ~$ take 4 to 8 tablets every 4 hours while symptoms last
 - o do not take more than 48 tablets in 24 hours unless directed by a doctor
- children under 12 years:ask a doctor

Storage Conditions

- store at room temperature 20-25°C (68-77°F)
- TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Inactive ingredients

Starch, Povidone, Microcrystalline Cellulose, Croscarmellose Sodium, Stearic Acid, Colloidal Silicon Dioxide, Methacrylic Acid and Ethyl acrylate Co-polymer, Triethyl Citrate, Talc, Fumed Silica, Titanium Dioxide, D&C Yellow #10, FD&C Yellow #6.

Questions or comments?

Please contact us at info@heivy.com

Distributed by:

Heivy ® of Felicific Inc

Queens, NY 11435

www.heivy.com

Made in the USA

PRINCIPAL DISPLAY PANEL

NDC 84014-531-03

Heivy ®

Compare to Bayer ®Low Dose Aspirin active ingredients*

Adult Low Dose Aspirin Delayed Release Tablets, 81 mg

Pain Reliever (NSAID)

Enteric Coated

300 Tablets

*This product is not manufactured or distributed by Bayer HealthCare LLC., owner of the registered trademark Bayer $^{\$}$ Low Dose Aspirin.



Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for

children even if you do not notice any signs or symptoms.

■ take 4 to 8 tablets every 4 hours while symptoms last ■ do not take more than 48 tablets in 24 hours

children under 12 years: ask a doctor

unless directed by a doctor

drink a full glass of water with each dose adults and children 12 years and over:

Directions

Storage Conditions

store at room temperature 20-25°C (68-77°F)

Starch, Povidone, Microcystalline cellulose, Croscarmellose sodium. Stearic acid. Collodal alion dioxide, Methacrylir Add and Ethyl acrystate co-colymer. Triethyl diztate, Tale. Furned silica, Titanium dioxide, D.& C. yellow #10, FD & C. yellow #6.

Inactive ingredients:

Questions or comments? Please contact us at info@heivy.com

not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

ask a health professional before use. It is especially importan

if pregnant or breast-feeding, if pregnant or breast-feeding,

These could be signs of a serious condition.

new symptoms occur

stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn

Drug Facts (Continued)

Ask a doctor before use if

you have asthma

you have high blood pressure, heart disease, liver cirrhosis

you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit

Stop use and ask a doctor if

taking any other drug

blood ■ have stomach pain that does not get better

■ pain gets worse or lasts for more than 10 days
■ fever gets worse or lasts for more than 3 days
■ redness or swelling present in the painful area

■ringing in the ears or loss of hearing occurs

Seek medical help right away.

■an allergic reaction occurs.

■ taking a prescription drug for diabetes, gout, or arthritis ■ under a doctor's care for any serious condition

ADULT LOW DOSE ASPIRIN

Ask a doctor or pharmacist before use if you are

you are taking a diuretic

or kidney disease

aspirin tablet, delayed release **Product Information Product Type HUMAN OTC DRUG** Item Code (Source) NDC:84014-531 **Route of Administration ORAL Active Ingredient/Active Moiety Basis of Strength Ingredient Name** Strength ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) **ASPIRIN** 81 mg **Inactive Ingredients Ingredient Name** Strength STARCH, CORN (UNII: O8232NY3SJ)

POVIDONE (UNII: FZ 989GH94E)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics			
Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	A1
Contains			

Ш	Packaging			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:84014-531- 03	300 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2024	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2024	

Labeler - FELICIFIC INC. (081178920)

Establishment				
Name	Address	ID/FEI	Business Operations	
Health Pharma USA LLC		080804485	manufacture(84014-531)	

Revised: 5/2024 FELICIFIC INC.