

**COLD AND FLU RELIEF NIGHTTIME- acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose

Pain reliever/Fever Reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists more than 2 days, is

accompanied or followed by fever, headache rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do

not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

Other information

- store between 15^o-30^oC (59^o-86^oF)
- avoid excessive heat

Inactive ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

Call **1-888-309-9030**

Principal Display Panel

Compare to the active ingredients in Vicks® Nyquil® Cold & Flu LiquiCaps®†
Night Time

Cold and Flu Relief

Acetaminophen- Pain Reliever/Fever Reducer

Dextromethorphan HBr- Cough suppressant

Doxylamine Succinate- Antihistamine

Relieves:

- Aches, fever, sore throat
- Cough
- Runny nose, sneezing

Alcohol Free

Softgels**

(**Liquid-Filled Capsules)

**This product is not manufactured or distributed by the Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademark of the Procter & Gamble

Company

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT
INFORMATION.**

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

Package Label

<p>Drug Facts (continued)</p> <p>Ask a doctor before use if you have liver disease or glaucoma</p> <p>trouble urinating due to an enlarged prostate gland</p> <p>a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema</p> <p>cough that occurs with too much phlegm (mucus)</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</p> <p>taking sedatives or tranquilizers</p> <p>When using this product do not exceed recommended dosage</p> <p>excitability may occur, especially in children</p> <p>avoid alcoholic drinks</p> <p>marked drowsiness may occur</p> <p>alcohol, sedatives, and tranquilizers may increase drowsiness</p> <p>Be careful when driving a motor vehicle or operating machinery</p> <p>Stop use and ask a doctor if</p> <p>pain or cough gets worse or lasts more than 7 days</p> <p>fever gets worse or lasts more than 3 days</p> <p>redness or swelling is present</p> <p>new symptoms occur</p> <p>cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <p>do not take more than directed (see Overdose warning)</p> <p>do not take more than 4 doses in 24 hours</p>	<p>Drug Facts</p> <p>Active ingredients (in each softgel)</p> <p>Acetaminophen 325 mg.....Pain reliever/fever reducer</p> <p>Dextromethorphan HBr 15 mg.....Cough suppressant</p> <p>Doxylamine succinate 6.25 mg.....Antihistamine</p> <p>Uses</p> <p>temporarily relieves common cold and flu symptoms:</p> <p>sore throat</p> <p>headache</p> <p>minor aches and pains</p> <p>runny nose and sneezing</p> <p>fever</p> <p>cough due to minor throat and bronchial irritation</p> <p>Warnings</p> <p>Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <p>more than 4,000 mg of acetaminophen in 24 hours</p> <p>with other drugs containing acetaminophen</p> <p>3 or more alcoholic drinks every day while using this product</p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening</p> <p>blisters</p> <p>rash</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <p>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <p>If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p>
---	---

DG health

Night Time Cold & Flu Relief



Acetaminophen - Pain Reliever/Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Doxylamine Succinate - Antihistamine

DG health

Compare to the active ingredients of Vicks® NyQuil® Cold & Flu LiquiCaps®†

Night Time Cold & Flu Relief



Acetaminophen - Pain Reliever/Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Doxylamine Succinate - Antihistamine

Relieves:

- Aches, fever, sore throat
- Cough
- Runny nose, sneezing

Alcohol Free



48 Softgels**

(*Liqui-Filled Capsules)

Actual Softgel Size

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

PARENTS:
Learn about how medicine abuse
www.StopMedicineAbuse.org

Lot No.:
Exp. Date:



100% Satisfaction Guaranteed! (888) 309-9030

DISTRIBUTED BY OLD EAST MAIN CO. 100 MISSION RIDGE GOODLETTSVILLE, TN 37072

This product is not manufactured or distributed by The Procter & Gamble Company, Vicks®, MyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

<p>Drug Facts (continued)</p> <p>adults and children 12 years and over: take 2 softgels with water every 6 hours</p> <p>swallow whole; do not crush, chew, or dissolve</p> <p>children under 12 years: do not use</p> <p>when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</p>	<p>Other Information</p> <p>store between 15-30°C (59-86°F)</p> <p>avoid excessive heat</p>
<p>Drug Facts (continued)</p> <p>adults and children 12 years and over: take 2 softgels with water every 6 hours</p> <p>swallow whole; do not crush, chew, or dissolve</p> <p>children under 12 years: do not use</p> <p>when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</p>	<p>Other Information</p> <p>store between 15-30°C (59-86°F)</p> <p>avoid excessive heat</p>
<p>Questions or comments?</p> <p>Call 1-888-309-9030</p>	<p>Drug Facts (continued)</p> <p>adults and children 12 years and over: take 2 softgels with water every 6 hours</p> <p>swallow whole; do not crush, chew, or dissolve</p> <p>children under 12 years: do not use</p> <p>when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</p>

DOLLAR GENERAL Cold & Flu Relief

COLD AND FLU RELIEF NIGHTTIME

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-149
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	20mm

Flavor		Imprint Code	P30	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-149-48	48 in 1 CARTON	12/27/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	12/27/2019		

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 3/2024

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)