

**NAPROXEN SODIUM- naproxen sodium tablet, film coated**  
**CVS Pharmacy**

-----  
**CVS Pharmacy, Inc. Naproxen Sodium Tablets Drug Facts**

**Active ingredient (in each caplet)**

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

### **Adults and children 12 years and older:**

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

### **Children under 12 years:**

- ask a doctor

### **Other information**

- **each caplet contains:** sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

### **Inactive ingredients**

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

### **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

Compare to the active ingredient in Aleve®

Caplets†

Naproxen Sodium

TABLETS, 220 mg

STRENGTH TO LAST 12 HOURS

Pain reliever/Fever reducer (NSAID)

Actual Size

270 CAPLETS†

†Capsule-shaped tablets

**Drug Facts (continued)**

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery

**Ask a doctor before use if** the stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke you are taking a diuretic you have problems or serious side effects from taking pain relievers or fever reducers



Compare to the active ingredient in Aleve®\*\*

**Caplets†**

NDC 69842-010-37

# Naproxen Sodium

TABLETS, 220 mg

**Pain reliever/Fever reducer (NSAID)**



Actual Size



**270 CAPLETS†**

†Capsule-shaped tablets

**Drug Facts**

**Active ingredient (in each caplet)**  
Naproxen sodium 220 mg  
(naproxen 200 mg) (NSAID)\*.....fever reducer  
\*nonsteroidal anti-inflammatory drug

**Purposes**  
Pain reliever/  
fever reducer

**Uses** temporarily relieves minor aches and pains due to:  
 minor pain of arthritis muscular aches backache  
 menstrual cramps headache toothache  
 the common cold temporarily reduces fever

**Warnings**  
**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug

#879502



0 50428 29435 53462 5

**DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING**

368H8 17 F2

PEEL BACK HERE

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are**  
 under a doctor's care for any serious condition taking any other drug

**When using this product** take with food or milk; if stomach upset occurs

**Stop use and ask a doctor if**  
 you experience any of the following signs of stomach bleeding:  
 feel faint vomit blood have bloody or black stools  
 have stomach pain that does not get better  
 you have symptoms of heart problems or stroke:  
 chest pain trouble breathing weakness in one part or side of body  
 slurred speech leg swelling  
 pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days you have difficulty swallowing it feels like the pill is stuck in your throat redness or swelling is present in the painful area any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** do not take more than directed the smallest effective dose should be used drink a full glass of water with each dose  
**Adults and children 12 years and older:** take 1 caplet every 8 to 12 hours while symptoms last for the first dose you may take 2 caplets within the first hour do not exceed 2 caplets in any 8- to 12-hour period do not exceed 3 caplets in a 24-hour period  
**Children under 12 years:** ask a doctor

**Other information** each caplet contains: sodium 21 mg store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients:** FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?** 1-800-719-9280

\*\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve®.

**Distributed by: CVS Pharmacy, Inc.**  
 One CVS Drive, Woonsocket, RI 02895  
 © 2019 CVS/pharmacy  
 CVS.com® 1-800-SHOP CVS  
 V-17160

**CVS Quality**  
 Money Back Guarantee

**STRENGTH TO LAST 12 HOURS**

## NAPROXEN SODIUM

naproxen sodium tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69842-010
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

**Inactive Ingredients**

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

**Product Characteristics**

Color	BLUE (Light Blue)	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	L368
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-010-62	1 in 1 CARTON	03/15/2016	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69842-010-71	1 in 1 CARTON	10/18/2016	09/30/2019
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:69842-010-78	1 in 1 CARTON	10/18/2016	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:69842-010-37	1 in 1 CARTON	10/18/2016	
4		270 in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	03/15/2016	

**Labeler** - CVS Pharmacy (062312574)