

RITE AID FIBER CAPLETS- calcium polycarbophil tablet, film coated
RITE AID CORPORATION

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rite Aid Fiber Caplets Calcium Polycarbophil

Drug Facts

Active Ingredient (in each caplet)

Calcium polycarbophil 625 mg
equivalent to 500 mg polycarbophil

PURPOSE

Bulk-forming laxative

USES

- relieves occasional constipation to help restore and maintain regularity
- this product generally produces bowel movement in 12 to 72 hours

Choking Warning: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that persists over a period of 2 weeks

Ask your doctor or pharmacist before use if you are taking any other drug.

Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product

- do not use for more than 7 days unless directed by a doctor
- do not take more than 8 caplets in a 24 hour period unless directed by a doctor

Stop use and ask a doctor if

rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

DIRECTIONS

- Take each dose of this product with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. (See choking warning).
- This product works naturally so continued use for one to three days is normally required to provide full benefit. Dosage may vary according to diet, exercise, previous laxative use or severity of constipation.

| Age | Recommended | Daily maximum |
|--|----------------------|----------------------|
| adults & children 12 years of age and over | 2 caplets once a day | up to 4 times a day |
| children under 12 years | consult a physician | |

OTHER INFORMATION

- **each caplet contains:** calcium 125 mg
- store at room temperature
- protect contents from moisture

SAFETY SEALED: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

calcium carbonate, caramel color, crospovidone, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium lauryl sulfate.

Questions or Comments?

1-866-467-2748

PRINCIPAL DISPLAY PANEL

NDC# 11822-2025-0

Compare to the active ingredients FiberCon[®] Caplets*

Fiber Caplets

Calcium Polycarbophil 625 mg

Bulk-Forming

Fiber Laxative

- Gentle Relief of Constipation
- Won't Cause Gas or Bloating
- Effective as Fiber Powders

250 Caplets

Distributed By:

*This product is not manufactured or distributed by Pfizer Inc., owner of the registered trademark Fibercon[®] Caplets.

Compare to the active ingredient in
FiberCon® Caplets*

NDC 11822-2025-0

Fiber Caplets

Calcium Polycarbophil

**Bulk-Forming
Fiber Laxative**

- Gentle Relief of Constipation
- Won't Cause Gas or Bloating
- Effective as Fiber Powders

250 Caplets



DO NOT USE IF PRINTED INNER SEAL UNDER CAP IS TORN OR MISSING

| | |
|---|-------------------------|
| Drug Facts | Purpose |
| Active ingredient (in each caplet) Calcium polycarbophil 625 mg equivalent to 500 mg polycarbophil | Bulk-forming laxative |
| Uses ■ relieves occasional constipation to help restore and maintain regularity ■ this product generally produces bowel movement in 12 to 72 hours. | |
| Warnings | |
| Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting or difficulty in swallowing or breathing after taking this product, seek immediate medical attention. | |
| Ask your doctor before use if you have ■ abdominal pain, nausea or vomiting ■ a sudden change in bowel habits that persists over a period of 2 weeks. | |
| Ask your doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work. | |
| When using this product ■ do not use for more than 7 days unless directed by a doctor ■ do not take more than 8 caplets in a 24 hour period unless directed by a doctor | |
| Stop use and ask a doctor if rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition. | |
| Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222. | |
| Directions ■ take each dose of this product with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning. ■ this product works naturally so continued use for one to three days is normally required to provide full benefit. Dosage may vary according to diet, exercise, previous laxative use or severity of constipation | |
| age | recommended dose |
| adults & children | 2 caplets |
| 12 years of age and over | once a day |
| children under 12 years | consult a physician |
| | daily maximum |
| | up to 4 times |
| | a day |
| | |
| Other information ■ each caplet contains: calcium 125 mg ■ store at room temperature ■ protect contents from moisture | |
| Inactive ingredients calcium carbonate, caramel color, croscovidone, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium lauryl sulfate | |
| Questions or comments? 1-866-467-2748 | |

*This product is not manufactured or distributed by the same company as the registered trademark FiberCon® Caplets.

120250XXLR



LOT:

EXP:

RITE AID FIBER CAPLETS

calcium polycarbophil tablet, film coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:11822-2025 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-----------------------|----------|
| CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49) (POLYCARBOPHIL - UNII:W25LM17A4W) | CALCIUM POLYCARBOPHIL | 625 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| CALCIUM CARBONATE (UNII: H0G9379FGK) | |
| CAMEL (UNII: T9D99G2B1R) | |
| CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| SODIUM LAURYL SULFATE (UNII: 368GB514J) | |

Product Characteristics

| | | | |
|---------------|-------|---------------------|----------|
| Color | WHITE | Score | no score |
| Shape | OVAL | Size | 19 mm |
| Flavor | | Imprint Code | RP120 |

Contains**Packaging**

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:11822-2025-0 | 250 in 1 BOTTLE; Type 0: Not a Combination Product | 04/20/2020 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part334 | 04/20/2020 | |

Labeler - RITE AID CORPORATION (014578892)

Revised: 5/2020

RITE AID CORPORATION