# PEPTO BISMOL CHEWABLES- bismuth subsalicylate tablet, chewable Procter & Gamble Manufactura S de RL de CV

-----

Pepto Bismol ® CHEWABLES

#### **Drug Facts**

#### **Active ingredient (in each tablet)**

Bismuth subsalicylate 262 mg

## Active ingredient (in each tablet) Purpose

Upset stomach reliever and antidiarrheal

#### Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

## Warnings

**Reye's syndrome:**Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

## Allergy alert: Contains salicylate. Do not take if you are

• allergic to salicylates (including aspirin) • taking other salicylate products

#### Do not use if

you have

- an ulcer
- a bleeding problem
- bloody or black stool

#### Ask a doctor before use if you have

- fever
- mucus in the stool

#### Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

#### When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

#### Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast feeding,** ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- chew or dissolve in mouth
- adults and children 12 years and over: 2 tablets (1 dose) every 1/2 hour or 4 tablets (2 doses) every hour as needed for diarrhea
- 2 tablets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

#### Other information

- each tablet contains:calcium 140 mg, magnesium 9 mg
- salicylate 99 mg
- low sodium
- avoid excessive heat (over 104°F or 40°C)

## Inactive ingredients

adipic acid, calcium carbonate, D&C Red No. 27 aluminum lake, FD&C Red No. 40 aluminum lake, flavor, magnesium stearate, mannitol, povidone, saccharin sodium, talc

### **Questions?**

1-800-717-3786

TAMPER EVIDENT: Do not use if the printed inner wrap is broken or missing.

MADE IN MEXICO

**DIST. BY PROCTER & GAMBLE** 

**CINCINNATI, OH 45202** 

#### PRINCIPAL DISPLAY PANEL - 30 Tablet Carton

Pepto Bismol ®

Bismuth Subsalicylate Upset Stomach Reliever/ Antidiarrheal

#### **CHEWABLES**

**CHERRY** 

**5 SYMPTOM** 

RELIEF

NAUSEA HEARTBURN INDIGESTION UPSET STOMACH DIARRHEA

30 CHEWABLE TABLETS



### **PEPTO BISMOL CHEWABLES**

bismuth subsalicylate tablet, chewable

<b>Product Information</b>	roduct Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58933-013	
Route of Administration	ORAL			

Active Ingredient/Active Moiety				
Ingredient Name	<b>Basis of Strength</b>	Strength		
	BISMUTH SUBSALICYLATE	525 mg in 2		

Inactive Ingredients		
Ingredient Name	Strength	
ADIPIC ACID (UNII: 76A0JE0FKJ)		
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)		

TALC (UNII: 7SEV7J4R1U)  FD&C RED NO. 40 (UNII: WZB9127XOA)  CALCIUM CARBONATE (UNII: H0G9379FGK)  MAGNESIUM STEARATE (UNII: 70097M6I30)  MANNITOL (UNII: 3OWL53L36A)  POVIDONE (UNII: FZ 989GH94E)  D&C RED NO. 27 (UNII: 2LRS185U6K)  ALUMINUM OXIDE (UNII: LMI2606933)	SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
CALCIUM CARBONATE (UNII: H0G9379FGK)  MAGNESIUM STEARATE (UNII: 70097M6I30)  MANNITOL (UNII: 30WL53L36A)  POVIDONE (UNII: FZ 989GH94E)  D&C RED NO. 27 (UNII: 2LRS185U6K)	TALC (UNII: 7SEV7J4R1U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)  MANNITOL (UNII: 30WL53L36A)  POVIDONE (UNII: FZ 989GH94E)  D&C RED NO. 27 (UNII: 2LRS185U6K)	FD&C RED NO. 40 (UNII: WZB9127XOA)	
MANNITOL (UNII: 30WL53L36A)  POVIDONE (UNII: FZ 989GH94E)  D&C RED NO. 27 (UNII: 2LRS185U6K)	CALCIUM CARBONATE (UNII: H0G9379FGK)	
POVIDONE (UNII: FZ 989GH94E) D&C RED NO. 27 (UNII: 2LRS185U6K)	MAGNESIUM STEARATE (UNII: 70097M6I30)	
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	MANNITOL (UNII: 30WL53L36A)	
	POVIDONE (UNII: FZ989GH94E)	
ALUMINUM OXIDE (UNII: LMI2606933)	<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	
	ALUMINUM OXIDE (UNII: LMI2606933)	

Product Characteristics			
Color	pink	Score	no score
Shape	ROUND	Size	16mm
Flavor	CHERRY	Imprint Code	Pepto;Bismol
Contains			

	ackaging				
,	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
	NDC:58933-013-	2 in 1 CELLO PACK; Type 0: Not a Combination Product	06/30/2004		
:	NDC:58933-013- 04	5 in 1 CARTON	06/30/2004		
:	2	6 in 1 BLISTER PACK; Type 0: Not a Combination Product			
	NDC:58933-013-	5959 in 1 BAG; Type 0: Not a Combination Product	06/30/2004		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	06/30/2004	

## Labeler - Procter & Gamble Manufactura S de RL de CV (812807550)

Revised: 10/2023 Procter & Gamble Manufactura S de RL de CV