PEPTO BISMOL CHEWABLES- bismuth subsalicylate tablet, chewable
Procter & Gamble Manufactura S de RL de CV

Pepto Bismol ®
CHEWABLES

Drug Facts

Active ingredient (in each tablet)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves
- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Contains salicylate. Do not take if you are
- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if

you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have
- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for
- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if
- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• chew or dissolve in mouth

• adults and children 12 years and over:
- 2 tablets (1 dose) every 1/2 hour or 4 tablets (2 doses) every hour as needed for diarrhea
- 2 tablets (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)

• do not exceed 8 doses (16 tablets) in 24 hours

• use until diarrhea stops but not more than 2 days

• children under 12 years: ask a doctor

• drink plenty of clear fluids to help prevent

dehydration caused by diarrhea

Other information
- **each tablet contains**: calcium 140 mg, magnesium 9 mg
- salicylate 101 mg
- low sodium
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

calcium carbonate, D&C Red No. 27 aluminum lake, flavor, magnesium stearate, mannitol, povidone, saccharin sodium, talc

Questions?

1-800-717-3786

TAMPER EVIDENT: Do not use if the printed inner wrap is broken or missing.

MADE IN MEXICO

DIST. BY PROCTER & GAMBLE

CINCINNATI OH 45202

PRINCIPAL DISPLAY PANEL - 12 Chewable Tablet Carton

**Pepto
Bismol®**

Bismuth Subsalicylate

Upset Stomach Reliever/ Antidiarrheal

CHEWABLES

5 SYMPTOM

RELIEF

**NAUSEA
HEARTBURN
INDIGESTION
UPSET STOMACH
DIARRHEA**

12 CHEWABLE TABLETS

Drug Facts

Active ingredient (in each tablet) — **Purpose**
Bismuth subsalicylate 262 mg .... Upset stomach reliever and antidiarrheal

Uses relieves • travelers' diarrhea • diarrhea
• upset stomach due to overindulgence in food and drink, including:
• heartburn • indigestion • nausea • gas • belching • fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are
• allergic to salicylates (including aspirin) • taking other salicylate products

Do not use if you have • an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have • fever • mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
• anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if • symptoms get worse or last more than 2 days
• ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• chew or dissolve in mouth
• adults and children 12 years and over:
• 2 tablets (1 dose) every ½ hour or 4 tablets (2 doses) every hour as needed for diarrhea
• 2 tablets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
• do not exceed 8 doses (16 tablets) in 24 hours
• use until diarrhea stops but not more than 2 days
• children under 12 years: ask a doctor
• drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Drug Facts (continued)
Inactive ingredients calcium carbonate, D&C Red No. 27 aluminum lake, flavor, magnesium stearate, mannitol, povidone, saccharin sodium, talc

Questions? 1-800-717-3786

Drug Facts (continued)
Other information
• each tablet contains: calcium 140 mg, magnesium 9 mg • salicylate 101 mg • low sodium • avoid excessive heat (over 104°F or 40°C)

Pepto Bismol®
Bismuth Subsalicylate
Upset Stomach Reliever/ Antidiarrheal
CHEWABLES

5 SYMPTOM RELIEF
NAUSEA
HEARTBURN
INDIGESTION
UPSET STOMACH
DIARRHEA

12 CHEWABLE TABLETS

TAMPER EVIDENT: Do not use if the printed inner wrap is broken or missing.

MADE IN MEXICO
DIST. BY PROCTER & GAMBLE, CINCINNATI OH 45202
www.pg.com
Patent: www.pg.com/patents
www.pepto-bismol.com
P&G
91510372

LOT   EXP

PEPTO BISMOL CHEWABLES
bismuth subsalicylate tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58933-011
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

MANNITOL (UNII: 3OWL53L36A)	
POVIDONE (UNII: FZ989GH94E)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	pink	**Score**	no score
Shape	ROUND	**Size**	16mm
Flavor	WINTERGREEN	**Imprint Code**	Pepto;Bismol
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-011-02	2 in 1 PACKET; Type 0: Not a Combination Product	06/30/2004	
2	NDC:58933-011-12	2 in 1 CARTON	06/30/2004	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:58933-011-30	5 in 1 CARTON	06/30/2004	
3		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:58933-011-48	8 in 1 CARTON	06/30/2004	
4		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:58933-011-04	4 in 1 PACKET; Type 0: Not a Combination Product	06/30/2004	
6	NDC:58933-011-90	3 in 1 BLISTER PACK	10/04/2022	
6		5 in 1 CARTON		
6		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	06/30/2004	

Labeler - Procter & Gamble Manufactura S de RL de CV (812807550)

Revised: 10/2025

Procter & Gamble Manufactura S de RL de CV