

**GOOD NEIGHBOR PHARMACY CHILDRENS PAIN AND FEVER-
acetaminophen suspension
Amerisource Bergen**

Amerisource Bergen Children's Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child’s dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL

60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- **each 5 mL contains:** sodium 3 mg
- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Children's Tylenol® Oral Suspension active ingredient

GOOD NEIGHBOR PHARMACY®

FOR AGES 2 TO 11 YEARS

children's Pain & Fever

acetaminophen 160 mg per 5 mL oral suspension

Fever Reducer – Pain Reliever

Aspirin Free


Ibuprofen Free

Alcohol Free

Paraben Free

Bubble Gum Flavor

4 fl oz (118 mL)



Drug Facts
Active ingredient (in each 5 mL) Acetaminophen 160 mg.....Pain reliever/fever reducer
Purpose
Uses temporarily:
 ■ reduces fever
 ■ relieves minor aches and pains due to:
 ■ the common cold ■ flu ■ headache
 ■ sore throat ■ toothache
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes
 ■ more than 5 doses in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if your child has ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if your child has liver disease
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin
When using this product do not exceed recommended dose (see overdose warning)
Stop use and ask a doctor if
 ■ pain gets worse or lasts more than 5 days
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur
 ■ redness or swelling is present
 These could be signs of a serious condition.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)
Directions
 ■ this product does not contain directions or complete warnings for adult use
 ■ do not give more than directed (see overdose warning)
 ■ shake well before using
 ■ mL = milliliter
 ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.
 ■ remove the child protective cap and squeeze your child's dose into the dosing cup
 ■ repeat dose every 4 hours while symptoms last
 ■ do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor
Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
Other information
 ■ each 5 mL contains: sodium 3 mg
 ■ store at 20-25°C (68-77°F)
 ■ do not use if printed neckband is broken or missing
Inactive ingredients
 anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate
Questions or comments? 1-800-719-9260
 *Good Neighbor Pharmacy® Children's Pain & Fever is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Tylenol® Oral Suspension.
 Distributed By
 AmerisourceBergen **GLUTEN FREE**
 1 West First Avenue
 Conshohocken, PA 19428
 Questions or Concerns?
 www.mygnp.com

NDC 46122-209-26
 Compare to Children's Tylenol® Oral Suspension active ingredient**
GOOD NEIGHBOR PHARMACY®

FOR AGES 2 TO 11 YEARS
 children's
Pain & Fever
 acetaminophen
 160 mg per 5 mL
 oral suspension
Fever Reducer - Pain Reliever
 Aspirin Free
 Ibuprofen Free
 Alcohol Free
 Paraben Free
Bubble Gum Flavor
 4 fl oz (118 mL)

GOOD NEIGHBOR PHARMACY®
 children's
Pain & Fever
DIRECTIONS:
STEP 1. Shake well before using

STEP 2. Remove child protective cap

STEP 3. Squeeze your child's dose into the dosing cup

IMPORTANT: Please see dosing chart or ask a doctor for your child's correct dose.



ABC#: 10052416

 10526 29 C12

GOOD NEIGHBOR PHARMACY CHILDRENS PAIN AND FEVER			
acetaminophen suspension			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-209
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	

Product Characteristics

Color	RED (opaque)	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-209-25	1 in 1 CARTON	08/15/2013	06/15/2015
1		148 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:46122-209-26	1 in 1 CARTON	08/15/2013	
2		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/15/2013	

