

MEIJER SPF 50 SUNSCREEN- avobenzone 3%, homosalate 10%, octisalate 5%, octocrylene 10% lotion

Meijer

Avobenzone 3%, Homosalate 15%, Ocitsalate 5%, Octocrylene 10%

Sunscreen

Helps prevent sunburn.

If used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun.

For external use only.

Do not use on damaged or broken skin.

When using this product avoid contact with eyes. Rinse with water to remove.

Stop use and ask a doctor if rash or irritation develops and lasts.

If product is swallowed, get medical help or contact a Poison Control Center right away.

Apply liberally 15 minutes before sun exposure and as needed. Reapply: after 80 minutes of swimming or sweating, immediately after towel drying, at least every 2 hours.

Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: limit time in the sun, especially from 10 a.m. - 2 p.m., wear long-sleeved shirts, pants, hats, and sunglasses, children under 6 months of age: ask a doctor.

Water, sorbitol, aluminum starch octenylsuccinate, VP/Eicosense copolymer, stearic acid, dimethicone, tocopherol, ethylhexylglycerin, phenoxyethanol, polyglyceryl-3 distearate, sorbitan isostearate, carbomer, disodium EDTA, triethanolamine, benzyl alcohol, fragrance.

NDC 79481-1830-8

meijer.

sunscreen lotion

BROAD SPECTRUM
SPF 50 SUNSCREEN



UVA/UVB PROTECTION

MOISTURIZING

PARABEN FREE

WATER RESISTANT (80 MINUTES)

NET WT 8 OZ (227 g)



Drug Facts

Active ingredients	Purpose
Avobenzone 3%	Sunscreen
Homosalate 13%	Sunscreen
Octisalate 5%	Sunscreen
Octocrylene 10%	Sunscreen

Uses • helps prevent sunburn • if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only.

Do not use on damaged or broken skin.

When using this product avoid contact with eyes. Rinse with water to remove.

Stop use and ask a doctor if rash or irritation develops and lasts.

Keep out of the reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions • apply liberally 15 minutes before sun exposure and as needed. Reapply: • after 80 minutes of swimming or sweating • immediately after towel drying • at least every 2 hours

Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m.-2 p.m.
- wear long-sleeved shirts, pants, hats, and sunglasses
- children under 6 months of age: ask a doctor

Other information • protect the product in this container from excessive heat and direct sun • may stain fabrics

Inactive ingredients water, sorbitol, aluminum starch octenylsuccinate, VP/eicosene copolymer, stearic acid, dimethicone, tocopherol, ethylhexylglycerin, phenoxylethanol, polyglyceryl-3 distearate, sorbitan isotearate, carbomer, disodium EDTA, triethanolamine, benzyl alcohol, fragrance

Questions? 1-844-987-2727

DIST BY: MEIJER DISTRIBUTION INC.
GRAND RAPIDS, MI 49544
www.meijer.com

Made in the USA with Domestic and Imported Ingredients.



www.meijer.com/
satisfaction

PID 2869972



NDC 79481-1830-9

meijer.

kids
suncscreen lotion

BROAD SPECTRUM
SPF 50 SUNSCREEN



UVA/UVB PROTECTION

MOISTURIZING

PARABEN FREE

WATER RESISTANT (80 MINUTES)

NET WT 8 OZ (227 g)



Drug Facts

Active ingredients	Purpose
Avobenzone 3%	Sunscreen
Homosalate 15%	Sunscreen
Octisalate 5%	Sunscreen
Octocrylene 10%	Sunscreen

Uses • helps prevent sunburn • if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only.
Do not use on damaged or broken skin.
When using this product avoid contact with eyes. Rinse with water to remove.
Stop use and ask a doctor if rash or irritation develops and lasts.

Keep out of the reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions • apply liberally 15 minutes before sun exposure and as needed. Reapply: • after 80 minutes of swimming or sweating • immediately after towel drying • at least every 2 hours

Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
• limit time in the sun, especially from 10 a.m. - 2 p.m.
• wear long-sleeved shirts, pants, hats, and sunglasses
• children under 6 months of age: ask a doctor

Other information • protect the product in this container from excessive heat and direct sun • may stain fabrics

Inactive ingredients water, sorbitol, aluminum starch octenylsuccinate, VP/eicosene copolymer, stearic acid, dimethicone, tocopherol, ethylhexylglycerin, phenoxyethanol, polyglyceryl-3 distearate, sorbitan isostearate, carbomer, disodium EDTA, triethanolamine, benzyl alcohol, fragrance

Questions? 1-844-987-2727

DIST BY: MEIJER DISTRIBUTION INC.
GRAND RAPIDS, MI 49544
www.meijer.com

Made in the USA with Domestic and Imported Ingredients.



PID 3828348



MEIJER SPF 50 SUNSCREEN

avobenzone 3%, homosalate 10%, octisalate 5%, octocrylene 10% lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-1830
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	15 g in 100 g
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	3 g in 100 g
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	10 g in 100 g
OCTISALATE (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE 200 (UNII: RGS4T2AS00)	
CARBOMER 940 (UNII: 4Q93RCW27E)	
POLYGLYCERYL-3 DISTEARATE (UNII: ZI1LK470XV)	
TROLAMINE (UNII: 9O3K93S3TK)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	
SORBITAN ISOSTEARATE (UNII: 01S2G2C1E4)	
EICOSYL POVIDONE (UNII: XQQ9MKE2BJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
TOCOPHEROL (UNII: ROZB2556P8)	
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
ALUMINUM STARCH OCTENYLSUCCINATE (UNII: I9PJ006294)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-1830-8	227 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/01/2024	
2	NDC:79481-1830-9	227 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	02/28/2024	

Labeler - Meijer (006959555)

Registrant - Derma Care Research Labs, LLC (116817470)

Establishment

Name	Address	ID/FEI	Business Operations
Derma Care Research Labs, LLC		116817470	manufacture(79481-1830)

Revised: 2/2024

Meijer