

KANKA TRIPLE ACTION- benzocaine and menthol, unspecified form gel

Blistex Inc.

Kanka® Triple Action Gel

Drug Facts

| Active ingredients | Purpose |
|---------------------------|--|
| Benzocaine 20.0% (w/w) | Oral anesthetic/analgesic (oral pain reliever) |
| Menthol 2.0% (w/w) | Oral anesthetic/analgesic (oral pain reliever) |

Uses

- for the temporary relief of pain associated with canker sores, minor irritation of the mouth and gums caused by dentures or orthodontic appliances, or minor injury of the mouth or gums.

Warnings

METHEMOGLOBINEMIA WARNING

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert

do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use

- for teething
- in children under 2 years of age

When using this product

- do not use this product for more than 7 days unless directed by a dentist or doctor. If sore mouth symptoms do not improve in 7 days; if irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your dentist or doctor promptly.

- do not exceed recommended dosage.

Keep out of reach of children.

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions

| | |
|--|--|
| Adults and children 2 years of age and older | Apply to affected area up to 4 times daily or as directed by a dentist or doctor |
| Children between 2 and 12 years of age | Ask a doctor before use. Should be supervised in the use of this product |
| Children under 2 years of age | Do not use |

Other information

- www.kank-a.com

Inactive ingredients

ascorbyl palmitate, cetylpyridinium chloride, PEG-75, phenoxyethanol, polyethylene glycol, sucralose, thymol

PRINCIPAL DISPLAY PANEL - 11.9 g Tube Carton

MAXIMUM STRENGTH

KANKA®

TRIPLE ACTION GEL
ORAL PAIN RELIEVER

Mouth Pain Relief for Canker Sores

Net Wt. 0.42 oz. (11.9 g)

NEW!

MAXIMUM STRENGTH

KANK-A[®]

TRIPLE ACTION GEL

ORAL PAIN RELIEVER

Mouth Pain Relief for Canker Sores

Fast-Acting + Long-Lasting*

Kills 99% of Germs for a Fresh Feel

Cools + Soothes Irritation



Helps Relieve Mouth Pain From:

- Canker Sores
- Mouth Irritation
- Gum Discomfort
- Mouth Sores



Net Wt. 0.42 oz. (11.9 g)



#C030342
 ♻️ Carton is 100% Recyclable.
 ©2024 Blistex Inc., P.O. Box 5392
 Oak Brook, IL 60522-5392

* Long-lasting pain relief

Warnings
 METHEMOGLOBINEMIA WARNING: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops pale, gray, or blue colored skin (cyanosis), headache, rapid heart rate, shortness of breath, dizziness or lightheadedness, fatigue or lack of energy.

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Uses
 For the temporary relief of pain associated with canker sores, minor irritation of the mouth and gums caused by dentures or orthodontic appliances, or minor injury of the mouth or gums.

Directions
 Adults and children 2 years of age and older: Apply to affected area up to 4 times daily or as directed by a dentist or doctor.
 Children between 2 and 12 years of age: Ask a doctor before use. Should be supervised in the use of this product.
 Children under 2 years of age: Do not use.

Other information
 • www.kank-a.com

Inactive ingredients
 phenylethanolamine hydrochloride, polyethylene glycol, sucralose, thymol, ascorbyl palmitate, cetylpyridinium chloride, PEG-75.

When using this product
 • do not use this product for more than 7 days unless directed by a dentist or doctor; if sore mouth symptoms do not improve in 7 days; if irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your dentist or doctor promptly.
 • do not exceed recommended dosage.
 • keep out of reach of children.
 • if more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Methyl 2.0% (w/w)
 Oral anesthetic (analgesic)
 (oral pain reliever)



KANKA TRIPLE ACTION

benzocaine and menthol, unspecified form gel

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:10157-2127 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|------------------------------|------------------|
| BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) | BENZOCAINE | 20 g in 100 g |
| MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A) (MENTHOL, UNSPECIFIED FORM - UNII:L7T10EIP3A) | MENTHOL, UNSPECIFIED FORM | 2 g in 100 g |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A) | |
| POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| PHENOXYETHANOL (UNII: H1E492ZZ3T) | |
| ASCORBYL PALMITATE (UNII: QN83US2B0N) | |
| THYMOL (UNII: 3J50XA376E) | |
| CETYLPIRIDINIUM CHLORIDE (UNII: D9OM4SK49P) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:10157-2127-1 | 1 in 1 CARTON | 01/06/2025 | |
| 1 | | 11.9 g in 1 TUBE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC monograph drug | M022 | 01/06/2025 | |

Labeler - Blistex Inc. (005126354)

Establishment

| Name | Address | ID/FEI | Business Operations |
|--------------|---------|-----------|-------------------------|
| Blistex Inc. | | 005126354 | MANUFACTURE(10157-2127) |

Revised: 1/2025

Blistex Inc.