

**ANTI-DIARRHEAL- bismuth subsalicylate tablet**  
**Lil' Drug Store Products, Inc.**

-----  
**Anti-Diarrheal**

***Drug Facts***

**Active ingredient (in each caplet)**

Bismuth subsalicylate 262 mg

**Purpose**

Upset stomach reliever and anti-diarrheal

**Uses**

relieves:

- indigestion
- nausea
- heartburn
- diarrhea
- upset stomach associated with these symptoms

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

**Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use** if you have

- bloody or black stool
- an ulcer
- a bleeding problem

**Ask a doctor before use if** you have

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are** taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

**When using this product** a temporary, but harmless darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.

### **Directions**

- swallow caplet(s) with water, do not chew
- adults and children 12 years and over: 2 caplets every 1/2 to 1 hour as needed
- do not take more than 8 doses (16 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### **Other information**

- **each caplet contains:** calcium 27 mg, salicylate 102 mg and sodium 2 mg
- very low sodium
- store at 59°-77°F (15°-25°C), in a dry place

### **Inactive ingredients**

calcium carbonate, colloidal silicon dioxide, D&C red 27 lake, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, sodium starch glycolate

### **Questions or comments?**

call toll-free **1-877-507-6516** (M-F 8AM-4:30PM CST)

**Product distributed by:** Lil' Drug Store Products, Inc.  
9300 Earhart Lane SW, Cedar Rapids, IA 52404

**PRINCIPAL DISPLAY PANEL - 262 mg Caplet Bottle Carton**

QUALITY

★ GUARANTEED ★

Anti-  
Diarrheal

Bismuth subsalicylate 262 mg

Upset Stomach Reliever/  
Anti-diarrheal

- Relieves the Symptoms  
of Diarrhea

ACTUAL SIZE

24  
Caplets

Lil'  
Drug Store®

# Anti-Diarrheal

Bismuth subsalicylate 262 mg  
Upset Stomach Reliever/  
Anti-diarrheal

24  
Caplets



# Anti-Diarrheal

Bismuth subsalicylate 262 mg  
Upset Stomach Reliever/  
Anti-diarrheal

- Relieves the Symptoms of Diarrhea



ACTUAL SIZE

24  
Caplets



Product distributed by: Lil' Drug Store Products, Inc.  
9300 Earhart Lane SW, Cedar Rapids, IA 52404



Blank Area  
for  
LOT & EXP.  
Information

**TAMPER EVIDENT: DO NOT USE IF  
PRINTED SEAL UNDER CAP  
IS TORN OR MISSING**

**KEEP OUTER PACKAGE FOR  
COMPLETE PRODUCT INFORMATION**

## Drug Facts

Active ingredient (in each caplet)	Purpose
---------------------------------------	---------

Bismuth subsalicylate 262 mg.....	Upset stomach reliever and anti-diarrheal
-----------------------------------	---

## Uses

- relieves:
- indigestion
  - heartburn
  - nausea
  - diarrhea
  - upset stomach associated with these symptoms

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

68414C-US-11-18

**Drug Facts** (continued)

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use** if you have

- bloody or black stool
- an ulcer
- a bleeding problem

**Ask a doctor before use** if you have

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use** if you are taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

**When using this product** a temporary, but harmless darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor** if

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately. ▶

**Drug Facts** (continued)

**Directions**

- swallow caplet(s) with water, do not chew
- adults and children 12 years and over:  
2 caplets every 1/2 to 1 hour as needed  
(16 caplets) in 24 hours
- do not take more than 8 doses  
(16 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**

- **each caplet contains:** calcium 27 mg, salicylate 102 mg and sodium 2 mg
- very low sodium
- store at 59°-77°F (15°-25°C), in a dry place

**Inactive ingredients**

calcium carbonate, colloidal silicon dioxide, D&C red 27 lake, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, sodium starch glycolate

**Questions or comments?** call toll-free 1-877-507-6516 (M-F 8AM-4:30PM CST)

# ANTI-DIARRHEAL

bismuth subsalicylate tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-6841
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	

## Product Characteristics

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	G172
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6841-4	1 in 1 CARTON	02/21/2019	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/21/2019	

---

**Labeler** - Lil' Drug Store Products, Inc. (093103646)

Revised: 11/2024

Lil' Drug Store Products, Inc.