

**TOPCARE ALL DAY PAIN RELIEF- naproxen sodium tablet, film coated**  
**Topco Associates LLC**

-----  
**Topco Associates LLC. All Day Pain Relief Drug Facts**

**Active ingredient (in each caplet)**

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

**Adults and children 12 years and older:**

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

**Children under 12 years:**

- ask a doctor

**Other information**

- **each caplet contains:** sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients**

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?**

**1-888-423-0139**

**Principal Display Panel**

COMPARE TO ALEVE® ACTIVE INGREDIENT

SEE NEW WARNINGS

All Day Pain Relief

NAPROXEN SODIUM TABLETS, 220 mg EACH

PAIN RELIEVER  
 FEVER REDUCER (NSAID)  
 12 HOUR  
 Strength to last 12 hours  
 400 CAPLETS†  
 †CAPSULE-SHAPED TABLETS  
 actual size



NDC 36800-368-79

COMPARE TO ALEVE® ACTIVE INGREDIENT\*\*  
 SEE NEW WARNINGS

# All Day Pain Relief

NAPROXEN SODIUM TABLETS,  
 220 mg EACH  
 PAIN RELIEVER • FEVER REDUCER (NSAID)

**12 HOUR** Strength to last 12 hours

**400 CAPLETS†**  
 †CAPSULE-SHAPED TABLETS

actual size

**Drug Facts**

**Active ingredient (in each caplet)**  
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\* Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Purposes**  
 Pain reliever/fever reducer

**Uses** ■ temporarily relieves minor aches and pains due to: ■ minor pain of arthritis  
 ■ muscular aches ■ backache ■ menstrual cramps ■ headache ■ toothache  
 ■ the common cold ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing)  
 ■ shock ■ skin reddening ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 ■ right before or after heart surgery

**Ask a doctor before use if** ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are** ■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

**Drug Facts (continued)**

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding:  
 ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling  
 ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days  
 ■ you have difficulty swallowing ■ it feels like the pill is stuck in your throat ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used ■ drink a full glass of water with each dose  
**Adults and children 12 years and older:** ■ take 1 caplet every 8 to 12 hours while symptoms last ■ for the first dose you may take 2 caplets within the first hour ■ do not exceed 2 caplets in any 8- to 12-hour period ■ do not exceed 3 caplets in a 24-hour period  
**Children under 12 years:** ■ ask a doctor

**Other information** ■ each caplet contains: sodium 21 mg ■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients** FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?** 1-888-423-0139

DISTRIBUTED BY TOPCARE ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007  
 ©TOPCARE PERM1119 QUESTIONS? 1-888-423-0139  
 topcare@topco.com www.topcarebrand.com

Visit here or call 1-888-423-0139 \*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve®.

GLUTEN FREE  
 QUALITY GUARANTEED

0 36800 45953 3

36879 88 F1

## TOPCARE ALL DAY PAIN RELIEF

naproxen sodium tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-368
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

### Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	

<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	BLUE (Light Blue)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	L368
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-368-62	1 in 1 CARTON	07/22/1997	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:36800-368-71	1 in 1 CARTON	01/14/1997	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:36800-368-78	1 in 1 CARTON	06/11/2002	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:36800-368-82	1 in 1 CARTON	01/27/2004	
4		200 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:36800-368-76	1 in 1 CARTON	06/29/2007	
5		120 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:36800-368-79	400 in 1 BOTTLE; Type 0: Not a Combination Product	12/30/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	01/14/1997	

**Labeler** - Topco Associates LLC (006935977)