

MAXIMUM STRENGTH ACID REDUCER- famotidine tablet
TARGET CORPORATION

Target Famotidine Tablets USP, 20 mg

Drug Facts

Active ingredient (in each tablet)

Famotidine USP 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

adults and children 12 years and over:

- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours

children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture
- FDA approved dissolution test specifications differ from USP

Inactive ingredients carnauba wax, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Questions or comments? Call **1-800-910-6874**

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

NDC 11673-697-20

Maximum Strength Acid Reducer

Famotidine Tablets USP, 20 mg

Just one tablet prevents and relieves heartburn due to acid indigestion



200 TABLETS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Purpose
Famotidine USP 20 mg.....Acid reducer
Uses ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages.

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. **Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ adults and children 12 years and over: ■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. ■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn ■ do not use more than 2 tablets in 24 hours ■ children under 12 years: ask a doctor

Other information

■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20°-25°C (68°-77°F) ■ protect from moisture ■ FDA approved dissolution test specifications differ from USP

Questions or comments? Call 1-800-910-6874

245 05 9360 R00 C-002262-01-073-0000

Distributed by Target Corporation

Minneapolis, MN 55403

Made in India TM & ©2024 Target Brands, Inc.

Lot No.:

Exp. Date:

720R 0324

Varnish Omit Area

Print/Varnish Omit Area

Compare to active ingredient in Maximum Strength Pepcid® AC*

Maximum Strength Acid Reducer

Famotidine Tablets USP, 20 mg

Just one tablet prevents and relieves heartburn due to acid indigestion



Actual Size
200 Tablets

200 TABLETS

Drug Facts

Active ingredient (in each tablet) Purpose
Famotidine USP 20 mg.....Acid reducer

Uses

■ relieves heartburn associated with acid indigestion and sour stomach
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
■ with other acid reducers

Ask a doctor before use if you have

■ had heartburn over 3 months. This may be a sign of a more serious condition.

■ heartburn with lightheadedness, sweating, or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

■ frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss ■ nausea or vomiting
■ stomach pain ■ kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Drug Facts (continued)

Stop use and ask a doctor if

■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ adults and children 12 years and over:

■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.

■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn

■ do not use more than 2 tablets in 24 hours
■ children under 12 years: ask a doctor

Other information

■ read the directions and warnings before use

■ keep the carton. It contains important information.

■ store at 20°-25°C (68°-77°F)

■ protect from moisture

■ FDA approved dissolution test specifications differ from USP

Inactive ingredients carnauba wax, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Questions or comments? Call 1-800-910-6874

ACTUAL SIZE



*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc. McNeil Consumer Healthcare Division, owner of the registered trademark Maximum Strength Pepcid® AC

Print/Varnish Omit Area

Print/Varnish Omit Area

Distributed by Target Corporation
Minneapolis, MN 55403
Made in India TM & ©2024 Target Brands, Inc.

245 05 9360 R00 C-002262-01-073-0000



720R 0324

LOT NO:

EXP DATE:

Print/Varnish Omit Area

ACTUAL SIZE

Lot No.:
Exp. Date:

7208 0324



Coating/Print Omit Area

Drug Facts

Active ingredient (in each tablet)
Famotidine USP 20 mg..... Acid reducer

Purpose

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may

Drug Facts (continued)

interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture
- FDA approved dissolution test specifications differ from USP

Inactive ingredients

carnauba wax, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Questions or comments?

Call 1-800-910-6874

THIS PRODUCT IS NOT TO BE USED IN CHILDREN UNDER 12 YEARS OF AGE. SEE THE BACK OF THE BOTTLE FOR MORE INFORMATION.

Satisfaction guaranteed - Love it or your money back.

This product is not a manufacturer or distributor of Johnson & Johnson Consumer Inc. McNeil Consumer Healthcare Division, owner of the registered trademark Maximum Strength Pepcid® AC.

Distributed by: Targem Corporation
Minneapolis, MN 55403

Made in India
TM & © 2024 Targem Brands, Inc.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

- JUST ONE TABLET** prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

NDC 11673-697-01

Compare to active ingredient in Maximum Strength Pepcid® AC*

Maximum Strength Acid Reducer

Famotidine Tablets USP, 20 mg

Just one tablet prevents and relieves heartburn due to acid indigestion



100 TABLETS (50 + 50)



Twin Pack



Actual Size
50 + 50 Tablets

Coating/
Print Omit Area

Coating/
Print Omit Area

Maximum Strength Acid Reducer

Famotidine Tablets USP, 20 mg

Just one tablet prevents and relieves heartburn due to acid indigestion



50 TABLETS

NDC 11673-897-05

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL UNDER BOTTLE CAP OR BAND AROUND THE CENTER OF EACH CAPSULE IS BROKEN OR MISSING

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet)
Famotidine USP 20 mg, Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert Do not use if you are allergic to famotidine or other acid reducers
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers
Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition.

■ heartburn with lightheadedness, sweating, or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting
■ stomach pain ■ kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children

Directions ■ adults and children 12 years and over: ■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. ■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn ■ do not use more than 2 tablets in 24 hours ■ children under 12 years: ask a doctor

Questions or comments? Call 1-800-910-6874
Dist. by Target Corp., Mpls., MN 55403
245161064 R0D
Made in India. TM & ©2024 Target Brands, Inc. C-02282-01-073-000

720R 0324

Lot No.:

Exp. Date:

Varnish Omit Area



ACTUAL SIZE

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

- JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

LOT NO.:
EXP. DATE:

720R 0324



Coating/Print Omit Area

Drug Facts

Active ingredient (in each tablet)
Famotidine USP 20 mg, Acid reducer

Uses
■ relieves heartburn associated with acid indigestion and sour stomach
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
■ with other acid reducers

Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition.

Drug Facts (continued)

■ heartburn with lightheadedness, sweating, or dizziness
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain
■ kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Drug Facts (continued)

Directions
■ adults and children 12 years and over:
■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
■ do not use more than 2 tablets in 24 hours
■ children under 12 years: ask a doctor

Other information

Drug Facts (continued)

■ store at 20°-25°C (68°-77°F)
■ protect from moisture
■ FDA approved dissolution test specifications differ from USP

Inactive ingredients carnauba wax, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Questions or comments? Call 1-800-910-6874

Satisfaction guaranteed – Love it or your money back.

Distributed by Target Corporation
Minneapolis, MN 55403
Made in India
TM & ©2024 Target Brands, Inc.

■ read the directions and warnings before use
 ■ keep the carton. It contains important information. ▶

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Maximum Strength Pepcid® AC.

Compare to active ingredient in
 Maximum Strength Pepcid® AC*

NDC 11673-697-05

Maximum Strength Acid Reducer

Famotidine Tablets USP, 20 mg

Just one tablet prevents and
 relieves heartburn due to
 acid indigestion



50 TABLETS



Actual Size
 50 Tablets

Coating/
 Print Omit Area

Coating/
 Print Omit Area

Love it or your money back.
 Satisfaction guaranteed –

245 050 165 R00 C-002262-01-073-0000
 Distributed by Target Corporation
 Minneapolis, MN 55403
 Made in India TM & ©2024 Target Brands, Inc.
 720R0324

DO NOT USE IF BLISTERPACK IS TORN OR SEALS ARE BROKEN

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Pepcid® AC.

Call 1-800-910-6874
 (Questions or comments?)
 Drug Facts (continued)

Drug Facts
 (in each tablet)
 Active ingredient
 Famotidine USP 20mg
 Acid reducer

Purpose
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days

Uses
 ■ relieves heartburn associated with acid indigestion and heartburn
 ■ soothes stomach
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
 ■ Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ Ask a doctor before use if you have had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating, or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain ■ kidney disease

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Other information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20°-25°C (68°-77°F) ■ protect from moisture
 ■ FDA approved dissolution test specifications differ from USP
 ■ inactive ingredients: camphor wax, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Directions
 ■ adults and children 12 years and over:
 ■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
 ■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
 ■ do not use more than 2 tablets in 24 hours
 ■ children under 12 years: ask a doctor

Other information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20°-25°C (68°-77°F) ■ protect from moisture
 ■ FDA approved dissolution test specifications differ from USP
 ■ inactive ingredients: camphor wax, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

- JUST ONE TABLET prevents and relieves each and every acid indigestion brought on by eating and drinking certain foods and beverages.
- Tip: It will help you by eating the tablet before you eat.
- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Lot No.:
 Exp Date:

Varnish Omit

3 11673 697521 8

Maximum Strength Acid Reducer



MAXIMUM STRENGTH ACID REDUCER

famotidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-697
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
MICROCRYSTALLINE CELLULOSE 102 (UNII: PNR0YF693Y)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics			
Color	white	Score	no score
Shape	SEMI-CIRCLE (D-shaped)	Size	9mm
Flavor		Imprint Code	87
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-697-52	5 in 1 CARTON	05/15/2024	
1		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11673-697-05	1 in 1 CARTON	05/15/2024	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:11673-697-01	2 in 1 CARTON	05/15/2024	
3		50 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:11673-697-20	1 in 1 CARTON	05/15/2024	
4		200 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA217543	05/15/2024	

Labeler - TARGET CORPORATION (006961700)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment			
Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(11673-697)