

5-SYMPTOM DIGESTIVE RELIEF- bismuth subsalicylate tablet
TARGET CORPORATION

Target Bismuth subsalicylate caplets 262mg

Active ingredient (in each caplet)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:

-heartburn

-indigestion

-nausea

-gas

-belching

-fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- swallow with water, do not chew
- adults and children 12 years and over:

-2 caplets (1 dose) every ½ hour or 4 caplets (2 doses) every hour as needed for diarrhea

-2 caplets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)

- do not exceed 8 doses (16 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each caplet contains:** calcium 27 mg
- salicylate 99 mg
- low sodium
- store at room temperature

Inactive ingredients

calcium carbonate, D&C red #27 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, povidone K-30, silicon dioxide, sodium starch glycolate

Questions or comments?


Call 1-877-290-4008

NDC 11673-880-44

5-Symptom Digestive Relief

Bismuth Subsalicylate, 262 mg
Upset Stomach Reliever / Antidiarrheal

Relieves upset stomach, nausea, heartburn, indigestion and diarrhea



40 CAPLETS**
(**CAPSULE-SHAPED TABLETS)

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet)
Bismuth subsalicylate 262 mg.....Upset stomach reliever and antidiarrheal

Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: heartburn, indigestion, nausea, gas, belching, fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms, should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are allergic to salicylates (including aspirin) taking other salicylate products

If pregnant or breast feeding, ask a health professional before use. **Keep out of reach of children.**

Directions ■ swallow with water, do not chew ■ adults and children 12 years and over: ■ 2 caplets (1 dose) every ½ hour or 4 caplets (2 doses) every hour as needed for diarrhea ■ 2 caplets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (16 caplets) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each caplet contains: calcium 27 mg ■ salicylate 99 mg ■ low sodium

■ store at room temperature

Questions or comments? Call 1-800-910-6874

24 505 0375 R00 C-002262-01-073-0000
Div. by Target Brands, Inc. MMS500
11/18/2024
Target Brands, Inc.

47970294

Varnish Omit Area

Varnish Omit

Varnish/Print Omit Area

NDC 11673-880-44

Compare to active ingredient in Pepto-Bismol® Caplets®

5-Symptom Digestive Relief

Bismuth Subsalicylate, 262 mg
Upset Stomach Reliever / Antidiarrheal

Relieves upset stomach, nausea, heartburn, indigestion and diarrhea



40 CAPLETS**
(**CAPSULE-SHAPED TABLETS)

Varnish Omit

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient Purpose (in each caplet)
Bismuth subsalicylate 262 mg.....Upset stomach reliever and antidiarrheal

Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

47580424

This product's minimum shelf life is 24 months. For more information, visit www.upandup.com.
© 2024 Target Brands, Inc. All rights reserved.

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts (continued)

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- swallow with water, do not chew
- adults and children 12 years and over: ■ 2 caplets (1 dose) every ½ hour or 4 caplets (2 doses) every hour as needed

Varnish Omit

24 505 0375 R00 C-002262-01-073-0000



3 11673 88044 9

Lot #: _____ Exp. Date: _____

Print/Varnish Omit Area

Drug Facts (continued)

For diarrhea

- 2 caplets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other Information

- each caplet contains: calcium 27 mg
- salicylate 99 mg
- low sodium
- store at room temperature

Inactive ingredients calcium carbonate, D&C red #27 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, povidone K-30, silicon dioxide, sodium starch glycolate

Questions or comments?
Call 1-800-910-6874

Distributed by Target Corporation
Minneapolis, MN 55409
TM & ©2024 Target Brands, Inc.

5-SYMPTOM DIGESTIVE RELIEF

bismuth subsalicylate tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-880

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE K30 (UNII: U725QWY32X)	
MANNITOL (UNII: 3OWL53L36A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

Product Characteristics

Color	pink	Score	no score
Shape	CAPSULE	Size	16mm
Flavor		Imprint Code	127
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-880-44	1 in 1 CARTON	06/15/2024	
1		40 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	06/15/2024	

Labeler - TARGET CORPORATION (006961700)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(11673-880)

Revised: 6/2024

TARGET CORPORATION