

**MUCINEX SINUS-MAX PRESSURE, PAIN AND COUGH MAXIMUM STRENGTH-
acetaminophen, dextromethorphan hydrobromide, guaifenesin, and
phenylephrine hydrochloride capsule, liquid filled
RB Health (US) LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

**Mucinex® Sinus-Max® Pressure, Pain & Cough
Maximum Strength**

Drug Facts

<i>Active ingredients (in each liquid gel)</i>	<i>Purposes</i>
Acetaminophen 325 mg	Pain reliever
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - cough
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Questions?**1-866-MUCINEX (1-866-682-4639)**

You may also report side effects to this phone number.

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224

PRINCIPAL DISPLAY PANEL - 16 Capsule Blister Pack Carton

Fast Dissolving Liquid Gels!

NDC 72854-202-16

MAXIMUM STRENGTH

Mucinex®
SINUS-MAX®

PRESSURE, PAIN
& COUGH

Acetaminophen – Pain Reliever
Dextromethorphan HBr - Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

- ✓ Relieves Sinus Pressure & Congestion
- ✓ Relieves Headache
- ✓ Controls Cough
- ✓ Thins & Loosens Mucus

Actual Size

LIQUID GELS
(Liquid Filled Capsules)

DAY TIME
FOR AGES 12+

MAXIMUM STRENGTH

Mucinex
SINUS-MAX

PRESSURE, PAIN
& COUGH

Tamper evident: Do not use if carton is damaged
or if printed seal on blister is broken or missing.

16 LIQUID GELS

Fast Dissolving Liquid Gels!

MAXIMUM STRENGTH

NDC 72854-202-16


Mucinex
SINUS-MAX

PRESSURE, PAIN
& COUGH

Acetaminophen – Pain Reliever
Dextromethorphan HBr - Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

- ✓ Relieves Sinus Pressure & Congestion
- ✓ Relieves Headache
- ✓ Controls Cough
- ✓ Thins & Loosens Mucus

Actual Size



16 LIQUID GELS
(Liquid Filled Capsules)

DAY TIME
FOR AGES 12+

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2021 RB Health

022721 3176794



HEALTH • HYGIENE • HOME

www.mucinex.com

Patents: www.rb.com/patents



LOT:

EXP.:

3176794



Drug Facts

Active ingredients (in each liquid gel)

Purposes

Acetaminophen 325 mg.....	Pain reliever
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - cough
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 12 liquid gels in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

MAXIMUM STRENGTH

Mucinex
SINUS-MAX

PRESSURE, PAIN
& COUGH

Maximum Strength per 4-hour dose
Do not take more than a total of
12 liquid gels in a 24-hour period.
Take only as directed.
Keep carton for full information.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

MUCINEX SINUS-MAX PRESSURE, PAIN AND COUGH MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-202
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C yellow no. 6 (UNII: H77VEI93A8)	
gelatin, unspecified (UNII: 2G86QN327L)	
glycerin (UNII: PDC6A3C0OX)	
hypromellose, unspecified (UNII: 3NXW29V3WO)	
isopropyl alcohol (UNII: ND2M416302)	
light mineral oil (UNII: N6K5787QVP)	
polyethylene glycol, unspecified (UNII: 3WJQ0SDW1A)	
povidone, unspecified (UNII: FZ989GH94E)	
propylene glycol (UNII: 6DC9Q167V3)	
titanium dioxide (UNII: 15FIX9V2JP)	
water (UNII: 059QF0KO0R)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	OVAL	Size	24mm
Flavor		Imprint Code	AR01
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-202-16	2 in 1 CARTON	07/26/2021	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination		

1		Product		
2	NDC:72854-202-08	1 in 1 CARTON	07/26/2021	
2		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL		part341	07/26/2021	

Labeler - RB Health (US) LLC (081049410)

Revised: 7/2021

RB Health (US) LLC