

UPSET STOMACH RELIEF-BISMUTH SUBSALICYLATE 262 MG- bismuth subsalicylate 262 mg capsule, liquid filled
Pioneer Life Sciences, LLC

Gencare-Upset Stomach Relief-Bismuth Subsalicylate 262 mg

Active Ingredient (in each softgel)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves:

■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- swallow with water, do not chew
- adults and children 12 years and over:
 - 2 softgels every ½ hour or 4 softgels every hour as needed for diarrhea
 - 2 softgels every ½ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 softgels) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information


- each softgel contains:
 - salicylate 99 mg
 - low sodium
- do not exceed 25°C

Inactive ingredients

Polyethylene Glycol-400, Polysorbate 80, Soya Lecithin, Povidone (PVPK 30), Gelatin, Glycerin, Sorbitol Sorbitan Solution, D & C Red 33 , Titanium Dioxide, Purified Water

Questions or Comments?

Call +1 (732) 994-2808 or email support@gencare.health

to compare to PEPTO BISMOL® LIQUICAPS® active ingredient


Upset Stomach Relief

Antidiarrheal

Bismuth Subsalicylate 262 mg

Symptom relief of:

- Nausea
- Heartburn
- Indigestion
- Diarrhea



48 softgels

TAMPER EVIDENT Do not use if safety seal under cap is broken or missing.

Drug Facts

Active Ingredient (in each softgel)
Bismuth subsalicylate 262 mg

Uses relieves:
 ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are
 ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

(continued under label)

Distributed by: **GenCare Consumer Products, LLC**
 40E Collers Ln, Suite A, East Brunswick, NJ 08816
PL0275-00



3 72090 00474 0

Lot No. Exp. Dt.

PEEL HERE

Drug Facts (cont)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- swallow with water, do not chew
- adults and children 12 years and over:
 ■ 2 softgels every ½ hour or 4 softgels every hour as needed for diarrhea
- 2 softgels every ½ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 softgels) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each softgel contains:
 ■ salicylate 99 mg
- low sodium
- do not exceed 25°C

Inactive ingredients

Polyethylene Glycol-400, Polysorbate 80, Soya Lecithin, Povidone (PVPK 30), Gelatin, Glycerin, Sorbitol, Sorbitan Solution, D & C Red 33, Titanium Dioxide, Purified Water

Questions or Comments?
 Call +1 (732) 994-2808 or email support@gencare.health

†This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Pepto Bismol® Liquicaps®

.5" Glue Area

UPSET STOMACH RELIEF-BISMUTH SUBSALICYLATE 262 MG			
bismuth subsalicylate 262 mg capsule, liquid filled			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72090-053
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)		BISMUTH SUBSALICYLATE	262 mg
Inactive Ingredients			
Ingredient Name			Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			
POLYSORBATE 80 (UNII: 6OZP39ZG8H)			
SOYBEAN LECITHIN (UNII: 1DI56QDM62)			
POVIDONE K30 (UNII: U725QWY32X)			

Gelatin (UNII: 2G86QN327L)				
Glycerin (UNII: PDC6A3C0OX)				
SORBITOL SOLUTION (UNII: 8KW3E207O2)				
SORBITAN (UNII: 6O92ICV9RU)				
D&C RED NO. 33 (UNII: 9DBA0SBB0L)				
Titanium Dioxide (UNII: 15FIX9V2JP)				
WATER (UNII: 059QF0KO0R)				
Product Characteristics				
Color	pink (Opaque baby pink)		Score	no score
Shape	OVAL (Oblong)		Size	20mm
Flavor			Imprint Code	SD30
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72090-053-01	48 in 1 BOTTLE; Type 0: Not a Combination Product	06/27/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M008	06/27/2025	

Labeler - Pioneer Life Sciences, LLC (014092742)

Revised: 6/2025

Pioneer Life Sciences, LLC