

**IBUPROFEN- ibuprofen tablet**  
**Walgreens**

-----  
**Ibuprofen 200**  
**Ibuprofen Tablets USP, 200mg**  
**Pain reliever/Fever reducer (NSAID)**

**Active ingredient (in each caplet)**

Ibuprofen 200 mg (NSAID)

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/ fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

**Allergy alert:**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives □ facial swelling □ asthma (wheezing)
- shock □ skin reddening □ rash □ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chances is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

### **Heart attack and stroke warning:**

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed

### **Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- you have symptoms of heart problems or stroke
  - chest pain • trouble breathing • weakness in one part or side of body • slurred speech • leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and older
- take 1 caplet every 4 to 6 hours while symptoms persist • If pain or fever does not respond to 1 caplet, 2 caplets may be used • do not exceed 6 caplets in 24 hours, unless directed by a doctor.
- children under 12 years
- ask a doctor

### **Other Information**

- store between 20 - 25°C (68 - 77°F).
- do not use if neck wrap or foil inner seal is broken or missing

### **Inactive ingredients**

colloidal silicon dioxide, FD&C Yellow #6, maize starch, polyethylene glycol, polyvinyl alcohol, povidone k30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

### **Questions or comments?**

Call 1-877- 770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST

### **Principal display panel**

**Walgreens**  WALGREENS STORES  
PHARMACY

Compare to the active ingredient in Motrin® IB

**EASY OPEN**  
**Ibuprofen 200**  
IBUPROFEN TABLETS USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)\*

**225** COATED CAPLETS\*\* (\*\*capsule-shaped tablets) **ACTUAL SIZE** 

**NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN**

**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING**  
**DO NOT USE IF FOIL INNER SEAL IS BROKEN OR MISSING**

**Drug Facts**  
**Active ingredient (in each caplet)** Ibuprofen 200 mg (NSAID)\* Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**  
temporarily relieves minor aches and pains due to:  
 ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache  
 ■ backache ■ the common cold ■ menstrual cramps  
 ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
 ■ skin redness ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product  
 ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
 ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if**  
 ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a

Walgreens.com NDC 0363-9593-21  
 MADE IN INDIA  
 Contains no ingredient made from 3 grain-containing grain (wheat, barley or rye). Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. \*This product is not manufactured or distributed by Henkel Inc., owner of the registered trademark Motrin® IB.  
 W3ORG0124-F

ITEM 712470 W10761-0721-F  
 11917 04698 3  
 Lot Exp. 20000004964 70000003712

**(CONTINUED ON BACK OF LABEL)**

label,  
ie.

**Inside (adhesive side)**

**Drug Facts (continued)**  
 history or stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
 ■ you experience any of the following signs of stomach bleeding: • feel faint or vomit blood • have bloody or black stools • have stomach pain that does not get better  
 ■ you have symptoms of heart problems or stroke: • chest pain • trouble breathing • weakness in one part or side of body • slurred speech  
 ■ leg swelling  
 ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area  
 ■ any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**  
 ■ do not take more than directed  
 ■ the smallest effective dose should be used

**Adults and children 12 years and older** ■ take 1 caplet every 4 to 6 hours while symptoms persist  
 ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used  
 ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

**Children under 12 years** ■ ask a doctor

**Other information** ■ store between 20-25°C (68-77°F)  
 ■ do not use if foil inner seal is broken or missing

**Inactive ingredients**  
 colloidal silicon dioxide, FD&C yellow #6, maize starch, polyethylene glycol, polyvinyl alcohol, polyvidone K30, pregelatinized starch, sodium starch glycolate, stearic acid, TBK, titanium dioxide

**Questions or comments?**  
 call 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST

**IBUPROFEN**

ibuprofen tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-9593
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	G2
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9593-21	225 in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202312	05/01/2024	

**Labeler** - Walgreens (008965063)

Revised: 12/2025

Walgreens