

**DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled
Strive Pharmaceuticals Inc.**

RIGHT REMEDIES DayTime Cold & Flu

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant
Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms: • nasal congestion • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • if you are now taking a prescription monoamine oxidase inhibitor (MAOI), (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if

your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless
• pain, nasal congestion or cough get worse or last more than 7 days
• fever gets worse or lasts more than 3 days
• redness or swelling is present
• new symptoms occur
• cough comes back, or occurs with rash or headache that lasts.
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away.

Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see **overdose warning**
- do not exceed 4 doses per 24 hrs

adults and children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12	ask a doctor
children under 4 yrs	do not use

Other information

- store at room temperature between 20-25°C (68-77°F)

Inactive ingredients

edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments

1-888-577-8033 Monday - Friday 8am - 4pm EST

Compare to the active ingredients of **Vicks® DayQuil™** Cold & Flu Relief LiquiCaps™*

RIGHT REMEDIES

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

*This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademarks Vicks® and DayQuil™.

Distributed by: Strive Pharmaceuticals Inc., East Brunswick, NJ 08816

Product of UAE

Packaged and Quality Assured in the USA

DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

REV.00-072022

Packaging

Compare to the active ingredients of **Vicks® DayQuil™ Cold & Flu Relief LiquiCaps™**

NDC 70692-807-48

RIGHT REMEDIES

DayTime Cold & Flu

Acetaminophen 325 mg
Pain reliever/ Fever reducer
Dextromethorphan HBr 10 mg
Cough Suppressant
Phenylephrine HCL 5 mg
Nasal Decongestant

48 softgels

temporarily relieves common cold/flu symptomatic headache, fever, sore throat, minor aches and pains, nasal congestion, and cough



RIGHT REMEDIES

DayTime Cold & Flu

Acetaminophen 325 mg
Pain reliever/ Fever reducer
Dextromethorphan HBr 10 mg
Cough Suppressant
Phenylephrine HCL 5 mg
Nasal Decongestant

temporarily relieves common cold/flu symptomatic headache, fever, sore throat, minor aches and pains, nasal congestion, and cough



READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/ Fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses
temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ cough due to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches & pains ■ fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hrs, which is the maximum daily amount for this product
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks daily while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI), (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough get worse or last more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF IMPAIRMENT



3 70692 00052 8

CT7069280748
REV.00-072022

Drug Facts (continued)

Directions
■ take only as directed - see overdose warning
■ do not exceed 4 doses per 24 hrs

adults and children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12	ask a doctor
children under 4 yrs	do not use

Other information
■ store at room temperature between 20-25°C (68-77°F)

Inactive ingredients
edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments
1-888-577-8033 Monday - Friday 8am - 4pm EST

*This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademarks Vicks® and DayQuil™.
Distributed by: Shire Pharmaceuticals Inc., East Brunswick, NJ 08816
Product of UAE
Packaged and Quality Assured in the USA

DRUG FACTS TABLE

Drug Facts	
Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/Fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant
Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ cough due to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches & pains ■ fever	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI), (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.	
When using this product, do not use more than directed.	
Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough get worse or last more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.	

Drug Facts (continued)	
Directions ■ take only as directed - see overdose warning ■ do not exceed 4 doses per 24 hrs	
adults and children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12	ask a doctor
children under 4 yrs	do not use
Other information ■ store at room temperature between 20-25°C (68-77°F)	
Inactive ingredients edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution	
Questions or comments 1-888-577-8033 Monday - Friday 8am - 4pm EST	

DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-807
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	807
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-807-48	4 in 1 CARTON	04/05/2023	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/05/2023	

Labeler - Strive Pharmaceuticals Inc. (080028013)

Revised: 12/2024

Strive Pharmaceuticals Inc.