

**DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled  
Strive Pharmaceuticals Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**RIGHT REMEDIES DayTime Cold & Flu**

**Drug Facts**

**Active ingredients (in each softgel)**

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Phenylephrine HCl 5 mg

**Purpose**

Pain reliever/Fever reducer  
Cough suppressant  
Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms: • nasal congestion • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a

doctor or pharmacist • if you are now taking a prescription monoamine oxidase inhibitor (MAOI), (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if** • you get nervous, dizzy or sleepless  
• pain, nasal congestion or cough get worse or last more than 7 days  
• fever gets worse or lasts more than 3 days  
• redness or swelling is present  
• new symptoms occur  
• cough comes back, or occurs with rash or headache that lasts.  
These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away.

Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed - see **overdose warning**
- do not exceed 4 doses per 24 hrs

adults and children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12	ask a doctor
children under 4 yrs	do not use

### **Other information**

- store at room temperature between 20-25°C (68-77°F)

### **Inactive ingredients**

edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

***Questions or comments***

**1-888-577-8033** Monday - Friday 8am - 4pm EST

Compare to the active ingredients of **Vicks® DayQuil™** Cold & Flu Relief LiquiCaps™\*

***RIGHT REMEDIES***

**READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

\*This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademarks Vicks® and DayQuil™.

Distributed by: Strive Pharmaceuticals Inc., East Brunswick, NJ 08816

Product of UAE

Packaged and Quality Assured in the USA

**DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

REV.00-072022

**Packaging**

Compare to the active ingredients of **Vicks® DayQuil™** Cold & Flu Relief LiquiCaps™

NDC 70692-807-48

**RIGHT REMEDIES**

# DayTime Cold & Flu

Acetaminophen 325 mg  
Pain reliever/ Fever reducer  
Dextromethorphan HBr 10 mg  
Cough Suppressant  
Phenylephrine HCL 5 mg  
Nasal Decongestant

**48 softgels**

**temporarily relieves common cold/flu symptomatic** headache, fever, sore throat, minor aches and pains, nasal congestion, and cough



**RIGHT REMEDIES**

# DayTime Cold & Flu

Acetaminophen 325 mg  
Pain reliever/ Fever reducer  
Dextromethorphan HBr 10 mg  
Cough Suppressant  
Phenylephrine HCL 5 mg  
Nasal Decongestant

**temporarily relieves common cold/flu symptomatic** headache, fever, sore throat, minor aches and pains, nasal congestion, and cough



**READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**Drug Facts**

Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/ Fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**  
temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ cough due to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches & pains ■ fever

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI), (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if** ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough get worse or last more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF IMPAIRMENT**



3 70692 00052 8

CT7069280748  
REV.00-072022

**Drug Facts (continued)**

**Directions**  
■ take only as directed - see overdose warning  
■ do not exceed 4 doses per 24 hrs

adults and children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12	ask a doctor
children under 4 yrs	do not use

**Other information**  
■ store at room temperature between 20-25°C (68-77°F)

**Inactive ingredients**  
edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

**Questions or comments**  
1-888-577-8033 Monday - Friday 8am - 4pm EST

\*This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademarks Vicks® and DayQuil™.  
Distributed by: Shire Pharmaceuticals Inc., East Brunswick, NJ 08816  
Product of UAE  
Packaged and Quality Assured in the USA

## DRUG FACTS TABLE

<b>Drug Facts</b>	
<b>Active ingredients (in each softgel)</b>	<b>Purpose</b>
Acetaminophen 325 mg.....	Pain reliever/Fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant
<b>Uses</b> temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ cough due to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches & pains ■ fever	
<b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product <b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. <b>Sore throat warning:</b> If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
<b>Do not use</b> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI), (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
<b>Ask a doctor before use if you have</b> ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema	
<b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</b>	
<b>When using this product, do not use more than directed.</b>	
<b>Stop use and ask a doctor if</b> ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough get worse or last more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.	
<b>If pregnant or breast-feeding, ask a health professional before use.</b> <b>Keep out of reach of children.</b> <b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.	

<b>Drug Facts (continued)</b>	
<b>Directions</b> ■ take only as directed - see overdose warning ■ do not exceed 4 doses per 24 hrs	
adults and children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12	ask a doctor
children under 4 yrs	do not use
<b>Other information</b> ■ store at room temperature between 20-25°C (68-77°F)	
<b>Inactive ingredients</b> edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution	
<b>Questions or comments</b> 1-888-577-8033 Monday - Friday 8am - 4pm EST	

# DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70692-807
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	807
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-807-48	4 in 1 CARTON	04/05/2023	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part341	04/05/2023	

**Labeler** - Strive Pharmaceuticals Inc. (080028013)

Revised: 5/2023

Strive Pharmaceuticals Inc.