

ERYTHROMYCIN- erythromycin ointment
ARMAS PHARMACEUTICALS INC.

Erythromycin Ophthalmic Ointment USP, 0.5%


Sterile

Rx only

DESCRIPTION:

Erythromycin Ophthalmic Ointment belongs to the macrolide group of antibiotics. It is basic and readily forms a salt when combined with an acid. The base, as crystals or powder, is slightly soluble in water, moderately soluble in ether, and readily soluble in alcohol or chloroform. Erythromycin ((3R*, 4S*, 5S*, 6R*, 7R*, 9R*, 11R*, 12R*, 13S*, 14R*)-4-[(2,6-dideoxy-3-C-methyl-3- O-methyl- α -L- *ribo*-hexopyranosyl)-oxy]-14-ethyl-7,12,13-trihydroxy-3,5,7,9,11,13-hexamethyl-6-[[3,4,6-trideoxy-3-(dimethyl-amino)- β -D-xylo-hexopyranosyl]oxy]oxacyclotetradecane-2,10-dione)) is an antibiotic produced from a strain of *Streptomyces erythraeus*.

It has the following structural formula:



Molecular Formula: C₂₇H₄₅NO₁₃
Molecular Weight: 733.94

Each gram contains Erythromycin USP 5 mg in a sterile ophthalmic base of mineral oil and white petrolatum.

CLINICAL PHARMACOLOGY:

Microbiology: Erythromycin inhibits protein synthesis without affecting nucleic acid synthesis. Erythromycin is usually active against the following organisms *in vitro* and in clinical infections:

Streptococcus pyogenes (group A β -hemolytic)

Alpha-hemolytic streptococci (viridans group)

Staphylococcus aureus, including penicillinase-producing strains (methicillin-resistant staphylococci are uniformly resistant to erythromycin)

Streptococcus pneumoniae

Mycoplasma pneumoniae (Eaton Agent, PPLO)

Haemophilus influenzae (not all strains of this organism are susceptible at the erythromycin concentrations ordinarily achieved)

Treponema pallidum

Corynebacterium diphtheriae

Neisseria gonorrhoeae

Chlamydia trachomatis

INDICATIONS AND USAGE:

For the treatment of superficial ocular infections involving the conjunctiva and/or cornea caused by organisms susceptible to erythromycin.

For prophylaxis of ophthalmia neonatorum due to *N. gonorrhoeae* or *C. trachomatis*.

The effectiveness of erythromycin in the prevention of ophthalmia caused by penicillinase-producing *N. gonorrhoeae* is not established.

For infants born to mothers with clinically apparent gonorrhea, intravenous or intramuscular injections of aqueous crystalline penicillin G should be given; a single dose of 50,000 units for term infants or 20,000 units for infants of low birth weight. Topical prophylaxis alone is inadequate for these infants.

CONTRAINDICATIONS:

This drug is contraindicated in patients with a history of hypersensitivity to erythromycin.

PRECAUTIONS:

General: The use of antimicrobial agents may be associated with the overgrowth of nonsusceptible organisms including fungi; in such a case, antibiotic administration should be stopped and appropriate measures taken.

Information for Patients: Avoid contaminating the applicator tip with material from the eye, fingers, or other source.

Carcinogenesis, Mutagenesis, Impairment of Fertility: Two year oral studies conducted in rats with erythromycin did not provide evidence of tumorigenicity. Mutagenicity studies have not been conducted. No evidence of impaired fertility that appeared related to erythromycin was reported in animal studies.

Pregnancy: Teratogenic Effects: Pregnancy Category B. Reproduction studies have been performed in rats, mice, and rabbits using erythromycin and its various salts and esters, at doses that were several multiples of the usual human dose. No evidence of harm to the fetus that appeared related to erythromycin was reported in these studies. There are, however, no adequate and well controlled studies in pregnant women. Because animal reproductive studies are not always predictive of human response, the erythromycins should be used during pregnancy only if clearly needed.

Nursing Mothers: Caution should be exercised when erythromycin is administered to a nursing woman.

Pediatric Use -See **INDICATIONS AND USAGE** and **DOSAGE AND ADMINISTRATION**.

Geriatric Use:No overall differences in safety or effectiveness have been observed between elderly and younger patients.

ADVERSE REACTIONS:

The most frequently reported adverse reactions are minor ocular irritations, redness, and hypersensitivity reactions.

DOSAGE AND ADMINISTRATION:

In the treatment of superficial ocular infections, a ribbon approximately 1 cm in length of Erythromycin Ophthalmic Ointment should be applied directly to the infected structure up to 6 times daily, depending on the severity of the infection.

For prophylaxis of neonatal gonococcal or chlamydial conjunctivitis, a ribbon of ointment approximately 1 cm in length should be instilled into each lower conjunctival sac. The ointment should not be flushed from the eye following instillation. A new tube should be used for each infant.

HOW SUPPLIED:

Sterile Erythromycin Ophthalmic Ointment USP, 0.5% is available as follows:

3.5 g (1/8 oz) sterile tamper-resistant tube (NDC 72485-670-35)

Carton of fifty (50) Unit Dose 1 g tube (NDC 72485-670-31)

STORAGE:Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Avoid excessive heat.

Protect from freezing.

Distributed by:

Armas Pharmaceuticals, Inc.

Freehold, NJ 07728 (USA)

Manufactured by:

Ophtapharm AG, Riethofstrasse 1, Hellingen,

8442, Switzerland (CHE) for Sentiss

Made in Switzerland

83008101 Rev. 07/23

Principal Display Panel Text for Container Label

NDC 72485-670-35

ERYTHROMYCIN OPHTHALMIC

OINTMENT USP, 0.5%

Rx only Sterile Net Wt. 3.5 g (1/8 oz.)

NDC 72485-670-35

**ERYTHROMYCIN OPHTHALMIC
OINTMENT USP, 0.5%**

Rx only Sterile Net Wt. 3.5 g (1/8 oz.)

Each Gram Contains: Erythromycin USP, 5 mg in a sterile ophthalmic base of mineral oil and white petrolatum.

Usual Dosage: See package insert for dosage information.

Storage: Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. KEEP TIGHTLY CLOSED.

See crimp for Lot Number and Expiration Date.

KEEP OUT OF REACH OF CHILDREN.

FOR OPHTHALMIC USE ONLY.

DO NOT USE IF CAP SEAL IS BROKEN

Distributed by:
Armas Pharmaceuticals, Inc.
Freehold, NJ 07728 (USA)
Made in Switzerland
Rev. 07/23


(01) 00372485670352

EXP

LOT



80008102



NDC 72485-670-31

ERYTHROMYCIN OPHTHALMIC

OINTMENT USP, 0.5%

Rx only Sterile Net Wt. 1 g

NDC 72485-670-31

**Erythromycin Ophthalmic
Ointment USP, 0.5% (5 mg/g)**

Rx only Sterile Net Wt. 1 g

DO NOT RE-USE. ONCE OPENED, DISCARD.

Each Gram Contains: Erythromycin USP, 5 mg in a sterile ophthalmic base of mineral oil and white petrolatum.

FOR OPHTHALMIC USE ONLY.

KEEP OUT OF REACH OF CHILDREN.

See crimp for Lot Number and Expiration Date.

DO NOT USE IF CAP SEAL IS BROKEN

Distributed by:
Armas Pharmaceuticals, Inc.
Freehold, NJ 07728 (USA)
Made in Switzerland
Rev. 07/23


(01) 00372485670314

EXP

LOT



80008202




Principal Display Panel Text for Carton Label

3.5 g NDC 72485-670-35

ERYTHROMYCIN OPHTHALMIC

OINTMENT USP, 0.5%

Rx only Sterile Net Wt 3.5 g (1/8 oz.)




1 g NDC 72485-670-31

ERYTHROMYCIN OPHTHALMIC

OINTMENT USP, 0.5%

Rx only Sterile Net Wt 1 g



ERYTHROMYCIN

erythromycin ointment

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:72485-670
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ERYTHROMYCIN (UNII: 63937KV33D) (ERYTHROMYCIN - UNII:63937KV33D)	ERYTHROMYCIN	5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
MINERAL OIL (UNII: T5L8T28FGP)	
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72485-670-35	1 in 1 CARTON	10/18/2023	
1		3.5 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:72485-670-31	50 in 1 CARTON	10/18/2023	

2	1 g in 1 TUBE; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA064030	10/18/2023	

Labeler - ARMAS PHARMACEUTICALS INC. (098405973)

Registrant - SENTISS AG (486920486)

Establishment

Name	Address	ID/FEI	Business Operations
Ophtapharm AG		482198285	analysis(72485-670) , pack(72485-670) , manufacture(72485-670) , label(72485-670)

Revised: 10/2023

ARMAS PHARMACEUTICALS INC.