OMEPRAZOLE- omeprazole tablet, orally disintegrating, delayed release Kroger Company

Kroger Co. Omeprazole Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs <u>2 or more</u> days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

• your heartburn continues or worsens

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- do not crush or chew tablets
- place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F); keep product out of high heat and moisture

Inactive ingredients

amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, crospovidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize

maltodextrin, mannitol, microcrystalline cellulose, propylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Package/Label Principal Display Panel

COMPARE TO PRILOSEC OTC[®] See side panel OUR PHARMACIST RECOMMENDED Omeprazole Delayed Release Orally Disintegrating Tablets 20 mg ACID REDUCER MELTech[™] Melts In Your Mouth **Treats FREQUENT Heartburn!** 24 HR MELTS IN YOUR MOUTH Dissolves without water actual size Strawberry Flavor 42 TABLETS Three 14-Day Courses of Treatment May Take 1 to 4 Days for Full Effect



Tips for Managing Heartburn
- Do not lie fict or bend over after eating
- Do not wear tight-fitting dathing
around the stranach
- Do not ext before bedtime
- Raise the head of your bed
- Avaid heartburn-cousing foods such
as rich, spicy, fatty or field foods,
chocoldze, caffeine, alcohol and
certain fulls and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit emoking

COMPARE TO PRILOSEC OTC® *See side panel



SAFETY FEATURE - DO NOT USE IF PRINTED TABLET BLISTER UNIT IS OPEN OR TORN.





OMEPRAZOLE						
omeprazole tablet, orally disintegrating, delayed release						
Product Information						
Product Type	HUMAN OTC DRUG	Item Code (Source) NDO		NDC:301	C:30142-394	
Route of Administration	ORAL					
Active Ingredient/Active	Moiety					
Ingredient Name		Basis of Strength		Strength		
OMEPRAZOLE (UNII: KG60484QX9) (OMEPRAZOLE - UNII:KG60484QX9)		OMEPRAZ OLE		20 mg		
					-	
Inactive Ingredients						
Ingredient Name					Strength	
DIMETHYLAMINOETHYL METHA COPOLYMER (UNII: 905HNO1SIH)	CRYLATE - BUTYL METHA	CRYLATE - MET	HYL METHACRY	LATE		
ASCORBIC ACID (UNII: PQ6CK8PD	OR)					
CETYL ALCOHOL (UNII: 936JST6JC	CN)					
SILICON DIOXIDE (UNII: ETJ7Z6XE	3U4)					
CROSPOVIDONE (15 MPA.S AT 5	5%) (UNII: 68401960MK)					
FERRIC OXIDE RED (UNII: 1K09F3	G675)					
HYPROMELLOSE, UNSPECIFIED	(UNII: 3NXW29V3WO)					

MANNITOL (UNII: 30WL53L36A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	RED (reddish)	Score	no score
Shape	ROUND	Size	9mm
Flavor	STRAWBERRY	Imprint Code	20
Contains			

Packaging

#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-394- 74	14 in 1 CARTON	04/10/2018	09/01/2022
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:30142-394- 55	3 in 1 CARTON	04/10/2018	
2		14 in 1 CARTON		
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
M	larketing	Information		
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NE	A	NDA209400	04/10/2018	

Labeler - Kroger Company (006999528)

Revised: 10/2023

Kroger Company