

**EQUATE ORASOL- 20% benzocaine gel  
WAL-MART STORES INC.**

---

**Maximum Strength Equate Orasol**

**Drug Facts**

**Active ingredient**

Benzocaine 20%

**Purpose**

Oral Pain Reliever

**Uses**

- temporary relief of occasional minor irritation, pain and sore mouth

**Warnings**

**Methemoglobinemia Warning:** Use of this product may cause Methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops;

- pale, gray, or blue colored skin(cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:** Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

**Do Not Use**

- more than directed
- for more than 7 days unless told to do so by a dentist or doctor

**Stop use and ask a doctor if**

- swelling, rash or fever develops
- irritation, pain, or redness persists or worsens

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- do not use tube if it is cut prior to opening
- cut open tip of tube on score mark
- use your fingertip or cotton applicator to apply a small pea-size amount of Oral Pain Relief Gel
- apply to affected area up to four times daily or as directed by a dentist or physician
- Adults and children 2 years of age and older: Apply to affected area
- Children under 12 years of age should be supervised in the use of this product
- Children under 2 years of age: Consult a doctor

### **Inactive ingredients**

Benzyl Alcohol, Carbomer, D&C Yellow No 10, FD&C Blue No 1, FD&C Red No 40, Flavor, Glycerin, Methylparaben, Polyethylene Glycol, Propylene Glycol, Sodium Saccharin

### **Other information**

store at a controlled room temperature 68°- 77°F (20°-25°C)

### **Principal Display Panel**

NDC 79903-966-09

Oral Pain Relief

MAXIMUM STRENGTH GEL

Benzocaine 20% NET WT. 0.33 OZ (9.35g)





**MAXIMUM STRENGTH**  
**Orasol**  
**Benzocaine 20%**  
**Oral Anesthetic Gel**  
**Instant Pain Relief**

NDC79903-966-09

NET WT  
0.33 OZ (9.35g)

**Active Ingredient** Benzocaine 20% **Purpose** Oral anesthetic **Uses** for the temporary relief of pain associated with: • toothache • canker sores • minor dental procedures • minor injury of the mouth and gums caused by dentures or orthodontic appliances **Warnings METHEMOGLOBINEMIA WARNING:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy **Allergy alert:** do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away. **Do not use:** • for teething • in children under 2 years of age **When using this product:** • avoid contact with the eyes • do not exceed recommended dosage • do not use for more than 7 days unless told to do so by a dentist or doctor **Stop use and ask a doctor if:** • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash or fever develops **Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions:** • adults and children 2 years of age or older: apply to affected area up to 4 times daily or as directed by doctor/dentist • children between 2 and 12 years of age: adult supervision should be given in use of this product • children under 2 years of age: do not use **Other information:** • do not use tube if it is cut prior to opening • to open tube, cut open tip of tube on score mark with scissors • store at a controlled room temperature 68°-77°F (20°-25°C) Questions? Call 1-888-287-1915

**DISTRIBUTED BY:** Walmart Inc., Bentonville, AR 72716

Satisfaction guaranteed - For questions or comments please call **1-888-287-1915**.

Not For Individual Retail Sale.

290131

734935

## EQUATE ORASOL

20% benzocaine gel

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79903-966
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>CARBOMER 940</b> (UNII: 4Q93RCW27E)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C00X)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	

**PROPYLENE GLYCOL** (UNII: 6DC9Q167V3)

**SACCHARIN SODIUM** (UNII: SB8ZUX40TY)

### Product Characteristics

Color		Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-966-09	1 in 1 CARTON	04/30/2024	
1		9.35 g in 1 TUBE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	04/30/2024	

**Labeler** - WAL-MART STORES INC. (051957769)

Revised: 1/2026

WAL-MART STORES INC.