

**ASPIRIN- aspirin tablet, coated
H-E-B**

HEB Aspirin 325 mg tablets

DRUG FACTS

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

INDICATIONS AND USAGE

Uses

for the temporary relief of minor aches and pains due to

- headache
- colds
- menstrual pain
- muscle pain
- toothache
- minor pain of arthritis

or as directed by your doctor

Warnings

Reye's syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning:This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs (aspirin,

ibuprofen, naproxen, or others)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have not been drinking fluids
- you have lost a lot of fluid due to vomiting or diarrhea

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug
- under a doctor's care for any serious condition

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

drink a full glass of water with each dose

- adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

PRINCIPAL DISPLAY PANEL

Adhesive Area

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless directly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions or comments? Call 1-877-290-4008

Inside

1/16"
Margin

Compare to Bayer® Genuine Aspirin active ingredient†

NDC 37808-174-53

Aspirin

325 mg (NSAID)

Pain Reliever/Fever Reducer

Safe Pain Relief

- Headaches
- Everyday Aches & Pains

actual size

500 COATED TABLETS

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscle pain
 - toothache
 - menstrual pain
- colds
- minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if clear or bloody vomit, nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Drug Facts (continued under label)

This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of Bayer® Genuine Aspirin.

MADE WITH PRIDE & CARE FOR H-E-B® 45260 - 2403

SAN ANTONIO, TX 78204

100% GUARANTEE

11R 0324

LOT: 1 11R 0324

EXP: 11/23/24

Varnish Omit Area

Outside

Drug Facts (continued)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding, warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:

- vomit blood
- have bloody or black stools
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions or comments? Call 1-877-290-4008

Adhesive Area

Inside

1/16" Margin

Varnish Omit Area

LOT:
EXP.:



43260 - 2403

MADE WITH PRIDE & CARE FOR H-E-B®

Distributor of Bayer® Genuine Aspirin.

SAN ANTONIO, TX 78204

100% GUARANTEE

11R 0324

Drug Facts (continued under label)

*This product is not manufactured or distributed by Bayer Healthcare LLC., distributor of Bayer® Genuine Aspirin.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

are age 60 or older

have had stomach ulcers or bleeding problems

take a blood thinning (anticoagulant) or steroid drug

take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

take 3 or more alcoholic drinks every day while using this product

take more or for a longer time than directed

may include:

■ facial swelling

■ asthma (wheezing)

■ shock

■ severe stomach bleeding

■ are age 60 or older

have had stomach ulcers or bleeding problems

take a blood thinning (anticoagulant) or steroid drug

take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

take 3 or more alcoholic drinks every day while using this product

take more or for a longer time than directed

may include:

■ facial swelling

■ asthma (wheezing)

■ shock

■ severe stomach bleeding

■ are age 60 or older

have had stomach ulcers or bleeding problems

take a blood thinning (anticoagulant) or steroid drug

take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

take 3 or more alcoholic drinks every day while using this product

take more or for a longer time than directed

may include:

■ facial swelling

■ asthma (wheezing)

■ shock

■ severe stomach bleeding

■ are age 60 or older

have had stomach ulcers or bleeding problems

take a blood thinning (anticoagulant) or steroid drug

take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

take 3 or more alcoholic drinks every day while using this product

take more or for a longer time than directed

may include:

Drug Facts

Active ingredient (in each tablet) Purposes

Aspirin 325 mg (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to:

- headache
- muscle pain
- toothache
- menstrual pain
- colds
- minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- facial swelling
- asthma (wheezing)
- shock
- severe stomach bleeding
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Compare to Bayer® Genuine Aspirin active ingredient!

NDC 37808-174-32



Aspirin
325 mg (NSAID)

Pain Reliever/Fever Reducer

Safe Pain Relief

- Headaches
- Everyday Aches & Pains



actual size

300 COATED TABLETS

Talk to your doctor or other healthcare provider before using this product for your heart. TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE UNPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.

Outside

Drug Facts (continued)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours or 3 tablets every 6 hours; not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions or comments? Call 1-877-290-4008

Adhesive Area

Inside

1/16" Margin

Compare to Bayer® Genuine Aspirin active ingredient¹

NDC 37808-174-73

H-E-B

Aspirin

325 mg (NSAID)

Pain Reliever/Fever Reducer

Safe Pain Relief

- Headaches
- Everyday Aches & Pains

actual size

100 COATED TABLETS

Talk to your doctor or other healthcare provider before using this product for your heart.

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)

Aspirin, 325 mg (NSAID)* Pain reliever/fever reducer

* nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscle pain
 - toothache
 - menstrual pain
 - colds
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from Chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued under label)

¹This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of Bayer® Genuine Aspirin.

MADE WITH PRIDE & CARE FOR H-E-B®

43260 - 2403

SAN ANTONIO, TX 78204

100% GUARANTEE

11R 0324

1 9 7 8 7 0 1 0 2 3 6 8 7

PEEL HERE

LOT:

EXP:

Varnish Omit Area

ASPIRIN

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-174
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL011
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-174-73	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2024	
2	NDC:37808-174-32	300 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2024	
3	NDC:37808-174-53	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2024	

Labeler - H-E-B (007924756)

Registrant - TIME CAP LABS INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABS INC		037052099	manufacture(37808-174)

Revised: 6/2024

H-E-B