

GLY-OXIDE- carbamide peroxide liquid
Medtech Products Inc.

Gly-Oxide

Drug Facts

Active ingredient

Carbamide peroxide 10% non USP*

*pH differs from USP specifications

Purposes

Oral debriding agent

Oral wound cleanser

Uses

- temporary use in cleansing:
 - canker sores
 - minor wounds or gum inflammation resulting from minor dental procedures, dentures, orthodontic appliances, accidental injury, or other irritations of the mouth and gums
- aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

Do not use

this product for more than 7 days unless directed by a dentist or doctor.

When using this product

- do not swallow
- avoid contact with eyes

Stop use and ask a doctor if:

- swelling, rash or fever develops
- irritation, pain, or redness persists or worsens
- sore mouth symptoms do not improve in 7 days

Keep out of the reach of children.

If more than used for rinsing is swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- replace tip on bottle when not in use

Adults and children 2 years and older:

- use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor

For direct application

- do not dilute
- apply several drops directly from the bottle onto affected area
- spit out after 2 to 3 minutes

As a mouthwash (oral rinse)

- place 10 to 20 drops onto tongue
- mix with saliva
- swish around in the mouth over the affected area for at least 1 minute and spit out

Children under 12 years: should be supervised in the use of this product

Children under 2 years: consult a dentist or doctor

Other information

- do not store above 25°C (77°F)
- store bottle in the outer carton

Inactive ingredients

citric acid, flavor, glycerin, propylene glycol, sodium lauroyl sarcosinate, water

Questions?

1-866-255-5202 weekdays

PRINCIPAL DISPLAY PANEL

Gly-Oxide LIQUID
(Carbamide peroxide 10%) Non USP*
ANTISEPTIC ORAL CLEANSER
2FL OZ (60 mL)

Read and retain carton for full product information.

Drug Facts	
Active ingredient	Purposes
Carbamide peroxide 10% non USP*	Oral debriding agent Oral wound cleanser
*pH differs from USP specifications.	
Uses	
• temporary use in cleansing:	
• canker sores	
• minor wounds or gum inflammation resulting from minor dental procedures, dentures, orthodontic appliances, accidental injury, or other irritations of the mouth and gums	
• aids in the removal of plaque, mucus, or other secretions associated with occasional sore mouth	
Warnings	
Do not use this product for more than 7 days unless directed by a dentist or doctor.	
When using this product	
• do not swallow	
• avoid contact with eyes	
Stop use and ask a doctor if	
• swelling, rash, or fever develops	
• irritation, pain, or redness persists or worsens	
• sore mouth symptoms do not improve in 7 days	
Keep out of reach of children.	
If more than used for rinsing is swallowed, get medical help or contact a Poison Control Center (1-800-322-1222) right away.	
Directions	
• replace cap on bottle when not in use	
Adults and children 2 years and older:	
• use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor	
For direct application	
• do not dilute	
• apply several drops directly from the bottle onto affected area	
• spit out after 2 to 3 minutes	



Carbamide peroxide 10%
Non USP*



Carbamide peroxide 10%
Non USP*



Carbamide peroxide 10%
Non USP*

Drug Facts (continued)

As a mouthwash (oral rinse)

- place 10 to 20 drops onto tongue
- mix with saliva
- swish around in the mouth over the affected area for at least 1 minute and spit out

Children under 12 years: should be supervised in the use of this product

Children under 2 years: consult a dentist or doctor

Other information

- do not store above 25°C (77°F)
- store bottle in the outer carton

Inactive ingredients
citric acid, flavor, glycerin, propylene glycol, sodium lauryl sarcosinate, water

Questions?
1-800-255-5202 weekdays

SAFE TRUSTED CARE EFFECTIVE

ANTISEPTIC ORAL CLEANSER

FOR MINOR MOUTH OR GUM IRRITATIONS

3 IN 1 ACTION

- ✓ **CLEANSES** minor mouth + gum irritation
- ✓ **SOOTHES** canker sores + braces irritation
- ✓ **KILLS** odor forming germs

UP TO 125 DOSES

2 FL OZ (60 mL)

TAMPER EVIDENT: Do Not Use if outer carton is open or torn.

Dist. by Medtech Products Inc.
Tarrytown, NY 10591
A Prestige Consumer Healthcare company.
©2021 Trade dress is owned by Medtech Products Inc.
All rights reserved.
Made in USA
GOU5003101



0 42037 10476 4

PRINCIPAL DISPLAY PANEL

Gly-Oxide LIQUID
(Carbamide peroxide 10%) Non USP*
ANTISEPTIC ORAL CLEANSER
0.5 FL OZ (15 mL)

Drug Facts
(continued)

onto affected area

- spit out after 2 to 3 minutes

As a mouthwash (oral rinse)

- place 10 to 20 drops onto tongue
- mix with saliva
- swish around in the mouth over the affected area for at least 1 minute and spit out

Children under 12 years: should be supervised in the use of this product

Children under 2 years: consult a dentist or doctor

Other information

- do not store above 25°C (77°F)
- store bottle in the outer carton

Inactive ingredients
citric acid, flavor, glycerin, propylene glycol, sodium lauryl sarcosinate, water

Questions?
1-866-255-5202 weekdays



Gly-Oxide
LIQUID
(Carbamide peroxide 10% Non USP)

ANTISEPTIC ORAL CLEANSER
FOR MINOR MOUTH OR GUM IRRITATIONS

3 IN 1 ACTION

- ✓ **CLEANSES** minor mouth + gum irritation
- ✓ **SOOTHES** canker sores + braces irritation
- ✓ **KILLS** odor forming germs

UP TO **30 DOSES**

0.5 FL OZ (15 mL)

Gly-Oxide
LIQUID

NOTE: Oversize package for ease in reading label information. Contains one bottle.

Tamper Evident: Do not use if outer carton is open or torn.

Dist. by Medtech Products Inc.
Tarrytown, NY 10591
A Prestige Consumer Healthcare company.
©2021 Trade dress is owned by Medtech Products Inc.
All rights reserved.
Made in USA
GOU5002101



Read and retain carton for full product information. Bottle image is actual size.

Drug Facts

Active ingredient	Purposes
Carbamide peroxide 10% non USP*	Oral debriding agent Oral wound cleanser
*pH differs from USP specifications	

Uses

- temporary use in cleansing:
 - canker sores
 - minor wounds or gum inflammation resulting from minor dental procedures, dentures, orthodontic appliances, accidental injury, or other irritations of the mouth and gums
 - aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

Do not use this product for more than 7 days unless directed by a dentist or doctor.

When using this product

- do not swallow
- avoid contact with eyes

Stop use and ask a doctor if

- swelling, rash, or fever develops
- irritation, pain, or redness persists or worsens
- sore mouth symptoms do not improve in 7 days

Keep out of reach of children.
If more than used for rinsing is swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- replace lip on bottle when not in use

Adults and children 2 years and older:

- use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor

For direct application

- do not dilute
- apply several drops directly from the bottle ▶

GLY-OXIDE

carbamide peroxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63029-500
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CARBAMIDE PEROXIDE (UNII: 31PZ2VAU81) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	CARBAMIDE PEROXIDE	100 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

SODIUM LAUROYL SARCOSINATE (UNII: 632GS99618)

WATER (UNII: 059QF0KO0R)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-500-01	1 in 1 CARTON	06/01/2012	
1		15 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:63029-500-02	1 in 1 CARTON	06/01/2012	
2		60 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	06/01/2012	

Labeler - Medtech Products Inc. (122715688)

Revised: 10/2024

Medtech Products Inc.