

ADULT LOW DOSE ENTERIC COATED ASPIRIN- aspirin tablet, delayed release Bi-Mart

Active ingredient (in each tablet)

Aspirin 81mg (NSAID)**

**nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headache or other symptoms needing immediate relief.**

Warnings

Reye's syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for:

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- **adults and children 12 years and over:**take 4 to 8 tablets every 4 hours; not to exceed 48 tablets in 24 hours unless directed by a doctor
- **children under 12 years:**consult a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Inactive ingredients

Acrycoat, Colloidal Silicon Dioxide, Croscarmellose Sodium, D&C Yellow# 10, Lactose, Microcrystalline Cellulose, Polysorbate 80, Sodium Hydroxide, Talc, Titanium Dioxide,

Triethyl Citrate, Yellow Oxide of Iron.

Questions?

call toll free - **1-844-912-4012**

Distributed by:

BI-MART

Eugene, OR 97402

PRINCIPAL DISPLAY PANEL-120 Count

NDC 37835-571-12

Compare to the active ingredients in Bayer[®] Aspirin Low Dose 81 mg*

Adult Low Dose Enteric Coated Aspirin

81 mg

Pain Reliever (NSAID)

For Aspirin Regimen Users

Enteric Safety Coated

120 Enteric Coated Tablets

*This product is not manufactured or distributed by the owner of the registered trademark Bayer[®].

Drug Facts (continued)

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic or you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs

if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose ■ **adults and children 12 years and over:** take 4 to 8 tablets every 4 hours, not to exceed 48 tablets in 24 hours unless directed by a doctor ■ **children under 12 years:** consult a doctor

Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients Acrycoat, Colloidal Silicon Dioxide, Croscarmellose Sodium, D&C Yellow #10, Lactose, Microcrystalline Cellulose, Polysorbate 80, Sodium Hydroxide, Talk, Titanium Dioxide, Triethyl Citrate, Yellow Oxide of Iron.

Questions? call toll-free 1-844-912-4012

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE: If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion for a full refund. This price will be cheerfully refunded.

Distributed by:
BI-MART
Eugene, OR 97402

HINGE

0 111161ZSEL4

Compare to the active ingredient in BAYER® ASPIRIN LOW DOSE 81 MG*

BI-MART NDC 37835-571-12

ADULT LOW DOSE ENTERIC COATED ASPIRIN PAIN RELIEVER (NSAID)

FOR ASPIRIN REGIMEN USERS ENTERIC SAFETY COATED

120 ENTERIC COATED TABLETS • 81 MG EACH

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

*This product is not manufactured or distributed by the owner of the registered trademark Bayer®

Drug Facts

Active ingredient (in each tablet)
Aspirin 81mg (NSAID)**Pain reliever
**nonsteroidal anti-inflammatory drug

Purpose
■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

PEEL FOR DIRECTIONS 571-12-BM 03/25

LIFT HERE

**PRINCIPAL DISPLAY PANEL-500 Count
NDC 37835-571-05**

Compare to the active ingredients in Bayer® Aspirin Low Dose 81 mg*

**Adult Low Dose Enteric Coated Aspirin
81 mg**

Pain Reliever (NSAID)

For Aspirin Regimen Users

Enteric Safety Coated

500 Enteric Coated Tablets

*This product is not manufactured or distributed by the owner of the registered trademark Bayer®.

01116 751910 3

Compare to the active ingredient
in **BAYER® ASPIRIN**
LOW DOSE 81 MG*

BI-MART NDC 37835-571-05
**ADULT
LOW DOSE
ENTERIC COATED
ASPIRIN**
PAIN RELIEVER (NSAID)

**FOR ASPIRIN REGIMEN USERS
ENTERIC SAFETY COATED**
**500 ENTERIC COATED
TABLETS • 81 MG EACH**

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts
Active ingredient (in each tablet) Purpose
Aspirin 81mg (NSAID)* Pain reliever
**nonsteroidal anti-inflammatory drug

Uses
■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

PEEL FOR DIRECTIONS 571-05-BM 03/25

LIFT HERE

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose ■ **adults and children 12 years and over:** take 4 to 8 tablets every 4 hours, not to exceed 48 tablets in 24 hours unless directed by a doctor ■ **children under 12 years:** consult a doctor

Other information
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients Acrycoat, Colloidal Silicon Dioxide, Croscarmellose Sodium, D&C Yellow #10, Lactose, Microcrystalline Cellulose, Polysorbate 80, Sodium Hydroxide, Talc, Titanium Dioxide, Triethyl Citrate, Yellow Oxide of Iron.

Questions? call toll-free 1-844-912-4012

*This product is not manufactured or distributed by the owner of the registered trademark Bayer®

Distributed by:
BI-MART
Eugene, OR 97402

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE
If you are not completely satisfied with your BI-MART QUALITY PRODUCT, return it to the store within the unused portion and your entire purchase price will be cheerfully refunded.

ADULT LOW DOSE ENTERIC COATED ASPIRIN

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37835-571
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	

D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
LACTOSE (UNII: J2B2A4N98G)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	5mm
Flavor		Imprint Code	A1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-571-12	120 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	
2	NDC:37835-571-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/01/2025	

Labeler - Bi-Mart (027630078)

Establishment

Name	Address	ID/FEI	Business Operations
ELYSIUM PHARMACEUTICALS LIMITED		915664486	manufacture(37835-571)

Revised: 3/2025

Bi-Mart