LORATADINE- loratadine tablet, chewable KROGER COMPANY

LORATADINE tablet, chewable

Active ingredient (in each chewable tablet)

Loratadine USP, 5 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings

Do not useif you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you haveliver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctorif an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or beast-feeding, ask a health professional before use.

Keep Out of Reach of Children

In case of overdose, get medical help or contact a Poison Control Center right away (1800-222-1222).

Directions

• chew or crush tablets completely before swallowing.

adults and children 6 years and over	chew 2 tablets daily; not more than 2	
	chewable tablets in 24 hours	

children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 chewable tablet in 24 hours
children under 2 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- phenylketonurics: contains phenylalanine 1.25 mg per tablet.
- TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.
- store between 20° to 25°C (68° to 77°F).

Inactive ingredients

aspartame, citric acid anhydrous, colloidal silicon dioxide, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Questions?

1800-632-6900

Package/Label Principal Display Panel



children's Allergy Relief

Loratadine Chewable Tablets USP, 5 mg Antihistamine



COMPARE TO the active ingredient of CHILDREN'S CLARITIN® CHEWABLES 'See bottom panel



Ages 2+ years

Non-Drowsy* children's Allergy Relief

Loratadine Chewable Tablets USP, 5 mg **Antihistamine**

CHEWABLE

Indoor and Outdoor Allergies

n taken as directed. See Drug Facts Pr

NDC 30142-565-30



24 RELIEF OF:

Sneezing, Runny Nose Itchy, Watery Eyes **Itchy Throat or Nose**

> DYE-FREE **Grape Flavor**

30 CHEWABLE **TABLETS**




Information, Scan UPC Using Your Kroger App or Call 800-632-6900 DISTRIBUTED BY THE KROGER CO. CINCINNATI, OHIO 45202 For More Product

†Claritin* is a registered trademark of Bayer Healthcare LLC. Bayer Healthcare LLC is not affiliated with The Kroger Co.

Our Brands, Our Guarantee, Love It or Your Money Back. www.kroger.com







Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose itchy, watery eyes sneezing
 - itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed.

Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions

chew or crush tablets completely before swallowing. adults and children chew 2 tablets daily; not more than 2 chewable tablets in 24 hours 6 years and over children 2 to under chew 1 tablet daily; not more than 1 chewable tablet in 24 hours 6 years of age children under ask a doctor 2 years of age consumers with liver ask a doctor or kidney disease

Other information

- phenylketonurics: contains phenylalanine 1.25 mg per
- TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.

 ■ store between 20° to 25°C (68° to 77°F).


Inactive ingredients

aspartame, citric acid anhydrous, colloidal silicon dioxide, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Questions or comments? 1-800-632-6900

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION. SEE END PANEL FOR EXPIRATION DATE.

CENE - NO COATING



Non

Varnish Area

loratadine tablet, chewable

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:30142-565

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PATABONE (UNIV. TAICORGION) (LORATADINE, LINU TAICORGION)	LODATABINE	_

LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN) LORATADINE 5 mg

Inactive Ingredients Ingredient Name

Ingredient Name Strength
ASPARTAME (UNII: Z0H242BBR1)

ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)

MAGNESIUM STEARATE (UNII: 70097M6I30)

MANNITOL (UNII: 30WL53L36A)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)

STEARIC ACID (UNII: 4ELV7Z65AP)

Product Characteristics

Color	white (white to off-white)	Score	no score
Shape	ROUND	Size	10mm
Flavor	GRAPE	Imprint Code	LC5
Contains			

Packaging

Ш	1 dekaging				
-	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
	NDC:30142- 565-30	3 in 1 CARTON	03/15/2024		
	1	10 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information

Marketing A	Application Number or Monograph Citation	Marketing Start	Marketing End
Category		Date	Date
ANDA AN	NDA210088	03/15/2024	

Establishment				
Name	Address	ID/FEI	Business Operations	
Ohm Laboratories Inc.		184769029	manufacture(30142-565)	

Revised: 3/2024 KROGER COMPANY