

METRONIDAZOLE VAGINAL- metronidazole gel

Saptalis Pharmaceuticals LLC

Metronidazole Vaginal Gel, 0.75%


Rx only

FOR INTRAVAGINAL USE ONLY

NOT FOR OPHTHALMIC, DERMAL, OR ORAL USE

DESCRIPTION

Metronidazole vaginal gel is the intravaginal dosage form of the synthetic antibacterial agent, metronidazole, USP at a concentration of 0.75%. Metronidazole is a member of the imidazole class of antibacterial agents and is classified therapeutically as an antiprotozoal and antibacterial agent. Chemically, metronidazole is a 2-methyl-5-nitroimidazole-1-ethanol. It has a molecular formula of $C_6H_9N_3O_3$, a molecular weight of 171.2 g/mole, and has the following structure:



Metronidazole vaginal gel is a gelled, purified water solution, containing metronidazole, USP at a concentration of 7.5 mg/g (0.75%). The gel is formulated at pH 4.0. The gel also contains carbomer homopolymer, edetate disodium, methylparaben, propylparaben, propylene glycol, and sodium hydroxide.

Each applicator full of 5 grams of vaginal gel contains approximately 37.5 mg of metronidazole.

CLINICAL PHARMACOLOGY

Normal Subjects

Following a single, intravaginal 5 g dose of metronidazole vaginal gel (equivalent to 37.5 mg of metronidazole) to 12 normal subjects, a mean maximum serum metronidazole concentration of 237 ng/mL was reported (range: 152 ng/mL to 368 ng/mL). This is approximately 2% of the mean maximum serum metronidazole concentration reported in the same subjects administered a single, oral 500 mg dose of metronidazole (mean C_{max} = 12,785 ng/mL, range: 10,013 ng/mL to 17,400 ng/mL). These peak concentrations were obtained in 6 hours to 12 hours after dosing with metronidazole vaginal gel and 1 hour to 3 hours after dosing with oral metronidazole.

The extent of exposure [area under the curve (AUC)] of metronidazole, when administered as a single intravaginal 5 g dose of metronidazole vaginal gel (equivalent to

37.5 mg of metronidazole), was approximately 4% of the AUC of a single oral 500 mg dose of metronidazole (4,977 ng-hr/mL and approximately 125,000 ng-hr/mL, respectively).

Dose-adjusted comparisons of AUCs demonstrated that, on a mg to mg comparison basis, the absorption of metronidazole, when administered vaginally, was approximately half that of an equivalent oral dosage.

Patients with Bacterial Vaginosis

Following single and multiple 5 g doses of metronidazole vaginal gel to 4 patients with bacterial vaginosis, a mean maximum serum metronidazole concentration of 214 ng/mL on day 1 and 294 ng/mL (range: 228 ng/mL to 349 ng/mL) on day five were reported. Steady-state metronidazole serum concentrations following oral dosages of 400 mg to 500 mg BID have been reported to range from 6,000 ng/mL to 20,000 ng/mL.

Microbiology

The intracellular targets of action of metronidazole on anaerobes are largely unknown. The 5-nitro group of metronidazole is reduced by metabolically active anaerobes, and studies have demonstrated that the reduced form of the drug interacts with bacterial DNA. However, it is not clear whether interaction with DNA alone is an important component in the bactericidal action of metronidazole on anaerobic organisms.

Culture and sensitivity testing of bacteria are not routinely performed to establish the diagnosis of bacterial vaginosis. (See **INDICATIONS AND USAGE**.)

Standard methodology for the susceptibility testing of the potential bacterial vaginosis pathogens, *Gardnerella vaginalis*, *Mobiluncusspp* ., and *Mycoplasma hominis*, has not been defined. Nonetheless, metronidazole is an antimicrobial agent active *in vitro* against most strains of the following organisms that have been reported to be associated with bacterial vaginosis:

Bacteroidesspp .

Gardnerella vaginalis

Mobiluncusspp .

Peptostreptococcusspp .

INDICATIONS AND USAGE

Metronidazole vaginal gel is indicated in the treatment of bacterial vaginosis (formerly referred to as *Haemophilusvagininitis*, *Gardnerellavagininitis*, nonspecific vaginitis, *Corynebacteriumvagininitis*, or anaerobic vaginosis).

NOTE For purposes of this indication, a clinical diagnosis of bacterial vaginosis is usually defined by the presence of a homogeneous vaginal discharge that (a) has a pH of greater than 4.5, (b) emits a "fishy" amine odor when mixed with a 10% KOH solution, and (c) contains clue cells on microscopic examination. Gram's stain results consistent with a diagnosis of bacterial vaginosis include (a) markedly reduced or absent *Lactobacillus* morphology, (b) predominance of *Gardnerella* morphotype, and (c) absent or few white blood cells.

Other pathogens commonly associated with vulvovaginitis, e.g., *Trichomonas vaginalis*,

Chlamydia trachomatis, *N. gonorrhoeae*, *Candida albicans*, and *Herpes simplexvirus* should be ruled out.

CONTRAINDICATIONS

Metronidazole vaginal gel is contraindicated in patients with a prior history of hypersensitivity to metronidazole, parabens, other ingredients of the formulation, or other nitroimidazole derivatives.

WARNINGS

Convulsive Seizures and Peripheral Neuropathy

Convulsive seizures and peripheral neuropathy, the latter characterized mainly by numbness or paresthesia of an extremity, have been reported in patients treated with oral or intravenous metronidazole. The appearance of abnormal neurologic signs demands the prompt discontinuation of metronidazole vaginal gel therapy.

Metronidazole vaginal gel should be administered with caution to patients with central nervous system diseases.

Psychotic Reactions

Psychotic reactions have been reported in alcoholic patients who were using oral metronidazole and disulfiram concurrently. Metronidazole vaginal gel should not be administered to patients who have taken disulfiram within the last 2 weeks.

PRECAUTIONS

Metronidazole vaginal gel affords minimal peak serum levels and systemic exposure (AUCs) of metronidazole compared to 500 mg oral metronidazole dosing. Although these lower levels of exposure are less likely to produce the common reactions seen with oral metronidazole, the possibility of these and other reactions cannot be excluded presently. Data from well-controlled trials directly comparing metronidazole administered orally to metronidazole administered vaginally are not available.

General

Patients with severe hepatic disease metabolize metronidazole slowly. This results in the accumulation of metronidazole and its metabolites in the plasma. Accordingly, for such patients, metronidazole vaginal gel should be administered cautiously.

Known or previously unrecognized vaginal candidiasis may present more prominent symptoms during therapy with metronidazole vaginal gel. Approximately 6% to 10% of patients treated with metronidazole vaginal gel developed symptomatic *Candidavaginitis* during or immediately after therapy.

Disulfiram-like reaction to alcohol has been reported with oral metronidazole, thus the possibility of such a reaction occurring while on metronidazole vaginal gel therapy cannot be excluded.

Metronidazole vaginal gel contains ingredients that may cause burning and irritation of the eye. In the event of accidental contact with the eye, rinse the eye with copious

amounts of cool tap water.

Information for the Patient

The patient should be cautioned about drinking alcohol while being treated with metronidazole vaginal gel. While blood levels are significantly lower with metronidazole vaginal gel than with usual doses of oral metronidazole, a possible interaction with alcohol cannot be excluded.

The patient should be instructed not to engage in vaginal intercourse during treatment with this product.

Drug Interactions

Oral metronidazole has been reported to potentiate the anticoagulant effect of warfarin and other coumarin anticoagulants, resulting in a prolongation of prothrombin time. This possible drug interaction should be considered when metronidazole vaginal gel is prescribed for patients on this type of anticoagulant therapy.

In patients stabilized on relatively high doses of lithium, short-term oral metronidazole therapy has been associated with elevation of serum lithium levels and, in a few cases, signs of lithium toxicity.

Use of cimetidine with oral metronidazole may prolong the half-life and decrease plasma clearance of metronidazole.

Drug/Laboratory Test Interactions

Metronidazole may interfere with certain types of determinations of serum chemistry values, such as aspartate aminotransferase (AST, SGOT), alanine aminotransferase (ALT, SGPT), lactate dehydrogenase (LDH), triglycerides, and glucose hexokinase. Values of zero may be observed. All of the assays in which interference has been reported involve enzymatic coupling of the assay to oxidation-reduction of nicotinamide-adenine dinucleotides (NAD + NADH). Interference is due to the similarity in absorbance peaks of NADH (340 nm) and metronidazole (322 nm) at pH 7.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Metronidazole has shown evidence of carcinogenic activity in a number of studies involving chronic oral administration in mice and rats. Prominent among the effects in the mouse was the promotion of pulmonary tumorigenesis. This has been observed in all six reported studies in that species, including one study in which the animals were dosed on an intermittent schedule (administration during every fourth week only). At very high dose levels (approximately 500 mg/kg/day), there was a statistically significant increase in the incidence of malignant liver tumors in males. Also, the published results of one of the mouse studies indicate an increase in the incidence of malignant lymphomas as well as pulmonary neoplasms associated with lifetime feeding of the drug. All these effects are statistically significant. Several long-term oral dosing studies in the rat have been completed. There were statistically significant increases in the incidence of various neoplasms, particularly in mammary and hepatic tumors, among female rats administered metronidazole over those noted in the concurrent female control groups. Two lifetime tumorigenicity studies in hamsters have been performed and reported to be negative.

These studies have not been conducted with 0.75% metronidazole vaginal gel, which would result in significantly lower systemic blood levels than those obtained with oral formulations.

Although metronidazole has shown mutagenic activity in a number of *in vitro* assay systems, studies in mammals (*in vivo*) have failed to demonstrate a potential for genetic damage.

Fertility studies have been performed in mice up to six times the recommended human oral dose (based on mg/m²) and have revealed no evidence of impaired fertility.

Pregnancy

Teratogenic Effects

There has been no experience to date with the use of metronidazole vaginal gel in pregnant patients. Metronidazole crosses the placental barrier and enters the fetal circulation rapidly. No fetotoxicity or teratogenicity was observed when metronidazole was administered orally to pregnant mice at six times the recommended human dose (based on mg/m²); however, in a single small study where the drug was administered intraperitoneally, some intrauterine deaths were observed. The relationship of these findings to the drug is unknown.

There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, and because metronidazole is a carcinogen in rodents, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

Specific studies of metronidazole levels in human milk following intravaginally administered metronidazole have not been performed. However, metronidazole is secreted in human milk in concentrations similar to those found in plasma following oral administration of metronidazole.

Because of the potential for tumorigenicity shown for metronidazole in mouse and rat studies, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness in children have not been established.

ADVERSE REACTIONS

Clinical Trials

There were no deaths or serious adverse events related to drug therapy in clinical trials involving 800 non-pregnant women who received metronidazole vaginal gel.

In a randomized, single-blind clinical trial of 505 non-pregnant women who received metronidazole vaginal gel once or twice a day, 2 patients (one from each regimen) discontinued therapy early due to drug-related adverse events. One patient discontinued

drug because of moderate abdominal cramping and loose stools, while the other patient discontinued drug because of mild vaginal burning. These symptoms resolved after discontinuation of drug.

Medical events judged to be related, probably related, or possibly related to administration of metronidazole vaginal gel once or twice a day were reported for 195/505 (39%) patients. The incidence of individual adverse reactions were not significantly different between the two regimens. Unless percentages are otherwise stipulated, the incidence of individual adverse reactions listed below was less than 1%:

Reproductive:

Vaginal discharge (12%),
Symptomatic *Candidacervicitis/vaginitis* (10%),
Vulva/vaginal irritative symptoms (9%),
Pelvic discomfort (3%).

Gastrointestinal:

Gastrointestinal discomfort (7%),
Nausea and/or vomiting (4%),
Unusual taste (2%),
Diarrhea/loose stools (1%),
Decreased appetite (1%),
Abdominal bloating/gas; thirsty, dry mouth.

Neurologic:

Headache (5%),
Dizziness (2%),
Depression.

Dermatologic:

Generalized itching or rash.

Other:

Unspecified cramping (1%),
Fatigue,
Darkened urine.

In previous clinical trials submitted for approved labeling of metronidazole vaginal gel the following was also reported:

Laboratory:

Increased/decreased white blood cell counts (1.7%).

Other Metronidazole Formulations

Other effects that have been reported in association with the use of **topical (dermal)** formulations of metronidazole include skin irritation, transient skin erythema, and mild skin dryness and burning. None of these adverse events exceeded an incidence of 2% of patients.

Metronidazole vaginal gel affords minimal peak serum levels and systemic exposure (AUC) of metronidazole compared to 500 mg oral metronidazole dosing. Although these lower levels of exposure are less likely to produce the common reactions seen with oral metronidazole, the possibility of these and other reactions cannot be excluded presently.

The following adverse reactions and altered laboratory tests have been reported with the **oral or parenteral** use of metronidazole:

Cardiovascular: Flattening of the T-wave may be seen in electrocardiographic tracings.

Central Nervous System: (See **WARNINGS**.) Headache, dizziness, syncope, ataxia, confusion, convulsive seizures, peripheral neuropathy, vertigo, incoordination, irritability, depression, weakness, insomnia.

Gastrointestinal: Abdominal discomfort, nausea, vomiting, diarrhea, an unpleasant metallic taste, anorexia, epigastric distress, abdominal cramping, constipation, "furry" tongue, glossitis, stomatitis, pancreatitis, and modification of taste of alcoholic beverages.

Genitourinary: Overgrowth of *Candida* in the vagina, dyspareunia, decreased libido, proctitis.

Hematopoietic: Reversible neutropenia, reversible thrombocytopenia.

Hypersensitivity Reactions: Urticaria; erythematous rash; flushing; nasal congestion; dryness of the mouth, vagina, or vulva; fever; pruritus; fleeting joint pains.

Renal: Dysuria, cystitis, polyuria, incontinence, a sense of pelvic pressure, darkened urine.

To report SUSPECTED ADVERSE REACTIONS, contact Saptalis Pharmaceuticals, LLC. at 1-833-727-8254 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

There is no human experience with overdosage of metronidazole vaginal gel. Vaginally applied metronidazole gel, 0.75% could be absorbed in sufficient amounts to produce systemic effects. (See **WARNINGS**.)

DOSAGE AND ADMINISTRATION

The recommended dose is one applicator full of metronidazole vaginal gel (approximately 5 grams containing approximately 37.5 mg of metronidazole) intravaginally once or twice a day for 5 days. For once-a-day dosing, metronidazole vaginal gel should be administered at bedtime.

HOW SUPPLIED

Metronidazole vaginal gel, 0.75% is a clear colorless to yellow gel with no visible grit or particles, supplied in a 70 g aluminum tube with HDPE cap and packaged with 5 vaginal applicators.

The NDC number for the 70 g tube is 71656-067-70.

Store at controlled room temperature 15°C to 30°C (59°F to 86°F). Protect from freezing.

Clinical Studies

In a randomized, single-blind clinical trial of non-pregnant women with bacterial vaginosis who received metronidazole vaginal gel daily for 5 days, the clinical cure rates for evaluable patients determined at 4 weeks after completion of therapy for the QD and BID regimens were 98/185 (53%) and 109/190 (57%), respectively.

Distributed by:

Saptalis Pharmaceuticals, LLC.

Hauppauge, NY 11788

MADE IN USA

Rev.11/2024-R2

Metronidazole Vaginal Gel, 0.75% DIRECTIONS FOR USE

1. Filling the applicator

- Remove cap and puncture metal seal on tube with the pointed tip of cap. (See Figure 1)
- Screw end of applicator onto tube. (See Figure 2)
- Slowly squeeze gel out of tube and into applicator. Plunger will stop when the applicator is full. (See Figure 3)
- Unscrew applicator and replace cap on tube.




Figure 1




Figure 2




Figure 3

2. Inserting the applicator

- The applicator may be inserted while lying on your back with your knees bent or in any comfortable position.
- Hold filled applicator by barrel, and gently insert into vagina as far as it will comfortably go. (See Figure 4)
- Slowly press the plunger until it stops to deposit gel into vagina and then withdraw the applicator.




Figure 4

3. Care of the applicator

If physician prescribes twice-a-day dosing:

- After use, pull the plunger out of the barrel. (See Figure 5)
- Wash both plunger and barrel in warm soapy water and rinse thoroughly.
- To reassemble applicator, gently push plunger back into barrel.




Figure 5

IMPORTANT

For once-a-day dosing, apply one applicator full at bedtime. For twice-a-day dosing, apply one applicator full each morning and evening for 5 days, or as directed by physician.

WARNINGS

- If significant irritation develops from the use of this medication, discontinue use and consult your physician.
- Do not use during pregnancy except under the supervision of a physician.
- Keep this and all medications out of reach of children.
- For vaginal use only. Not for use in the eyes, on the skin or in the mouth.

Store at room temperature. Avoid exposure to extreme heat or cold.

See end of carton and bottom of tube for lot number and expiration date.

Distributed by:

Saptalis Pharmaceuticals, LLC.

Hauppauge, NY 11788

MADE IN USA

Rev. 11/2024-R3

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

Package/Label Display Panel - 70 g Label

NDC 71656-067-70

Metronidazole Vaginal Gel, 0.75%

FOR INTRAVAGINAL USE ONLY

NOT FOR OPHTHALMIC, DERMAL, OR ORAL USE

Rx Only

Net Wt. 70 g

The image shows the principal display panel for a 70g tube of Metronidazole Vaginal Gel, 0.75%. The panel features a green and blue header bar at the top. Below the header, the NDC number 71656-067-70 is printed. The product name, Metronidazole Vaginal Gel, 0.75%, is prominently displayed in large, bold, black font. Below the product name, the text 'FOR INTRAVAGINAL USE ONLY' and 'NOT FOR OPHTHALMIC, DERMAL, OR ORAL USE' is printed in bold, black font. To the right of this text is a small black rectangular mark. Below the usage instructions, the text 'Rx only' and 'Net Wt. 70 g' is printed. The Saptalis logo, which consists of a green mountain range above the word 'SAPTALIS' in blue, is positioned to the right of the net weight. At the bottom of the panel, there is a section with detailed instructions: 'To open: Use top of cap to puncture seal at opening.', 'Important: The opening of this tube is covered by a metal tamper-resistant seal. If the seal has been punctured or is not visible, do not use this product.', 'Each gram contains: 7.5 mg of metronidazole, USP in a gel consisting of carbomer homopolymer, edetate disodium, methylparaben, propylparaben, propylene glycol, purified water, and sodium hydroxide. Formulated at pH 4.0.', 'Usual Dosage: Insert one applicator full (approximately 5 grams) of gel into the vagina once or twice daily for 5 days or as directed by a physician. For once-a-day dosing. Metronidazole vaginal gel, 0.75% should be administered at bedtime. Use applicators supplied in package.', 'Store at controlled room temperature 15°C to 30°C (59°F to 86°F). Protect from freezing. KEEP OUT OF REACH OF CHILDREN.', and 'See package insert for complete product information. See crimp end for lot number and expiration date.' To the right of these instructions, the distribution information is provided: 'Distributed by: Saptalis Pharmaceuticals, LLC. Hauppauge, NY 11788 MADE IN USA Rev. 11/24-R2 PPM-0045'. At the bottom right of the panel is a barcode with the numbers 3 71656 06770 0 printed below it.

Package/Label Display Panel - 70 g Carton

NDC 71656-067-70

Metronidazole Vaginal Gel, 0.75%

FOR INTRAVAGINAL USE ONLY

NOT FOR OPHTHALMIC, DERMAL, OR ORAL USE

Rx Only

Net Wt. 70 g



METRONIDAZOLE VAGINAL

metronidazole gel

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71656-067
Route of Administration	VAGINAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METRONIDAZOLE (UNII: 140QMO216E) (METRONIDAZOLE - UNII:140QMO216E)	METRONIDAZOLE	7.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
CARBOMER HOMOPOLYMER TYPE B (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: HHT01ZNK31)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71656-067-70	1 in 1 CARTON	03/03/2025	
1		70 g in 1 TUBE, WTH APPLICATOR; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA216750	03/03/2025	

Labeler - Saptalis Pharmaceuticals LLC (080145868)

Revised: 1/2025

Saptalis Pharmaceuticals LLC