

**MOMMYS BLISS CHILDRENS PAIN AND FEVER- acetaminophen suspension
Mom Enterprises, LLC.**

Mommys Bliss Childrens Pain and Fever

Inactive ingredients:

anhydrous citric acid, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, natural flavors, organic elderberry juice concentrate, purified water, sodium benzoate, sucrose, xanthan gum

USE:

Purpose: Pain reliever/fever reducer

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Directions and Dosing:

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL=milliliter
- find right dose on chart below
- if possible, use weight to dose; otherwise, use age
- remove cap and seal, then insert the adapter plug by pushing down firmly until a tight seal is formed
- top of plug should be flush with bottle neck and not removeable
- squeeze dose into the provided dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

Dosing:

Weight (lb)	Age (yr)	Dose (mL)*
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL

60-71 lbs 9-10 years 12.5 mL

72-95 lbs 11 years 15 mL

*or as directed by a doctor

Attention: use only enclosed cup specifically designed for use with this product. Do not use any other dosing device.

Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription.) If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease

Ask a doctor before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if:

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Prompt medical attention is critical even if you do not notice any signs or symptoms.

Active ingredient:

Active ingredient: (in each 5 mL) Acetaminophen 160mg

WARNINGS:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription.) If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease

Ask a doctor before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if:

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Prompt medical attention is critical even if you do not notice any signs or symptoms.

Keep out of reach of Children.

Keep out of reach of children.

Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Prompt medical attention is critical even if you do not notice any signs or symptoms.

Purpose and Use:

Purpose:

Pain reliever/fever reducer

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Children's Pain and Fever Acetaminophen 160mg per 5mL oral suspension

Pain Reliever - Fever Reducer, 4 fl oz (118 mL) bottle in carton, Ages 2-11 years, Berry Flavor, Dosing Cup Included

NEW Medicine for Peace of Mind



mommy's BLISS

CHILDREN'S

pain & fever

acetaminophen
160 mg per 5 mL

Oral Suspension | Pain Reliever-Fever Reducer

Free Of:

- ✓ Dyes
- ✓ Alcohol
- ✓ Artificial flavors
- ✓ Artificial sweeteners
- ✓ High-fructose corn syrup
- ✓ Top 9 allergens*

clean label PROJECT CERTIFIED

Yummy Berry Flavor

DOSING CUP INCLUDED

4 FL OZ (118 mL)

AGES 2-11 YEARS

NDC 71444-703-01

Drug Facts

Active Ingredient **Purpose**
(In each 5 mL)
Acetaminophen 160mg.....Pain reliever/
fever reducer

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription.) If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease

Ask a doctor before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if:

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

These could be signs of a serious condition.

NDC 71444-703-01 **Drug Facts (continued)**

Drug Facts (continued)

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Prompt medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL=milliliter
- find right dose on chart below
- if possible, use weight to dose; otherwise, use age
- remove cap and seal, then insert the adapter plug by pushing down firmly until a tight seal is formed
- top of plug should be flush with bottle neck and not removable
- squeeze dose into the provided dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

Weight (lb)	Age (yr)	Dose (mL)*
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-83 lbs	11 years	15 mL

*or as directed by a doctor
Attention: use only enclosed cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton or bottle seal is broken or missing

Inactive ingredients
anhydrous citric acid, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, natural flavors, organic elderberry juice concentrate, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
call weekdays 9AM to 5PM PST at 1-877-457-4953 (toll-free)

Distributed by MOM Enterprises, LLC.
Richmond, CA 94804, USA | 1-877-457-4953
mommysbliss.com

Medicine for Peace of Mind

As moms, we want to quickly make things "all better" for our little ones, without compromising on the quality of the products we give them. When I became a mom, I saw an opportunity to create safe and effective pain relief that you can truly feel good about - no dyes or artificial sweeteners, delicious tasting and with a commitment to being planet conscious. To me, this is peace of mind.

- Yasmin Kaderali | Mom & CEO

CHILDREN'S
Pain & Fever

Safe and effective medicine meets yummy natural berry flavor - free of unnecessary additives.

Free Of:

- ✓ Dyes
- ✓ Alcohol
- ✓ Parabens
- ✓ Artificial flavors
- ✓ Artificial sweeteners
- ✓ High-fructose corn syrup
- ✓ Top 9 allergens

clean label PROJECT CERTIFIED

✓ Peanuts	✓ Milk	✓ Tree Nuts
✓ Wheat	✓ Fish	✓ Soy
✓ Eggs	✓ Shellfish	✓ Sesame

MOM
PLANET→FWD™
Founded + Led

See all of our health and wellness products at: www.mommysbliss.com

Free Of:

- ✓ Dyes
- ✓ Alcohol
- ✓ Parabens
- ✓ Artificial flavors & sweeteners
- ✓ High-fructose corn syrup
- ✓ Top 9 allergens†

† Top 9 allergens are peanuts, wheat, eggs, milk, fish, shellfish, tree nuts, soy, and sesame.



Safe and effective medicine meets yummy natural berry flavor - free of unnecessary additives.



acetaminophen (160 mg per 5 mL)

Oral Suspension
Pain Reliever-Fever Reducer

Yummy Berry Flavor

4 FL OZ (118 mL)

AGE 2-11 YEARS

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL=milliliter
- find right dose on chart below
- if possible, use weight to dose; otherwise, use age
- remove cap and seal, then insert the adapter plug by pushing down firmly until a tight seal is formed
- top of plug should be flush with bottle neck and not removable
- squeeze dose into the provided dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor
Attention: use only enclosed cup specifically designed for use with this product. Do not use any other dosing device.

Open to read complete product information

Drug Facts

Active ingredient (in each 5 mL) Purpose (in each 5 mL)

Acetaminophen 160mg Pain reliever/fever reducer

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug

or nonprescription.) if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease

Ask a doctor before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if:

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

These could be signs of a serious condition.



MOMMYS BLISS CHILDRENS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71444-793
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM BENZOATE (UNII: OJ245FE5EU)	
GLYCERIN (UNII: PDC6A3C0OX)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
WATER (UNII: 059QF0KO0R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	
AMERICAN ELDERBERRY (UNII: 7610Y219OC)	

Product Characteristics

Color	purple (purple to brown liquid)	Score	
Shape		Size	
Flavor	BERRY (berry flavor)	Imprint Code	

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71444-793-02	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/19/2024	
2	NDC:71444-793-01	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/19/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/19/2024	

Labeler - Mom Enterprises, LLC. (135903776)

Revised: 4/2024

Mom Enterprises, LLC.