

PERCY MEDICINE- bismuth subsalicylate liquid
Merrick Medicine Company, Inc

PERCY MEDICINE

Drug Facts Active Ingredient

Active Ingredients

Bismuth Subsalicylate

Purpose

Antidiarrheal

Warnings

Children and teenagers that have or are recovering from Chicken Pox, flu systems, or flue should NOT use this product. If nausea, vomiting, or fever occur, consult a doctor because these symptoms could be an early sign of Reye Syndrome, a rare but serious condition.

Allergy Alert: Contains Salicylates. Do Not take if you are - allergic to salicylates (including aspirin) - taking other salicylate products.

Do not use if you have - blood or black stool - an ulcer - bleeding problem.

Ask a doctor before use if you are taking any drug for anti coagulation (thinning of the blood), diabetes, gout, arthritis. - When using this product a temporary but harmless darkening of the stool and/or tongue may occur.

Stop use and consult a doctor if - symptoms get worse - ringing in the ears or loss of hearing occurs - diarrhea lasts more that 2 days.

if pregnant or breast feeding *ask a professional before use*. In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

Inactive Ingredients

INACTIVE INGREDIENTS: CALCIUM HYDROXIDE, CITRIC ACID, ETHYL ALCOHOL, GLYCERINE, GUM ARABIC, OIL OF CINNAMON, OIL OF ORANGE, POTASSIUM CARBONATE, RHUBARB FLUID EXTRACT, SUGAR, PURIFIED WATER

Directions

(shake well before using)

Adults and children 12 yrs and over : 2 teaspoonfuls every hour as needed , not to

exceed 4 doses in 24 hours and not more than 2 days.

Children under 12: Consult a doctor.

Keep out of reach of children

Keep out of reach of children.

Uses:

Relieves diarrhea, reduces number of bowel movements.

Helps firm stool.

Relieves nausea, heartburn, fullness due to overindulgence of food and drink.

Package Label

=====
**PERCY MEDICINE
FOR DIARRHEA**

**MANUFACTURED FOR
MERRICK MEDICINE COMPANY
WACO, TEXAS 76703
info@percymedicine.net
1-254-753-3461**

Carton Label

PERCY[®] MEDICINE

FOR DIARRHEA
PARA DIARREA

10 mL / 4 fl oz

UPC / NDC 0322-2222-03

PERCY[®] MEDICINE

SINCE 1900

FOR DIARRHEA
PARA DIARREA

BELIEVES UPSET STOMACH DUE TO
OVERINDULGENCE IN FOOD & DRINK

Relieves Diarrhea

Helps Firm Stool

Reduces Number of Bowel Movements

Relieves Travelers' Diarrhea

See Back Panel for All Ingredients

ALCOHOL 5%

10 mL / 4 fl oz

Drug Facts

Active ingredient	Purpose
Bismuth subsalicylate, 18% (25 mg/mL)	Antidiarrheal

Uses

- Relieves Diarrhea • Reduces the number of bowel movements
- Relieves upset stomach due to overindulgence in food and drink
- Helps firm stool • Relieves Travelers' Diarrhea

Warnings

- Children and teenagers who have or are recovering from Chicken Pox or Shingles symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness • Ask doctor before use if you have • liver • kidney • stomach • or other conditions

Warnings: Do not take if you are • allergic to salicylates (including aspirin) • taking other salicylate products. Do not use if you have • bloody or black stool • an ulcer • bleeding problems

Ask a doctor or pharmacist before use if you are taking any drug for • acid • indigestion (burning the stomach) • diabetes • gout • arthritis

When using this product a temporary but harmless darkening of the stool and/or tongue may occur

Stop use and consult a doctor if • symptoms get worse • diarrhea lasts for more than two days • ringing in the ears or loss of hearing occurs.

If pregnant or breast feeding ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

Directions: (shake well before using)
• Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.

• Adults and Children 12 and up: 2 teaspoons every hour as needed, not to exceed 4 doses in 24 hours and not more than 2 days.
• Children under 12: Consult a doctor

Other Information
• Store at room temperature. Protect from freezing and excessive heat (over 40°F or 104°F)
• Each teaspoon (5 mL) contains: Bismuth: 6 mg, total salicylate: 225 mg

TAMPER EVIDENT:
Do not use if inner cap seal is broken or missing.

Inactive Ingredients: Calcium Hydroxide, Citric Acid, Ethyl Alcohol, Glycerin, Gum Arabic, Citric Acid, Citric Acid, Potassium Carbonate, Potassium Bicarbonate, Sage, Purified Water

MANUFACTURED FOR
MERRICK MEDICINE COMPANY
WACO, TEXAS 76703
Tel: 800-999-8888 ext. 1
1-354-753-3461

Rev 10/24



SHAKE WELL BEFORE USING
For relief from Diarrhea

For children 12 years of age and over • 1050 milligrams (2 teaspoons) every hour as needed
Adults • Do not exceed 4200 milligrams or 4 doses in 24 hours.
Use until Diarrhea stops but not more than 2 days

Children under 12 • Ask a doctor

Bottle Label

Drug Facts			<i>If pregnant or breast feeding</i> ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center immediately.
Active ingredient Bismuth Subsalicylate, 10% 525 mg/tsp.....	Purpose Antidiarrheal		Directions (shake well before using) • Drink plenty of clear fluids to help prevent dehydration caused by diarrhea • Adults and Children 12 and up: 2 teaspoonfuls every hour as needed, not to exceed 4 doses in 24 hours and not more than 2 days. • Children under 12: Consult a doctor
Uses • Relieves diarrhea • Reduces the number of bowel movements • Relieves upset stomach due to overindulgences in food and drink • Helps firm stool • Relieves Travelers' Diarrhea			Other Information • Store at room temperature. Protect from freezing and excessive heat (over 40°F or 104°C) • Each tablespoon (5 mL) contains: Calcium: 8 mg, total salicylate: 200 mg TAMPER EVIDENT: Do not use if inner cap seal is broken or missing.
Warnings • Children and teenagers who have or are recovering from Chicken Pox or flu-like symptoms should not use this product. When using this if product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness • Ask doctor before use if you have • fever • mucus in the stool			Inactive ingredients • Calcium Hydroxide, Citric Acid, Ethyl Alcohol, Glycerin, Gum Arabic, Oil of Cinnamon, Oil of Orange, Potassium Carbonate, Rhubarb Fluid Extract, Sugar, Purified Water
Allergy alert: Do not take if you are • allergic to salicylates (including aspirin) • taking other salicylate products. Do not use if you have • bloody or black stool • an ulcer • bleeding problem			 MANUFACTURED FOR MERRICK MEDICINE COMPANY WACO, TEXAS 76703 • info@percymedicine.net 1-254-753-3461
Ask a doctor or pharmacist before use if you are taking any drug for • anti coagulation (thinning the blood) • diabetes • gout • arthritis			
When using this product a temporary but harmless darkening of the stool and/or tongue may occur			
Stop use and consult a doctor if • symptoms get worse • diarrhea lasts for more than two days • ringing in the ears or loss of hearing occurs.			

res

PERCY MEDICINE

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0322-2222
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	1050 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KOOR)	
CALCIUM HYDROXIDE (UNII: PF5DZ W74VN)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
ALCOHOL (UNII: 3K9958V90M)	
GLYCERIN (UNII: PDC6A3C0OX)	
ACACIA (UNII: 5C5403N26O)	
ORANGE OIL (UNII: AKN3KSD11B)	
CINNAMON OIL (UNII: E5GY4I6YCZ)	
POTASSIUM CARBONATE (UNII: BQN1B9B9HA)	
RHUBARB (UNII: G280W4MW6E)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0322-2222-00	1 in 1 BOX	09/09/2019	01/01/2025
1	NDC:0322-2222-03	90 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0322-2222-03	90 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/09/2019	01/01/2025
3	NDC:0322-2222-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/09/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	09/09/2019	

Labeler - Merrick Medicine Company, Inc (007331838)

Registrant - Pharma Nobis, LLC (118564114)

Establishment

Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	analysis(0322-2222) , manufacture(0322-2222) , pack(0322-2222) , label(0322-2222)

Revised: 11/2025

Merrick Medicine Company, Inc