

ACETAMINOPHEN- acetaminophen tablet
PUBLIX SUPERMARKETS, INC

Extra strength
Pain Relief

- **Pain reliever/fever reducer**
 - **For adults**
 - **Contains no aspirin**
- ACETAMINOPHEN, 500 mg**

Active ingredient (in each caplet)

Acetaminophen USP, 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use

Keep out of reach of children

Keep out of reach of children

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F). See USP Controlled Room Temperature.
- see end panel for lot number and expiration date

Inactive ingredients

hydroxypropyl methyl cellulose, polyethylene glycol, povidone k-30, pregelatinized starch, sodium starch glycolate, stearic acid

Questions or comments?

call **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST

Principal display panel



COATING
FREE AREA



READ AND KEEP CARTON FOR COMPLETE WARNINGS
AND INFORMATION

**TAMPER EVIDENT DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen USP, 500 mg... Pain reliever/fever reducer

Uses

- Temporarily relieve minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - menstrual and menstrual cramps
- Temporarily reduce fever

Warnings

- Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if you take:
 - more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more 800 mg tablets every day while using this product
- Allergy alerts:** acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin redness
 - hives
 - itchIf a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning).
- adults and children 12 years and over:
 - take 2 caplets every 6 hours while symptoms last.
 - do not take more than 8 caplets in 24 hours, unless directed by a doctor.
 - do not use for more than 10 days unless directed by a doctor.
- children under 12 years:
 - ask a doctor.

Other information

- Store between 20-25°C (68-77°F). See USP Controlled Room Temperature.
- See and read label for lot number and expiration date.

Inactive ingredients

hydroxypropyl methylcellulose, polyethylene glycol, povidone K15, pregelatinized starch, sodium starch glycolate, stearic acid

Questions or comments?

1-877-770-3388 Mon-Fri 9:00 AM EST to 5:00 PM PST

This product is not manufactured or distributed by Kenwa Inc., owner of the registered trademark Extra Strength Tylenol® Caplets.

DISTRIBUTED BY
PUBLIX SUPER MARKETS, INC.
3300 PUBLIX CORPORATE PARKWAY,
LAKE AND, FL 33011
1-888-263-2637 publix.com
PUBLIX GUARANTEE: COMPLETE
SATISFACTION OR YOUR MONEY BACK



**Extra Strength
Pain Relief**
ACETAMINOPHEN, 500 mg

- Pain reliever/fever reducer
- For adults
- Contains no aspirin

**Extra Strength
Pain Relief**
ACETAMINOPHEN, 500 mg

- Pain reliever/fever reducer
- For adults
- Contains no aspirin

100 Caplets!
Caplets-shaped tablets

Compare to the active ingredient in Extra Strength Tylenol® Caplets

ACTUAL SIZE



NDC 41415-177-10

COATING
FREE AREA



Publix.

Lot
Exp.

COATING
FREE AREA

**Extra Strength
Pain Relief**
ACETAMINOPHEN, 500 mg

- Pain reliever/fever reducer
- For adults
- Contains no aspirin

100 Caplets!
Caplets-shaped tablets

Compare to the active ingredient in Extra Strength Tylenol® Caplets

ACTUAL SIZE



NDC 41415-177-10



ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41415-177
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients	
Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics			
Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	G;551
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41415-177-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	08/02/2024	
2	NDC:41415-177-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/02/2024	
3	NDC:41415-177-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	08/02/2024	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/02/2024	

Labeler - PUBLIX SUPERMARKETS, INC (006922009)