

ACETAMINOPHEN- acetaminophen tablet
PUBLIX SUPERMARKETS, INC

Extra Strength
Pain Relief

ACETAMINOPHEN, 500 mg

- **Pain reliever/fever reducer**
- **Contains no aspirin**
- **For adults**

Active ingredient (in each coated tablet)

Acetaminophen USP, 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold ■ headache
- backache ■ minor pain of arthritis
- toothache ■ muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

if a skin reaction occurs, stop use and seek medical help right away

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition

If pregnant or breast-feeding

ask a health professional before use

Keep out of reach of children

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 tablets every 6 hours while symptoms last
 - do not take more than 6 tablets in 24 hours, unless directed by a doctor
 - do not use for more than 10 days unless directed by a doctor
- children under 12 years

- ask a doctor

Other information

- store at 20-25°C (68-77°F). See USP Controlled Room Temperature.
- see end panel for lot number and expiration date

Inactive ingredients

FD&C red no. 40, FD&C yellow no. 6, lecithin, polyethylene glycol, polyvinyl alcohol, povidone k-30, pregelatinized starch, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

call **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST

Principal display panel

COATING FREE AREA

700000003737
200000005004

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts
Active ingredient (in each coated tablet) **Purpose**
Acetaminophen USP 500 mg. Pain reliever/fever reducer.

Uses
temporarily relieves minor aches and pains due to:
the common cold, headache, backache, minor pain of arthritis, backache, muscular aches, premenstrual and menstrual cramps, temporary reduces fever.

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
more than 4,000 mg of acetaminophen in 24 hours
with other drugs containing acetaminophen
3 or more alcoholic drinks every day while using this product.
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
skin reddening, hives, rash, blisters, or swelling. If a skin reaction occurs, stop use and seek medical help right away.
Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
if you are allergic to acetaminophen or any of the inactive ingredients in this product.
Ask a doctor before use if you have liver disease.
Ask a doctor or pharmacist before use if you are taking the

Drug Facts (continued)
blood thinning drug warfarin.
Stop use and ask a doctor if
pain gets worse or lasts more than 10 days
fever gets worse or lasts more than 3 days
new symptoms occur
redness or swelling is present. These could be signs of a serious condition.
if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-322-2222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
do not take more than directed (see overdose warning).
adults and children 12 years and over: take 2 tablets every 6 hours while symptoms last. do not take more than 6 tablets in 24 hours, unless directed by a doctor. do not use for more than 10 days unless directed by a doctor.
children under 12 years: ask a doctor.

Other information
store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature.
see end panel for lot number and expiration date.

Inactive ingredients FD&C red no. 40, FD&C yellow no. 6, lecithin, polyethylene glycol, polyvinyl alcohol, povidone k-30, pregelatinized starch, stearic acid, sucralose, talc, titanium dioxide.

Questions or comments?
call 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST

Extra Strength Pain Relief
ACETAMINOPHEN, 500 mg
Pain reliever/fever reducer
Contains no aspirin

50 Coated tablets

Extra Strength Pain Relief
ACETAMINOPHEN, 500 mg
Pain reliever/fever reducer
Contains no aspirin
For adults

50 Coated tablets

ACTUAL SIZE
Compare to the active ingredient in Extra Strength Tylenol®

ACTUAL SIZE
Compare to the active ingredient in Extra Strength Tylenol®

COATING FREE AREA

200000005004
700000003737

COATING FREE AREA

DEVELOPED BY PUBLIX SUPER MARKETS, INC.
3008 PUBLIX CORPORATE PARKWAY
LAKELAND, FL 33811 1-888-881-9877
PUBLIX SUPER MARKETS, INC. is a registered trademark of Publix Super Markets, Inc.
PUBLIX SUPER MARKETS, INC. is a registered trademark of Publix Super Markets, Inc.


Publix

0 4 14 15 55 173 2

Lot
Exp.

COATING FREE AREA

NDC 41415-538-05



ACETAMINOPHEN			
acetaminophen tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41415-538
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg
Inactive Ingredients			
Ingredient Name			Strength

STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE K30 (UNII: U725QWY32X)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	L;1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41415-538-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	09/09/2024	
2	NDC:41415-538-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/09/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/09/2024	

Labeler - PUBLIX SUPERMARKETS, INC (006922009)

Revised: 9/2024

PUBLIX SUPERMARKETS, INC