

**ACETAMINOPHEN- acetaminophen tablet**  
**PUBLIX SUPERMARKETS, INC**

-----  
**Extra Strength**  
**Pain Relief**

**ACETAMINOPHEN, 500 mg**

- **Pain reliever/fever reducer**
- **Contains no aspirin**
- **For adults**

**Active ingredient (in each coated tablet)**

Acetaminophen USP, 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold ■ headache
  - backache ■ minor pain of arthritis
  - toothache ■ muscular aches
  - premenstrual and menstrual cramps
  - temporarily reduces fever

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

if a skin reaction occurs, stop use and seek medical help right away

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

### **Ask a doctor before use if you have**

liver disease

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - redness or swelling is present
- These could be signs of a serious condition

### **If pregnant or breast-feeding**

ask a health professional before use

### **Keep out of reach of children**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

#### **■ do not take more than directed (see overdose warning)**


adults and children 12 years and over

- take 2 tablets every 6 hours while symptoms last
  - do not take more than 6 tablets in 24 hours, unless directed by a doctor
  - do not use for more than 10 days unless directed by a doctor
- children under 12 years

- ask a doctor

### **Other information**





# ACETAMINOPHEN

acetaminophen tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41415-538-05
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

## Inactive Ingredients

Ingredient Name	Strength

<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	L;1
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41415-538-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	09/09/2024	
2	NDC:41415-538-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/09/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/09/2024	

**Labeler** - PUBLIX SUPERMARKETS, INC (006922009)

Revised: 12/2025

PUBLIX SUPERMARKETS, INC