

ZILACTIN EARLY RELIEF COLD SORE- benzyl alcohol gel
Blairex Laboratories, Inc.

Zilactin Red Blairex

Drug Facts

Active ingredient

Benzyl alcohol 10%

Purpose

Cold sore/fever blister treatment, Oral pain reliever

Use

temporarily relieves pain caused by:

- cold sores/fever blisters
- canker sores, mouth sores
- gum irritations

Warnings

Flammable:

- keep away from fire or flame

Stop use and ask a physician if

- sore mouth symptoms do not improve in 7 days
- condition worsens or symptoms clear up and occur again within a few days
- swelling, rash or fever develops
- irritation, pain or redness persists or worsens
- apply only to affected area
- do not exceed recommended dosage
- avoid contact with the eyes
- do not use for more than 7 days unless directed by a physician or dentist

Keep out of reach of children. If swallowed, get medical help or contact a poison control center immediately.

Directions

adults and children 2 years and older	dry affected area. Apply with cotton swab or clean finger up to 4 times daily. Allow to dry 30-60 seconds.
children under 12	adult supervision should be given in the use of this product

years	
children under 2 years	do not use, consult a physician or dentist


Other information

- do not peel off protective film. Attempting to peel off film may result in skin irritation or tenderness. To remove film, first apply another coat of Zilactin to film, and immediately wipe the area with a moist gauze pad or tissue.
- contains alcohol 68% by volume
- store at 15-30°C (59-86°F)

Inactive ingredients

boric acid, hydroxypropylcellulose, propylene glycol, purified water, salicylic acid, SD alcohol 37, tannic acid

Manufactured for: Blairex Laboratories, Inc. P.O. Box 2127 Columbus, IN 47202 USA

 Questions or Comments?

Call TOLL FREE 1-800-252-4739

www.zilactin.com

PROMOTES HEALING

Cold Sore/Fever Blister/Oral Pain Treatment

Zilactin[®] Early Relief Cold Sore Gel

- Bioadhesive Promotes Healing
- Relieves Pain for up to 6 Hours

Net wt. 0.25 oz (7.1g)

PROMOTES
HEALING

Cold Sore/
Fever Blister/
Oral Pain Treatment

zilactin[®]

Early Relief Cold Sore Gel

• **Bioadhesive**
Promotes Healing

• **Relieves Pain**
for up to 6 Hours

Net wt. 0.25 oz (7.1g)

Dries Clear

www.zilactin.com

zilactin[®]

Early Relief Cold Sore Gel



000-S00027-03



Manufactured for:
Blair Laboratories, Inc.

P.O. Box 2127 Columbus, IN 47202 USA

Questions or Comments?

Call TOLL FREE 1-800-252-4739

www.zilactin.com

Tamper Evident Feature: Do not use if glued carton has been opened or tampered with.

Inactive ingredients boric acid, hydroxypropylcellulose, propylene glycol, purified water, salicylic acid, SD alcohol 37, tannic acid

Other information ■ do not peel off protective film. Attempting to peel off film may result in skin irritation or tenderness. To remove film, first apply another coat of Zilactin to film, and immediately wipe the area with a moist gauze pad or tissue. ■ contains alcohol 68% by volume ■ store at 15-30°C (59-86°F)


adults and children 2 years and older	dry affected area. Apply with cotton swab or clean finger up to 4 times daily. Allow to dry 30-60 seconds.
children under 12 years	adult supervision should be given in the use of this product
children under 2 years	do not use, consult a physician or dentist

Directions

Keep out of reach of children. If swallowed, get medical help or contact a poison control center immediately.

■ apply only to affected area ■ do not exceed recommended dosage ■ avoid contact with the eyes ■ do not use for more than 7 days unless directed by a physician or dentist

develops ■ irritation, pain or redness persists or worsens



ZILACTIN EARLY RELIEF COLD SORE

benzyl alcohol gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50486-468
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH) (BENZYL ALCOHOL - UNII:LKG8494WBH)	BENZYL ALCOHOL	0.1 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
THYMOL (UNII: 3J50XA376E)	
MENTHOL (UNII: L7T10EIP3A)	
TANNIC ACID (UNII: 28F9E0DJY6)	
WATER (UNII: 059QF0KO0R)	
BORIC ACID (UNII: R57ZHV85D4)	
HYDROXYPROPYL CELLULOSE (TYPE M) (UNII: U3JF91U133)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
ALCOHOL (UNII: 3K9958V90M)	
EUCALYPTOL (UNII: RV6J6604TK)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50486-468-02	1 in 1 CARTON	06/30/2005	
1		7.1 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	06/30/2005	

Labeler - Blairex Laboratories, Inc. (092575133)

Revised: 11/2024

Blairex Laboratories, Inc.